

SPECIALIZATION AND MICROLOCALIZATION OF SUBANALYTIC SHEAVES

Luca Prelli

Abstract

In this paper we define the specialization and microlocalization functors for subanalytic sheaves. Then we specialize and microlocalize the sheaves of tempered and Whitney holomorphic functions generalizing some classical constructions.

Contents

Introduction	2
1 Review on sheaves on subanalytic sites	5
1.1 Sheaves on subanalytic sites	5
1.2 Modules over a $k_{X_{sa}}$ -algebra	8
2 Conic sheaves on subanalytic sites	9
2.1 Conic sheaves on topological spaces	9
2.2 Conic sheaves on subanalytic sites	11
2.3 An equivalence of categories	14
2.4 Derived category	17
2.5 Operations	21
3 Fourier-Sato transform for subanalytic sheaves.	26
3.1 Conic sheaves on vector bundles	26
3.2 Fourier-Sato transformation	29
4 Specialization of subanalytic sheaves	34
4.1 Review on normal deformation	34
4.2 Specialization of subanalytic sheaves	37

5	Microlocalization of subanalytic sheaves	41
5.1	Microlocalization of subanalytic sheaves	41
5.2	The functor μhom^{sa}	44
5.3	Microlocalization and microsupport	45
5.4	The link with the functor μ of microlocalization	48
6	Holomorphic functions with growth conditions	52
6.1	Review on temperate and formal cohomology	52
6.2	Tempered and Whitney holomorphic functions	53
6.3	Asymptotic expansions	54
6.4	Microlocalization of $\mathcal{O}_X^t$ and $\mathcal{O}_X^w$	59
7	Integral transforms	61
7.1	$\mathcal{E}_X$ -modules	62
7.2	Integral transforms	62
7.3	Microlocal integral transformations	75
7.4	Contact transformations	76
8	Cauchy-Kowaleskaya-Kashiwara theorem	78
8.1	Microsupport and characteristic variety	79
8.2	Cauchy-Kowaleskaya-Kashiwara theorem	82
A	Appendix	84
A.1	Review on subanalytic sets	84
A.2	Ind-sheaves and subanalytic sites	87
A.3	Inverse image for tempered holomorphic functions	89
A.4	Inverse image for Whitney holomorphic functions	93
	Bibliography	96

Introduction

After the fundamental works of Sato on hyperfunctions and microfunctions and the developement of algebraic analysis, the methods of cohomological theory of sheaves became very useful for studying systems of PDE on real or complex analytic manifolds. Unfortunately sheaf theory is not well suited to study objects which are not defined by local properties. Since the study of solutions of a system of PDE in these spaces is very important (Riemann-Hilbert correspondence, Laplace transform, etc.), many ways have been explored by the specialists to overcome this problem. First Kashiwara in [9]

defined the functor of tempered cohomology to solve the Riemann-Hilbert problem for holonomic systems. Its dual, the functor of formal cohomology was introduced by Kashiwara and Schapira in [13]. Their microlocal analogues were defined by Andronikof in [1] and Colin in [5]. These results present a problem: they are construction “ad hoc”, and they are not contained in a unifying theory based on the six Grothendieck operations. In [14] Kashiwara and Schapira introduced the notion of ind-sheaves, and defined the six Grothendieck operations in this framework. Then they defined the subanalytic site (a site whose open sets are subanalytic and the coverings are locally finite) and they proved the equivalence between subanalytic sheaves and ind- $\mathbb{R}$ -constructible sheaves. They obtained the formalism of the six Grothendieck operations by including subanalytic sheaves into the category of ind-sheaves. In [23] a direct, self-contained and elementary construction of the six Grothendieck operations for subanalytic sheaves is introduced without using the more sophisticated and much more difficult theory of ind-sheaves. In the beginning of this paper we extend some classical constructions for sheaves, as the Fourier-Sato transform and the functors of specialization and microlocalization. We introduce first the category of conic subanalytic sheaves on an analytic manifold. Then we extend the Fourier-Sato transform to the category of conic subanalytic sheaves on a vector bundle. At this point we can start studying subanalytic sheaves from a microlocal point of view by introducing the functors of specialization and microlocalization along a submanifold of a real analytic manifold. Roughly speaking, starting from a sheaf on a real analytic manifold, we construct conic sheaves on the tangent and the cotangent bundle respectively. We also show that the functor of microlocalization is related with the functor of ind-microlocalization defined in [16]. Then, applying specialization (resp. microlocalization) to the subanalytic sheaves of tempered and Whitney holomorphic functions, we generalize tempered and formal specialization (resp. microlocalization). In this way we get a unifying description of Andronikov’s and Colin’s “ad hoc” constructions. This is also a “simplification”, since the definitions of specialization and microlocalization for subanalytic sheaves are more intuitive for people which are familiar with the classical sheaf theory of [11], although working with the subanalytic site is delicate and requires results which are related to the geometry of subanalytic subsets.

In more details the contents of this work are as follows.

In **Section 1** we recall the results on subanalytic sheaves of [14] and [23].

In **Section 2** we construct the category of conic sheaves on a subanalytic site endowed with an action of $\mathbb{R}^+$.

In **Section 3** we consider a vector bundle E over a real analytic manifold and its dual E^* endowed with the natural action of $\mathbb{R}^+$. We define the Fourier-Sato transform which gives an equivalence between conic subanalytic sheaves on E and conic subanalytic sheaves on E^* .

Then we define the functor ν_M^{sa} of specialization along a submanifold M of a real analytic manifold X (**Section 4**) and its Fourier-Sato transform, the functor μ_M^{sa} of microlocalization (**Section 5**). We introduce the functor μhom^{sa} for subanalytic sheaves and we give an estimate of its support using the notion of microsupport of [15]. Then we study its relation with the functor of ind-microlocalization of [16].

We apply these results in **Section 6**. We study the connection between specialization and microlocalization for subanalytic sheaves and the classical ones. Specialization of subanalytic sheaves generalize tempered and formal specialization of [1] and [6], in particular when we specialize Whitney holomorphic functions we obtain the sheaves of functions asymptotically developable of [20] and [29]. Moreover, thanks to the functor of microlocalization, we are able to generalize tempered and formal microlocalization introduced by Andronikof in [1] and Colin in [5] respectively.

Section 7 is dedicated to the study of the microlocalization of tempered and Whitney holomorphic functions. We prove that the microlocalization of $\mathcal{O}^t$ and $\mathcal{O}^w$ have (in cohomology) a natural structure of $\mathcal{E}$ -module and that locally they are invariant under contact transformations.

In **Section 8** we study the Cauchy-Kowaleskaya-Kashiwara theorem with growth conditions. The proof is not a rephrasing of Kashiwara's proof, which consisted in a reduction to the case of one operator: we first prove a propagation result (already known for $\mathcal{O}_X^t$) and we use this propagation result to reduce the statements to the commutation of duality with non-characteristic inverse image.

We end this work with a short **Appendix** in which we recall the definitions and we collect some properties of subanalytic subsets and ind-sheaves, and then we study the inverse image of the subanalytic sheaves of tempered and Whitney holomorphic functions.

Acknowledgments. We would like to thank Prof. Pierre Schapira who encouraged us to develop specialization and microlocalization of subanalytic sheaves, and Prof. A. D'Agnolo for his many useful remarks.

1 Review on sheaves on subanalytic sites

In the following X will be a real analytic manifold and k a field. Reference are made to [12] for a complete exposition on sheaves on Grothendieck topologies, to [14] and [23] for an introduction to sheaves on subanalytic sites. We refer to [3] and [18] for the theory of subanalytic sets.

1.1 Sheaves on subanalytic sites

Let us recall some results of [14] and [23].

Denote by $\text{Op}(X_{sa})$ the category of open subanalytic subsets of X . One endows $\text{Op}(X_{sa})$ with the following topology: $S \subset \text{Op}(X_{sa})$ is a covering of $U \in \text{Op}(X_{sa})$ if for any compact K of X there exists a finite subset $S_0 \subset S$ such that $K \cap \bigcup_{V \in S_0} V = K \cap U$. We will call X_{sa} the subanalytic site, and for $U \in \text{Op}(X_{sa})$ we denote by $U_{X_{sa}}$ the category $\text{Op}(X_{sa}) \cap U$ with the topology induced by X_{sa} . We denote by $\text{Op}^c(X_{sa})$ the subcategory of $\text{Op}(X_{sa})$ consisting of relatively compact open subanalytic subsets.

Let $\text{Mod}(k_{X_{sa}})$ denote the category of sheaves on X_{sa} . Then $\text{Mod}(k_{X_{sa}})$ is a Grothendieck category, i.e. it admits a generator and small inductive limits, and small filtrant inductive limits are exact. In particular as a Grothendieck category, $\text{Mod}(k_{X_{sa}})$ has enough injective objects.

Let $\text{Mod}_{\mathbb{R}\text{-c}}(k_X)$ be the abelian category of $\mathbb{R}$ -constructible sheaves on X , and consider its subcategory $\text{Mod}_{\mathbb{R}\text{-c}}^c(k_X)$ consisting of sheaves whose support is compact.

We denote by $\rho : X \rightarrow X_{sa}$ the natural morphism of sites. We have functors

$$\text{Mod}(k_X) \begin{array}{c} \xrightarrow{\rho_*} \\ \xleftarrow[\rho_!]{\rho^{-1}} \end{array} \text{Mod}(k_{X_{sa}}).$$

The functors ρ^{-1} and ρ_* are the functors of inverse image and direct image respectively. The sheaf $\rho_! F$ is the sheaf associated to the presheaf $\text{Op}(X_{sa}) \ni$

$U \mapsto F(\overline{U})$. In particular, for $U \in \text{Op}(X)$ one has $\rho_! k_U \simeq \varinjlim_{V \subset \subset U} \rho_* k_V$, where $V \in \text{Op}(X_{sa})$. Let us summarize the properties of these functors:

- the functor ρ_* is fully faithful and left exact, the restriction of ρ_* to $\text{Mod}_{\mathbb{R}\text{-c}}(k_X)$ is exact,
- the functor ρ^{-1} is exact,
- the functor $\rho_!$ is fully faithful and exact,
- (ρ^{-1}, ρ_*) and $(\rho_!, \rho^{-1})$ are pairs of adjoint functors.

Notations 1.1.1 *Since the functor ρ_* is fully faithful and exact on $\text{Mod}_{\mathbb{R}\text{-c}}(k_X)$, we can identify $\text{Mod}_{\mathbb{R}\text{-c}}(k_X)$ with its image in $\text{Mod}(k_{X_{sa}})$. When there is no risk of confusion we will write F instead of $\rho_* F$, for $F \in \text{Mod}_{\mathbb{R}\text{-c}}(k_X)$.*

Let $F \in \text{Mod}(k_{X_{sa}})$. There exists a filtrant inductive system $\{F_i\}_{i \in I}$ in $\text{Mod}_{\mathbb{R}\text{-c}}^c(k_X)$ such that $F \simeq \varinjlim_i \rho_* F_i$.

Let X, Y be two real analytic manifolds, and let $f : X \rightarrow Y$ be a real analytic map. We have a commutative diagram

$$(1.1) \quad \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow \rho & & \downarrow \rho \\ X_{sa} & \xrightarrow{f} & Y_{sa} \end{array}$$

We get external operations f^{-1} and f_* , which are always defined for sheaves on Grothendieck topologies. For subanalytic sheaves we can also define the functor of proper direct image

$$\begin{aligned} f_{!!} : \text{Mod}(k_{X_{sa}}) &\rightarrow \text{Mod}(k_{Y_{sa}}) \\ F &\mapsto \varinjlim_U f_* F_U \simeq \varinjlim_K f_* \Gamma_K F \end{aligned}$$

where U ranges through the family of relatively compact open subanalytic subsets of X and K ranges through the family of subanalytic compact subsets of X . The notation $f_{!!}$ follows from the fact that $f_{!!} \circ \rho_* \not\simeq \rho_* \circ f_!$ in general. If f is proper on $\text{supp}(F)$ then $f_* F \simeq f_{!!} F$, in this case $f_{!!}$ commutes with ρ_* . While functors f^{-1} and $\otimes$ are exact, the functors $\mathcal{H}om$, f_* and $f_{!!}$ are left exact and admits right derived functors.

To derive these functors we use the category of quasi-injective objects. An object $F \in \text{Mod}(k_{X_{sa}})$ is quasi-injective if for $U, V \in \text{Op}^c(X_{sa})$ with $V \subset U$ the restriction morphism $\Gamma(U; F) \rightarrow \Gamma(V; F)$ is surjective or, equivalently, if the functor $\text{Hom}_{k_{X_{sa}}}(\cdot, F)$ is exact on $\text{Mod}_{\mathbb{R}\text{-c}}^c(k_X)$. Quasi injective objects are injective with respect to the functors f_* , $f_{!!}$ and, if $G \in \text{Mod}_{\mathbb{R}\text{-c}}(k_X)$, with respect to the functors $\text{Hom}_{k_{X_{sa}}}(G, \cdot)$, $\mathcal{H}om(G, \cdot)$.

The functor $Rf_{!!}$ admits a right adjoint, denoted by $f^!$, and we get the usual isomorphisms between Grothendieck operations (projection formula, base change formula, Künneth formula, etc.) in the framework of subanalytic sites.

Let Z be a subanalytic locally closed subset of X . As in classical sheaf theory we define

$$\begin{aligned} \Gamma_Z : \text{Mod}(k_{X_{sa}}) &\rightarrow \text{Mod}(k_{X_{sa}}) \\ F &\mapsto \mathcal{H}om(\rho_* k_Z, F) \\ (\cdot)_Z : \text{Mod}(k_{X_{sa}}) &\rightarrow \text{Mod}(k_{X_{sa}}) \\ F &\mapsto F \otimes \rho_* k_Z. \end{aligned}$$

Finally we recall the properties of the six Grothendieck operations and their relations with the functors ρ^{-1} , $R\rho_*$ and $\rho_!$. We refer to [23] for a detailed exposition.

- the functor $R^k \mathcal{H}om(\rho_* F, \cdot)$ commutes with filtrant $\varinjlim$ if $F \in \text{Mod}_{\mathbb{R}\text{-c}}(k_X)$,
- the functors $R^k f_{!!}$ and $H^k f^!$ commute with filtrant $\varinjlim$,
- the functor ρ^{-1} commutes with $\otimes$, f^{-1} and $Rf_{!!}$,
- the functor $R\rho_*$ commutes with $R\mathcal{H}om$, Rf_* and $f^!$,
- the functor $\rho_!$ commutes with $\otimes$ and f^{-1} ,
- the restriction of $\otimes$ and f^{-1} to the category of $\mathbb{R}$ -constructible sheaves commute with ρ_* ,
- if f is a topological submersion (i.e. it is locally isomorphic to a projection $Y \times \mathbb{R}^n \rightarrow Y$), then $f^! \simeq f^{-1} \otimes f^! k_Y$ commutes with ρ^{-1} and $Rf_{!!}$ commutes with $\rho_!$.

Moreover the functors Rf_* , $Rf_{!!}$ and $R\mathcal{H}om(F, \cdot)$ with $F \in \text{Mod}_{\mathbb{R}\text{-c}}(k_X)$ have finite cohomological dimension.

1.2 Modules over a $k_{X_{sa}}$ -algebra

A sheaf of $k_{X_{sa}}$ -algebras (or a $k_{X_{sa}}$ -algebra, for short) is an object $\mathcal{R} \in \text{Mod}(k_{X_{sa}})$ such that $\Gamma(U; \mathcal{R})$ is a k -algebra for each $U \in \text{Op}(X_{sa})$ and the restriction maps are algebra morphisms. A sheaf of (left) $\mathcal{R}$ -modules is a sheaf F such that $\Gamma(U; F)$ has a structure of (left) $\Gamma(U; \mathcal{R})$ -module for each $U \in \text{Op}(X_{sa})$. Let us denote by $\text{Mod}(\mathcal{R})$ the category of sheaves of (left) $\mathcal{R}$ -modules. The category $\text{Mod}(\mathcal{R})$ is a Grothendieck category and the forgetful functor $for : \text{Mod}(\mathcal{R}) \rightarrow \text{Mod}(k_{X_{sa}})$ is exact.

The functors

$$\begin{aligned} \mathcal{H}om_{\mathcal{R}} & : \text{Mod}(\mathcal{R})^{op} \times \text{Mod}(\mathcal{R}) \rightarrow \text{Mod}(k_{X_{sa}}), \\ \otimes_{\mathcal{R}} & : \text{Mod}(\mathcal{R}^{op}) \times \text{Mod}(\mathcal{R}) \rightarrow \text{Mod}(k_{X_{sa}}) \end{aligned}$$

are well defined. Remark that in the case of $\mathcal{R}$ -modules the functor $\otimes_{\mathcal{R}}$ is only right exact and commutes with $\varinjlim$.

Let X, Y be two real analytic manifolds, and let $f : X \rightarrow Y$ be a morphism of real analytic manifolds. Let $\mathcal{R}$ be a $k_{Y_{sa}}$ -algebra. The functors f^{-1} , Rf_* and $Rf_{!!}$ induce functors

$$\begin{aligned} f^{-1} & : \text{Mod}(\mathcal{R}) \rightarrow \text{Mod}(f^{-1}\mathcal{R}), \\ f_* & : \text{Mod}(f^{-1}\mathcal{R}) \rightarrow \text{Mod}(\mathcal{R}), \\ f_{!!} & : \text{Mod}(f^{-1}\mathcal{R}) \rightarrow \text{Mod}(\mathcal{R}). \end{aligned}$$

Now we consider the derived category of sheaves of $\mathcal{R}$ -modules.

Definition 1.2.1 *An object $F \in \text{Mod}(\mathcal{R})$ is flat if the functor $\text{Mod}(\mathcal{R}^{op}) \ni G \rightarrow G \otimes_{\mathcal{R}} F$ is exact.*

Thanks to flat objects we can find a left derived functor $\otimes_{\mathcal{R}}^L$ of the tensor product $\otimes_{\mathcal{R}}$.

Definition 1.2.2 *An object $F \in \text{Mod}(\mathcal{R})$ is quasi-injective if its image via the forgetful functor is quasi-injective in $\text{Mod}(k_{X_{sa}})$.*

Let X, Y be two real analytic manifolds, and let $f : X \rightarrow Y$ be a real analytic map. Let $\mathcal{R}$ be a $k_{Y_{sa}}$ -algebra. One can prove that quasi-injective objects are injective with respect to the functors f_* and $f_{!!}$. The functors Rf_* and $Rf_{!!}$ are well defined and projection formula, base change formula remain valid for $\mathcal{R}$ -modules. Moreover we have

Theorem 1.2.3 *The functor $Rf_{!!} : D^+(f^{-1}\mathcal{R}) \rightarrow D^+(\mathcal{R})$ admits a right adjoint. We denote by $f^! : D^+(\mathcal{R}) \rightarrow D^+(f^{-1}\mathcal{R})$ the adjoint functor.*

2 Conic sheaves on subanalytic sites

In this section we study the category of conic sheaves on a subanalytic site. Reference are made to [11] for the classical theory of conic sheaves.

2.1 Conic sheaves on topological spaces

Let k be a field. Let X be a real analytic manifold endowed with a subanalytic action μ of $\mathbb{R}^+$. In other words we have a subanalytic map

$$\mu : X \times \mathbb{R}^+ \rightarrow X,$$

which satisfies, for each $t_1, t_2 \in \mathbb{R}^+$:

$$\begin{cases} \mu(x, t_1 t_2) = \mu(\mu(x, t_1), t_2), \\ \mu(x, 1) = x. \end{cases}$$

Note that μ is open. Indeed let $U \in \text{Op}(X)$ and $W \in \text{Op}(\mathbb{R}^+)$. Then $\mu(U, W) = \bigcup_{t \in W} \mu(U, t)$, and $\mu(\cdot, t) : X \rightarrow X$ is a homeomorphism (with inverse $\mu(\cdot, t^{-1})$). We have a diagram

$$X \xrightarrow{j} X \times \mathbb{R}^+ \xrightleftharpoons[p]{\mu} X,$$

where $j(x) = (x, 1)$ and p denotes the projection. We have $\mu \circ j = p \circ j = \text{id}$.

Definition 2.1.1 (i) Let S be a subset of X . We set $\mathbb{R}^+ S = \mu(S, \mathbb{R}^+)$. If $U \in \text{Op}(X)$, then $\mathbb{R}^+ U \in \text{Op}(X)$ since μ is open.

(ii) Let S be a subset of X . We say that S is conic if $S = \mathbb{R}^+ S$. In other words, S is invariant by the action of μ .

(iii) An orbit of μ is the set $\mathbb{R}^+ x$ with $x \in X$.

Definition 2.1.2 We say that a subset S of X is $\mathbb{R}^+$ -connected if $S \cap \mathbb{R}^+ x$ is connected for each $x \in S$.

Definition 2.1.3 A sheaf $F \in \text{Mod}(k_X)$ is conic if $\mu^{-1} F \simeq p^{-1} F$.

(i) We denote by $\text{Mod}_{\mathbb{R}^+}(k_X)$ the subcategory of $\text{Mod}(k_X)$ consisting of conic sheaves.

(ii) We denote by $D_{\mathbb{R}^+}^b(k_{X_{sa}})$, the subcategory of $D^b(k_X)$ consisting of objects F such that $H^j(F)$ belongs to $\text{Mod}_{\mathbb{R}^+}(k_X)$ for all $j \in \mathbb{Z}$.

Proposition 2.1.4 *Let $U \in \text{Op}(X)$ be $\mathbb{R}^+$ -connected and let $F \in D_{\mathbb{R}^+}^b(k_X)$. Then*

$$\text{R}\Gamma(\mathbb{R}^+U; F) \xrightarrow{\sim} \text{R}\Gamma(U; F).$$

Let us assume the following hypothesis

$$(2.1) \quad \begin{cases} \text{(i) every point } x \in X \text{ has a fundamental neighborhood} \\ \text{system consisting of } \mathbb{R}^+\text{-connected open subsets;} \\ \text{(ii) for any } x \in X \text{ the set } \mathbb{R}^+x \text{ is contractible.} \end{cases}$$

In this situation (see [2]) either $\mathbb{R}^+x \simeq \mathbb{R}$ or $\mathbb{R}^+x = x$.

Denote by $X_{\mathbb{R}^+}$ the topological space X endowed with the conic topology, i.e. $U \in \text{Op}(X_{\mathbb{R}^+})$ if it is open for the topology of X and invariant by the action of $\mathbb{R}^+$.

Let us consider the natural map $\eta : X \rightarrow X_{\mathbb{R}^+}$. The restriction of η_* induces an exact functor denoted by $\tilde{\eta}_*$ and we obtain a diagram

$$(2.2) \quad \begin{array}{ccc} \text{Mod}_{\mathbb{R}^+}(k_X) & \xrightleftharpoons[\eta^{-1}]{\tilde{\eta}_*} & \text{Mod}(k_{X_{\mathbb{R}^+}}) \\ \downarrow & \nearrow \eta_* & \\ \text{Mod}(k_X) & & \end{array}$$

Let $F \in D_{\mathbb{R}^+}^b(k_X)$. Let φ be the natural map from $\text{R}\Gamma(\mathbb{R}^+U; F)$ to $\text{R}\Gamma(U; \eta^{-1}F)$ defined by

$$(2.3) \quad \begin{array}{ccc} \text{R}\Gamma(\mathbb{R}^+U; F) & \rightarrow & \text{R}\Gamma(\mathbb{R}^+U; R\eta_*\eta^{-1}F) \\ & \simeq & \text{R}\Gamma(\mathbb{R}^+U; \eta^{-1}F) \\ & \rightarrow & \text{R}\Gamma(U; \eta^{-1}F). \end{array}$$

Proposition 2.1.5 *Let F be a sheaf over $X_{\mathbb{R}^+}$. Let U be an open set of X and assume that U is $\mathbb{R}^+$ -connected. Then the morphism φ defined by (2.3) is an isomorphism.*

Theorem 2.1.6 *The functors $R\eta_*$ and η^{-1} in (2.2) induce equivalences of derived categories*

$$D_{\mathbb{R}^+}^b(k_X) \xrightleftharpoons[\eta^{-1}]{R\eta_*} D^b(k_{X_{\mathbb{R}^+}})$$

inverse to each others.

We need to introduce the subcategory of coherent conic sheaves.

Definition 2.1.7 Let $U \in \text{Op}(X_{\mathbb{R}^+})$. Then U is said to be relatively quasi-compact if, for any covering $\{U_i\}_{i \in I}$ of $X_{\mathbb{R}^+}$, there exists $J \subset I$ finite such that $U \subset \bigcup_{i \in J} U_i$. We write $U \subset\subset X_{\mathbb{R}^+}$.

We will denote by $\text{Op}^c(X_{\mathbb{R}^+})$ the subcategory of $\text{Op}(X_{\mathbb{R}^+})$ consisting of relatively quasi-compact open subsets.

One can check easily that if $U \in \text{Op}^c(X)$, then $\mathbb{R}^+U \in \text{Op}^c(X_{\mathbb{R}^+})$.

Definition 2.1.8 Let $F \in \text{Mod}(k_{X_{\mathbb{R}^+}})$ and consider the family $\text{Op}(X_{sa, \mathbb{R}^+})$.

- (i) F is $X_{sa, \mathbb{R}^+}$ -finite if there exists an epimorphism $G \twoheadrightarrow F$, with $G \simeq \bigoplus_{i \in I} k_{U_i}$, I finite and $U_i \in \text{Op}^c(X_{sa, \mathbb{R}^+})$.
- (ii) F is $X_{sa, \mathbb{R}^+}$ -pseudo-coherent if for any morphism $\psi : G \rightarrow F$, where G is $X_{sa, \mathbb{R}^+}$ -finite, $\ker \psi$ is $X_{sa, \mathbb{R}^+}$ -finite.
- (iii) F is $X_{sa, \mathbb{R}^+}$ -coherent if it is both $X_{sa, \mathbb{R}^+}$ -finite and $X_{sa, \mathbb{R}^+}$ -pseudo-coherent.

We will denote by $\text{Coh}(X_{sa, \mathbb{R}^+})$ the subcategory of $\text{Mod}(k_{X_{\mathbb{R}^+}})$ consisting of $X_{sa, \mathbb{R}^+}$ -coherent objects.

2.2 Conic sheaves on subanalytic sites

Definition 2.2.1 A sheaf of k -modules F on X_{sa} is conic if the restriction morphism $\Gamma(\mathbb{R}^+U; F) \rightarrow \Gamma(U; F)$ is an isomorphism for each $\mathbb{R}^+$ -connected $U \in \text{Op}^c(X_{sa})$ with $\mathbb{R}^+U \in \text{Op}(X_{sa})$.

- (i) We denote by $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ the subcategory of $\text{Mod}(k_{X_{sa}})$ consisting of conic sheaves.
- (ii) We denote by $D_{\mathbb{R}^+}^b(k_{X_{sa}})$, the subcategory of $D^b(k_{X_{sa}})$ consisting of objects F such that $H^j(F)$ belongs to $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ for all $j \in \mathbb{Z}$.

Remark 2.2.2 Let X be a real analytic manifold endowed with a subanalytic action μ of $\mathbb{R}^+$ and consider the following diagram

$$X \times \mathbb{R}^+ \xrightleftharpoons[p]{\mu} X,$$

where p denotes the projection. As in classical sheaf theory one can define the subcategory $\text{Mod}^\mu(k_{X_{sa}})$ of $\text{Mod}(k_{X_{sa}})$ consisting of sheaves satisfying

$\mu^{-1}F \simeq p^{-1}F$. The categories $\text{Mod}^\mu(k_{X_{sa}})$ and $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ are not equivalent in general.

Indeed, let $X = \mathbb{R}$, set $X^+ = \{x \in \mathbb{R}; x > 0\}$ and let μ be the natural action of $\mathbb{R}^+$ (i.e. $\mu(x, t) = tx$). Let us consider the sheaf $\rho_!k_{X^+} \in \text{Mod}(k_{X_{sa}})$. Then

$$\mu^{-1}\rho_!k_{X^+} \simeq \rho_!\mu^{-1}k_{X^+} \simeq \rho_!p^{-1}k_{X^+} \simeq p^{-1}\rho_!k_{X^+}.$$

Let $V = \{x \in \mathbb{R}; 1 < x < 2\}$ and set $W_m = \{x \in \mathbb{R}; \frac{1}{m} < x < m\}$, where $m \in \mathbb{N} \setminus \{0\}$. Recall that $\rho_!k_{X^+} \simeq \varinjlim_{U \subset \subset X^+} \rho_*k_U \simeq \varinjlim_m \rho_*k_{W_m}$. We have

$$\Gamma(V; \rho_!k_{X^+}) \simeq \varinjlim_m \Gamma(V; k_{W_m}) \simeq k,$$

since $V \subset W_m$ for $m \geq 2$. On the other hand, let $V_n^+ = \{x \in \mathbb{R}; 0 < x < n\}$, where $n \in \mathbb{N}$. Since $\mathbb{R}^+V = X^+$ we have

$$\Gamma(X^+; \rho_!k_{X^+}) \simeq \varprojlim_n \Gamma(V_n^+; \rho_!k_{X^+}) \simeq \varprojlim_n \varinjlim_m \Gamma(V_n^+; k_{W_m}),$$

(in the second isomorphism we used the fact that $V_n^+ \in \text{Op}^c(X_{sa})$ for each n) and $\Gamma(V_n^+; k_{W_m}) = 0$ for each $n, m \in \mathbb{N}$. Hence $\Gamma(V; \rho_!k_{X^+}) \not\simeq \Gamma(\mathbb{R}^+V; \rho_!k_{X^+})$.

Definition 2.2.3 We denote by $\text{Op}(X_{sa, \mathbb{R}^+})$ the full subcategory of $\text{Op}(X_{sa})$ consisting of conic subanalytic subsets, i.e. $U \in \text{Op}(X_{sa, \mathbb{R}^+})$ if $U \in \text{Op}(X_{sa})$ and it is invariant by the action of $\mathbb{R}^+$.

We denote by $X_{sa, \mathbb{R}^+}$ the category $\text{Op}(X_{sa, \mathbb{R}^+})$ endowed with the topology induced by X_{sa} .

We denote by $\rho_{\mathbb{R}^+} : X_{\mathbb{R}^+} \rightarrow X_{sa, \mathbb{R}^+}$ the natural morphism of sites.

Let $\eta : X \rightarrow X_{\mathbb{R}^+}$ and $\eta_{sa} : X_{sa} \rightarrow X_{sa, \mathbb{R}^+}$ be the natural morphisms of sites. We have a commutative diagram of sites

$$(2.4) \quad \begin{array}{ccc} X & \xrightarrow{\rho} & X_{sa} \\ \downarrow \eta & & \downarrow \eta_{sa} \\ X_{\mathbb{R}^+} & \xrightarrow{\rho_{\mathbb{R}^+}} & X_{sa, \mathbb{R}^+}. \end{array}$$

Lemma 2.2.4 Let $F \in \text{Coh}(X_{sa, \mathbb{R}^+})$. Then $\eta_{sa}^{-1}\rho_{\mathbb{R}^+*}F \simeq \rho_*\eta^{-1}F$.

Proof. Since all these functors are exact on $\text{Coh}(X_{sa, \mathbb{R}^+})$, we may reduce to the case $F = k_U$ with $U \in \text{Op}^c(X_{sa, \mathbb{R}^+})$. Then we have

$$\eta_{sa}^{-1}\rho_{\mathbb{R}^+*}k_U \simeq \eta_{sa}^{-1}k_U \simeq k_U,$$

on the other hand we have

$$\rho_* \eta^{-1} k_U \simeq \rho_* k_U \simeq k_U$$

and the result follows. $\square$

Replacing $\mathrm{Op}^c(X_{sa})$ with $\mathrm{Op}^c(X_{sa, \mathbb{R}^+})$, we can adapt the results of [14], [23] and we get the following results.

Theorem 2.2.5 (i) Let $G \in \mathrm{Coh}(X_{sa, \mathbb{R}^+})$ and let $\{F_i\}$ be a filtrant inductive system in $\mathrm{Mod}(k_{X_{sa, \mathbb{R}^+}})$. Then we have an isomorphism

$$\varinjlim_i \mathrm{Hom}_{k_{X_{sa, \mathbb{R}^+}}}(\rho_{\mathbb{R}^+*} G, F_i) \xrightarrow{\sim} \mathrm{Hom}_{k_{X_{sa, \mathbb{R}^+}}}(\rho_{\mathbb{R}^+*} G, \varinjlim_i F_i).$$

Moreover the functor of direct image $\rho_{\mathbb{R}^+*}$ associated to the morphism $\rho_{\mathbb{R}^+}$ in (2.4) is fully faithful and exact on $\mathrm{Coh}(X_{sa, \mathbb{R}^+})$.

(ii) Let $F \in \mathrm{Mod}(k_{X_{sa, \mathbb{R}^+}})$. There exists a small filtrant inductive system $\{F_i\}_{i \in I}$ in $\mathrm{Coh}(X_{sa, \mathbb{R}^+})$ such that $F \simeq \varinjlim_i \rho_{\mathbb{R}^+*} F_i$.

Notations 2.2.6 Since $\rho_{\mathbb{R}^+*}$ is fully faithful and exact on $\mathrm{Coh}(X_{sa, \mathbb{R}^+})$, we can identify $\mathrm{Coh}(X_{sa, \mathbb{R}^+})$ with its image in $\mathrm{Mod}(k_{X_{sa, \mathbb{R}^+}})$. When there is no risk of confusion we will write F instead of $\rho_{\mathbb{R}^+*} F$, for $F \in \mathrm{Coh}(X_{sa, \mathbb{R}^+})$.

We can also find a left adjoint to the functor $\rho_{\mathbb{R}^+}^{-1}$.

Proposition 2.2.7 The functor $\rho_{\mathbb{R}^+}^{-1}$ admits a left adjoint, denoted by $\rho_{\mathbb{R}^+!}$. It satisfies

(i) for $F \in \mathrm{Mod}(k_{X_{\mathbb{R}^+}})$ and $U \in \mathrm{Op}^c(X_{sa, \mathbb{R}^+})$, $\rho_{\mathbb{R}^+!} F$ is the sheaf associated to the presheaf $U \mapsto \varinjlim_{V \supset \supset U} \Gamma(V; F)$,

(ii) for $U \in \mathrm{Op}(X_{\mathbb{R}^+})$ one has $\rho_{\mathbb{R}^+!} k_U \simeq \varinjlim_{V \subset \subset U, V \in \mathrm{Op}^c(X_{sa, \mathbb{R}^+})} k_V$.

Remark 2.2.8 One can check that $\eta_{sa}^{-1} \circ \rho_{\mathbb{R}^+*} \simeq \rho_* \circ \eta^{-1}$ and $\rho^{-1} \circ \eta_{sa}^{-1} \simeq \eta^{-1} \circ \rho_{\mathbb{R}^+}^{-1}$. Remark that $\rho_! \circ \eta^{-1} \not\simeq \eta_{sa}^{-1} \circ \rho_{\mathbb{R}^+!}$. In fact with the notations of Remark 2.2.2 we have $\rho_! \eta^{-1} k_{X^+} \simeq \varinjlim_m \rho_* k_{W_m}$. On the other hand,

$$\eta_{sa}^{-1} \rho_{\mathbb{R}^+!} k_{X^+} \simeq k_{X^+}.$$

Let us extend the notion of quasi-injective object of [14], [23] to $\text{Mod}(k_{X_{sa, \mathbb{R}^+}})$.

Definition 2.2.9 *An object $F \in \text{Mod}(k_{X_{sa, \mathbb{R}^+}})$ is quasi-injective if the functor $\text{Hom}_{k_{X_{sa, \mathbb{R}^+}}}(\cdot, F)$ is exact in $\text{Coh}(X_{sa, \mathbb{R}^+})$ or, equivalently (see Theorem 8.7.2 of [12]) if for each $U, V \in \text{Op}^c(X_{sa, \mathbb{R}^+})$ with $V \subset U$ the restriction morphism $\Gamma(U; F) \rightarrow \Gamma(V; F)$ is surjective.*

The category of quasi-injective object is cogenerating since it contains injective objects. Moreover it is stable by filtrant $\varinjlim$ and $\prod$. We have the following result

Theorem 2.2.10 *The family of quasi-injective sheaves is injective with respect to the functor $\text{Hom}_{k_{X_{sa, \mathbb{R}^+}}}(G, \cdot)$ for each $G \in \text{Coh}(X_{sa, \mathbb{R}^+})$.*

In particular

Proposition 2.2.11 *The family of quasi-injective sheaves is injective with respect to the functor $\Gamma(U; \cdot)$ for any $U \in \text{Op}(X_{sa, \mathbb{R}^+})$.*

2.3 An equivalence of categories

Let X be a real analytic manifold endowed with an action μ of $\mathbb{R}^+$. In the following we shall assume the hypothesis below:

$$(2.5) \quad \left\{ \begin{array}{l} \text{(i) every } U \in \text{Op}_{sa}^c(X) \text{ has a finite covering consisting} \\ \text{of } \mathbb{R}^+\text{-connected subanalytic open subsets,} \\ \text{(ii) for any } U \in \text{Op}_{sa}^c(X) \text{ we have } \mathbb{R}^+U \in \text{Op}(X_{sa}), \\ \text{(iii) for any } x \in X \text{ the set } \mathbb{R}^+x \text{ is contractible,} \\ \text{(iv) there exists a covering } \{V_n\}_{n \in \mathbb{N}} \text{ of } X_{sa} \text{ such that} \\ V_n \text{ is } \mathbb{R}^+\text{-connected and } V_n \subset\subset V_{n+1} \text{ for each } n. \end{array} \right.$$

Let $U \in \text{Op}(X_{sa})$ such that $\mathbb{R}^+U$ is still subanalytic. Let φ be the natural map from $\Gamma(\mathbb{R}^+U; F)$ to $\Gamma(U; \eta_{sa}^{-1}F)$ defined by

$$(2.6) \quad \begin{aligned} \Gamma(\mathbb{R}^+U; F) &\rightarrow \Gamma(\mathbb{R}^+U; \eta_{sa*} \eta_{sa}^{-1}F) \\ &\simeq \Gamma(\mathbb{R}^+U; \eta_{sa}^{-1}F) \\ &\rightarrow \Gamma(U; \eta_{sa}^{-1}F). \end{aligned}$$

Proposition 2.3.1 *Let $F \in \text{Mod}(k_{X_{sa, \mathbb{R}^+}})$. Let $U \in \text{Op}(X_{sa})$ and assume that U is $\mathbb{R}^+$ -connected. Then the morphism φ defined by (2.6) is an isomorphism.*

Proof. (i) Assume that $U \in \text{Op}^c(X_{sa})$ is $\mathbb{R}^+$ -connected. Let $F \in \text{Mod}(k_{X_{sa, \mathbb{R}^+}})$, then $F = \varinjlim_i \rho_{\mathbb{R}^+*} F_i$, with $F_i \in \text{Coh}(X_{sa, \mathbb{R}^+})$. We have the chain of isomorphisms

$$\begin{aligned}
\text{Hom}_{k_{X_{sa}}}(k_U, \eta_{sa}^{-1} \varinjlim_i \rho_{\mathbb{R}^+*} F_i) &\simeq \text{Hom}_{k_{X_{sa}}}(k_U, \varinjlim_i \rho_* \eta^{-1} F_i) \\
&\simeq \varinjlim_i \text{Hom}_{k_X}(k_U, \eta^{-1} F_i) \\
&\simeq \varinjlim_i \text{Hom}_{k_{X_{\mathbb{R}^+}}}(k_{\mathbb{R}^+U}, F_i) \\
&\simeq \text{Hom}_{k_{X_{sa, \mathbb{R}^+}}}(k_{\mathbb{R}^+U}, \varinjlim_i \rho_{\mathbb{R}^+*} F_i),
\end{aligned}$$

where the first isomorphism follows since $\eta_{sa}^{-1} \circ \rho_{\mathbb{R}^+*} \simeq \rho_* \circ \eta^{-1}$ by Lemma 2.2.4 and the third one follows from the equivalence between conic sheaves on X and sheaves on $X_{\mathbb{R}^+}$. In the fourth isomorphism we used the fact that $\mathbb{R}^+U \in \text{Op}^c(X_{sa, \mathbb{R}^+})$.

(ii) Let $U \in \text{Op}(X_{sa})$ be $\mathbb{R}^+$ -connected. Let $\{V_n\}_{n \in \mathbb{N}} \in \text{Cov}(X_{sa})$ be a covering of X as in (2.5) (iv) and set $U_n = U \cap V_n$. We have

$$(2.7) \quad \Gamma(U; \eta_{sa}^{-1} F) \simeq \varprojlim_n \Gamma(U_n; \eta_{sa}^{-1} F) \simeq \varprojlim_n \Gamma(\mathbb{R}^+U_n; F) \simeq \Gamma(\mathbb{R}^+U; F).$$

□

We can extend Lemma 2.2.4 to $\text{Mod}(k_{X_{\mathbb{R}^+}})$.

Lemma 2.3.2 *Let $F \in \text{Mod}(k_{X_{\mathbb{R}^+}})$. Then $\eta_{sa}^{-1} \rho_{\mathbb{R}^+*} F \simeq \rho_* \eta^{-1} F$.*

Proof. Let $F \in \text{Mod}(k_{X_{\mathbb{R}^+}})$ and let $U \in \text{Op}^c(X_{sa})$ be $\mathbb{R}^+$ -connected. Then

$$\Gamma(U; \rho_* \eta^{-1} F) \simeq \Gamma(U; \eta^{-1} F) \simeq \Gamma(\mathbb{R}^+U; F),$$

where the second isomorphism follows from Proposition 2.1.5. On the other hand

$$\Gamma(U; \eta_{sa}^{-1} \rho_{\mathbb{R}^+*} F) \simeq \Gamma(\mathbb{R}^+U; \rho_{\mathbb{R}^+*} F) \simeq \Gamma(\mathbb{R}^+U; F),$$

where the second isomorphism follows from Proposition 2.3.1. Hence by (2.5)

$$(i) \quad \eta_{sa}^{-1} \rho_{\mathbb{R}^+*} F \simeq \rho_* \eta^{-1} F.$$

□

Let us consider the category $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ of conic sheaves on X_{sa} . The restriction of η_{sa*} induces a functor denoted by $\tilde{\eta}_{sa*}$ and we obtain a diagram

$$(2.8) \quad \begin{array}{ccc} \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}}) & \xrightleftharpoons[\eta_{sa}^{-1}]{\tilde{\eta}_{sa*}} & \text{Mod}(k_{X_{sa,\mathbb{R}^+}}) \\ \downarrow & \nearrow \eta_{sa*} & \\ \text{Mod}(k_{X_{sa}}) & & \end{array}$$

Theorem 2.3.3 *The functors $\tilde{\eta}_{sa*}$ and η_{sa}^{-1} in (2.8) are equivalences of categories inverse to each others.*

Proof. (i) Let $F \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$, and let $U \in \text{Op}^c(X_{sa})$ be $\mathbb{R}^+$ -connected. We have

$$\Gamma(U; F) \simeq \Gamma(\mathbb{R}^+U; F) \simeq \Gamma(\mathbb{R}^+U; \tilde{\eta}_{sa*}F) \simeq \Gamma(U; \eta_{sa}^{-1}\tilde{\eta}_{sa*}F).$$

The third isomorphism follows from Proposition 2.3.1. Then (2.5) (i) implies $\eta_{sa}^{-1}\tilde{\eta}_{sa*} \simeq \text{id}$.

(ii) For any $U \in \text{Op}^c(X_{sa,\mathbb{R}^+})$ we have:

$$\Gamma(U; \eta_{sa*}\eta_{sa}^{-1}F) \simeq \Gamma(U; \eta_{sa}^{-1}F) \simeq \Gamma(U; F)$$

where the second isomorphism follows from Proposition 2.3.1. This implies $\eta_{sa*}\eta_{sa}^{-1} \simeq \text{id}$. □

Notations 2.3.4 *Since η_{sa}^{-1} is fully faithful and exact we will often identify $\text{Coh}(X_{sa,\mathbb{R}^+})$ with its image in $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$. Hence, for $F \in \text{Coh}(X_{sa,\mathbb{R}^+})$ we shall often write F instead of $\eta_{sa}^{-1}F$.*

Thanks to Theorem 2.2.5 we can give another description of the category of conic sheaves.

Theorem 2.3.5 *Let $F \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$. Then there exists a small filtrant system $\{F_i\}$ in $\text{Coh}(X_{sa,\mathbb{R}^+})$ such that $F \simeq \varinjlim_i \rho_*\eta^{-1}F_i$.*

Remark 2.3.6 *Let $F \in \text{Coh}(X_{sa,\mathbb{R}^+})$. The functor of inverse image commutes with $\varinjlim$ and*

$$\mu^{-1}\rho_*\eta^{-1}F \simeq \rho_*\mu^{-1}\eta^{-1}F \simeq \rho_*p^{-1}\eta^{-1}F \simeq p^{-1}\rho_*\eta^{-1}F.$$

Hence $F \in \text{Coh}(X_{sa, \mathbb{R}^+})$ implies $F \in \text{Mod}^\mu(k_{X_{sa}})$, where $\text{Mod}^\mu(k_{X_{sa}})$ is the category introduced in Remark 2.2.2. Since $\text{Mod}^\mu(k_{X_{sa}})$ is stable by filtrant $\varinjlim$ we have that F belongs to $\text{Mod}^\mu(k_{X_{sa}})$. Hence $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ is a full subcategory of $\text{Mod}^\mu(k_{X_{sa}})$ but $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}}) \not\cong \text{Mod}^\mu(k_{X_{sa}})$ in general. We have the chain of fully faithful functors

$$\text{Coh}(X_{sa, \mathbb{R}^+}) \hookrightarrow \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}}) \hookrightarrow \text{Mod}^\mu(k_{X_{sa}}).$$

2.4 Derived category

Assume (2.5). Injective and quasi-injective objects of $\text{Mod}(k_{X_{sa}})$ are not contained in $\text{Mod}_{\mathbb{R}^+}(k_{sa})$. For this reason we are going to introduce a subcategory which is useful when we try to find acyclic resolutions.

Lemma 2.4.1 *Assume that X satisfies (2.5). Then the following property is satisfied:*

$$(2.9) \quad \begin{cases} \text{each finite covering of an } \mathbb{R}^+\text{-connected } U \in \text{Op}^c(X_{sa}) \\ \text{has a finite refinement } \{V_i\}_{i=1}^n \text{ such that each ordered} \\ \text{union } \bigcup_{i=1}^j V_i \text{ is } \mathbb{R}^+\text{-connected for each } j \in \{1, \dots, n\}. \end{cases}$$

Proof. Let $U \in \text{Op}^c(X_{sa})$ be $\mathbb{R}^+$ -connected. Then each finite covering of U admits a finite refinement consisting of $\mathbb{R}^+$ -connected open subanalytic subsets. Let $\{U_i\}_{i=1}^n$ be a finite covering of U , $U_i \in \text{Op}^c(X_{sa})$ $\mathbb{R}^+$ -connected for each i . We will construct a refinement satisfying (2.9).

For $k = 1, \dots, n$ set $V_{k11} := U_k$ and $V_{k1i} := U_{\sigma(i)} \cap \mathbb{R}^+(U_k \cap U_{\sigma(i)})$ for $i = 2, \dots, n$ and $\sigma(i) = i - 1$ if $i \leq k$, $\sigma(i) = i$ if $i > k$. Then set $U_{k2} := \bigcup_{i=1}^n V_{k1i}$ and $V_{k2i} := U_{\sigma(i)} \cap \mathbb{R}^+(U_{k2} \cap U_{\sigma(i)})$. For $j = 1, \dots, n$ define recursively $U_{kj} = \bigcup_{\ell=1}^j \bigcup_{i=1}^n V_{k\ell i}$ and $V_{kji} = U_{\sigma(i)} \cap \mathbb{R}^+(U_{kj} \cap U_{\sigma(i)})$. Remark that $\bigcup_{p=1}^j \bigcup_{\ell=1}^n \bigcup_{i=1}^n V_{p\ell i} = \bigcup_{p=1}^j \mathbb{R}^+ U_p \cap U$. By Lemma 2.4.2 below all the sets V_{kji} are $\mathbb{R}^+$ -connected and $\{V_{kji}\}_{i,k,j}$ is a refinement of $\{U_i\}_i$ satisfying (2.9) (with the lexicographic order). $\square$

Lemma 2.4.2 *Assume that X satisfies (2.5) (iii). Let U, V, W be open and $\mathbb{R}^+$ -connected. Then $U \cup (V \cap \mathbb{R}^+(U \cap V)) \cup (W \cap \mathbb{R}^+(U \cap W))$ is $\mathbb{R}^+$ -connected.*

Proof. In what follows, when we write $\mathbb{R}^+x$ we suppose that $\mathbb{R}^+x \simeq \mathbb{R}$. If $\mathbb{R}^+x = x$ everything becomes obvious.

(i) First remark that $U \cap V$ (resp. $U \cap W$, $V \cap W$) is $\mathbb{R}^+$ -connected. Indeed, let $x_1 \in U \cap \mathbb{R}^+x$, $x_2 \in V \cap \mathbb{R}^+x$ for some $x \in X$. Then $x_1 = \mu(x, a)$, $x_2 = \mu(x, b)$. Every path in $\mathbb{R}^+x$ connecting x_1 and x_2 contains $\mu(x, [a, b])$. Since U and V are $\mathbb{R}^+$ -connected then $U \cap V \supset \mu(x, [a, b])$.

(ii) Now let us prove that $U \cup (V \cap \mathbb{R}^+(U \cap V))$ is $\mathbb{R}^+$ -connected. Let $x_1, x_2 \in U \cup (V \cap \mathbb{R}^+(U \cap V)) \cap \mathbb{R}^+x$ for some $x \in X$. Then $x_1 = \mu(x, a)$, $x_2 = \mu(x, b)$. We want to prove that $\mu(x, [a, b]) \subset U \cup (V \cap \mathbb{R}^+(U \cap V))$. If $x_1, x_2 \in U$ it follows since U is $\mathbb{R}^+$ -connected and if $x_1, x_2 \in V \cap \mathbb{R}^+(U \cap V)$ it follows from (i). So we may assume that $x_1 \in U$ and $x_2 \in V \cap \mathbb{R}^+(U \cap V)$. Since U is $\mathbb{R}^+$ -connected and $x_2 \in \mathbb{R}^+x_1$, there exists $y = \mu(x, c) \in U \cap V$. Then $\mu(x, [a, c]) \subset U$. In the same way $\mu(x, [b, c]) \subset V \cap \mathbb{R}^+(U \cap V)$ and hence $\mu(x, [a, c] \cup [b, c]) \subset U \cup (V \cap \mathbb{R}^+(U \cap V))$.

(iii) Let us show that $U \cup (V \cap \mathbb{R}^+(U \cap V)) \cup (W \cap \mathbb{R}^+(U \cap W))$ is $\mathbb{R}^+$ -connected. Let $x_1, x_2 \in U \cup (V \cap \mathbb{R}^+(U \cap V)) \cup (W \cap \mathbb{R}^+(U \cap W)) \cap \mathbb{R}^+x$ for some $x \in X$. Then $x_1 = \mu(x, a)$, $x_2 = \mu(x, b)$. We want to prove that $\mu(x, [a, b]) \subset U \cup (V \cap \mathbb{R}^+(U \cap V)) \cup (W \cap \mathbb{R}^+(U \cap W))$. By (i) and (ii) we may reduce to the case $x_1 \in V$, $x_2 \in W$. As in (ii), there exist $y_1 = \mu(x, c) \in U \cap V$ and $y_2 = \mu(x, d) \in U \cap W$. Then $\mu(x, [c, d]) \subset U$, $\mu(x, [a, c]) \subset V \cap \mathbb{R}^+(U \cap V)$ and $\mu(x, [b, d]) \subset W \cap \mathbb{R}^+(U \cap W)$. Hence $\mu(x, [c, d] \cup [a, c] \cup [b, d]) \subset U \cup (V \cap \mathbb{R}^+(U \cap V)) \cup (W \cap \mathbb{R}^+(U \cap W))$ and the result follows. $\square$

Definition 2.4.3 A sheaf $F \in \text{Mod}(k_{X_{sa}})$ is $\mathbb{R}^+$ -quasi-injective if for each $\mathbb{R}^+$ -connected $U \in \text{Op}^c(X_{sa})$ the restriction morphism $\Gamma(X; F) \rightarrow \Gamma(U; F)$ is surjective.

Remark that the functor η_{sa}^{-1} sends quasi-injective objects of $\text{Mod}(k_{X_{sa, \mathbb{R}^+}})$ to $\mathbb{R}^+$ -quasi-injective objects since $\Gamma(U; \eta_{sa}^{-1}F) \simeq \Gamma(\mathbb{R}^+U; F)$ if $U \in \text{Op}^c(X_{sa})$ is $\mathbb{R}^+$ -connected. Moreover the category of $\mathbb{R}^+$ -injective objects is cogenerating since injective objects are cogenerating in $\text{Mod}(k_{X_{sa}})$.

Proposition 2.4.4 Let $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ be an exact sequence in $\text{Mod}(k_{X_{sa}})$ and assume that F' is $\mathbb{R}^+$ -quasi-injective. Let $U \in \text{Op}^c(X_{sa})$ be $\mathbb{R}^+$ -connected. Then the sequence

$$0 \rightarrow \Gamma(U; F') \rightarrow \Gamma(U; F) \rightarrow \Gamma(U; F'') \rightarrow 0$$

is exact.

Proof. Let $s'' \in \Gamma(U; F'')$, and let $\{V_i\}_{i=1}^n$ be a finite covering of U satisfying (2.9) and such that there exists $s_i \in \Gamma(V_i; F)$ whose image is $s''|_{V_i}$. For $n \geq 2$ on $V_1 \cap V_2$ $s_1 - s_2$ defines a section of $\Gamma(V_1 \cap V_2; F')$ which extends to $s' \in \Gamma(X; F')$. Replace s_1 with $s_1 - s'$. We may suppose that $s_1 = s_2$ on $V_1 \cap V_2$. Then there exists $t \in \Gamma(V_1 \cup V_2)$ such that $t|_{V_i} = s_i$, $i = 1, 2$. Thus the induction proceeds. $\square$

Proposition 2.4.5 *Let $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ be an exact sequence in $\text{Mod}(k_{X_{sa}})$ and assume that F' is $\mathbb{R}^+$ -quasi-injective. Let $U \in \text{Op}(X_{sa})$ be $\mathbb{R}^+$ -connected. Then the sequence*

$$0 \rightarrow \Gamma(U; F') \rightarrow \Gamma(U; F) \rightarrow \Gamma(U; F'') \rightarrow 0$$

is exact.

Proof. By (2.5) (iv) there exists a covering $\{V_n\}_{n \in \mathbb{N}}$ of X_{sa} such that V_n is $\mathbb{R}^+$ -connected and $V_n \subset \subset V_{n+1}$ for each n . For each n the sequence

$$0 \rightarrow \Gamma(U \cap V_n; F') \rightarrow \Gamma(U \cap V_n; F) \rightarrow \Gamma(U \cap V_n; F'') \rightarrow 0$$

is exact and the morphism $\Gamma(U \cap V_{n+1}; F') \rightarrow \Gamma(U \cap V_n; F')$ is surjective for each n since F' is $\mathbb{R}^+$ -quasi-injective. Then by the Mittag-Leffler property (see Proposition 1.12.3 of [11]) the sequence

$$0 \rightarrow \varprojlim_n \Gamma(U \cap V_n; F') \rightarrow \varprojlim_n \Gamma(U \cap V_n; F) \rightarrow \varprojlim_n \Gamma(U \cap V_n; F'') \rightarrow 0$$

is exact. Since $\varprojlim_n \Gamma(U \cap V_n; G) \simeq \Gamma(U; G)$ for each $G \in \text{Mod}(k_{X_{sa}})$ the result follows. $\square$

Proposition 2.4.6 *Let F', F be $\mathbb{R}^+$ -quasi-injective and consider the exact sequence $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ in $\text{Mod}(k_{X_{sa}})$. Then F'' is $\mathbb{R}^+$ -quasi-injective.*

Proof. Let $U \in \text{Op}^c(X_{sa})$ be $\mathbb{R}^+$ -connected and let us consider the diagram below

$$\begin{array}{ccc} \Gamma(X; F) & \longrightarrow & \Gamma(X; F'') \\ \downarrow \alpha & & \downarrow \gamma \\ \Gamma(U; F) & \xrightarrow{\beta} & \Gamma(U; F''). \end{array}$$

The morphism α is surjective since F is $\mathbb{R}^+$ -quasi-injective and β is surjective by Proposition 2.4.4. Then γ is surjective. $\square$

It follows from the preceding results that

Proposition 2.4.7 *$\mathbb{R}^+$ -quasi-injective objects are injective with respect to the functor $\Gamma(U; \cdot)$, with $U \in \text{Op}(X_{sa})$ and $\mathbb{R}^+$ -connected.*

Corollary 2.4.8 *$\mathbb{R}^+$ -quasi-injective objects are η_{sa*} -injective.*

Proof. Let $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ be an exact sequence in $\text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ and assume that F' is $\mathbb{R}^+$ -quasi-injective. By Proposition 2.4.7 the sequence

$$0 \rightarrow \Gamma(U; F') \rightarrow \Gamma(U; F) \rightarrow \Gamma(U; F'') \rightarrow 0$$

for any $U \in \text{Op}(X_{sa, \mathbb{R}^+})$. This implies that the sequence

$$0 \rightarrow \eta_{sa*} F' \rightarrow \eta_{sa*} F \rightarrow \eta_{sa*} F'' \rightarrow 0$$

is exact. $\square$

Theorem 2.4.9 *The categories $D^b(k_{X_{sa, \mathbb{R}^+}})$ and $D_{\mathbb{R}^+}^b(k_{X_{sa}})$ are equivalent.*

Proof. In order to prove this statement, it is enough to show that η_{sa}^{-1} is fully faithful. Let $F \in D^b(k_{X_{sa, \mathbb{R}^+}})$ and let F' be an injective complex quasi-isomorphic to F . Since η_{sa}^{-1} sends injective objects to $\mathbb{R}^+$ -quasi-injective objects which are η_{sa*} -injective we have $R\eta_{sa*}\eta_{sa}^{-1}F \simeq \eta_{sa*}\eta_{sa}^{-1}F' \simeq F' \simeq F$. This implies $R\eta_{sa*}\eta_{sa}^{-1} \simeq \text{id}$, hence η_{sa}^{-1} is fully faithful. $\square$

Hence for each $F \in D_{\mathbb{R}^+}^b(k_{X_{sa}})$ we have $F \simeq \eta_{sa}^{-1}F'$ with $F' \in D^b(k_{X_{sa, \mathbb{R}^+}})$.

Proposition 2.4.10 *Let $F \in D_{\mathbb{R}^+}^b(k_{X_{sa}})$ and let $U \in \text{Op}(X_{sa})$ be $\mathbb{R}^+$ -connected. There is an isomorphism $R\Gamma(\mathbb{R}^+U; F) \xrightarrow{\sim} R\Gamma(U; \eta_{sa}^{-1}F)$.*

Proof. Let F' be a complex of injective objects quasi-isomorphic to F . Since η_{sa}^{-1} sends injective objects to $\mathbb{R}^+$ -quasi-injective objects we have

$$\begin{aligned} R\Gamma(\mathbb{R}^+U; F) &\simeq \Gamma(\mathbb{R}^+U; F') \\ &\xrightarrow{\sim} \Gamma(U; \eta_{sa}^{-1}F') \\ &\simeq R\Gamma(U; \eta_{sa}^{-1}F), \end{aligned}$$

where the second isomorphism follows from Proposition 2.3.1. $\square$

We extend Lemma 2.3.2 to $D^b(k_{X_{\mathbb{R}^+}})$.

Lemma 2.4.11 *Let $F \in D^b(k_{X_{\mathbb{R}^+}})$. Then $\eta_{sa}^{-1}R\rho_{\mathbb{R}^+*}F \simeq R\rho_*\eta^{-1}F$.*

Proof. (i) Let $F \in \text{Mod}(k_{X_{\mathbb{R}^+}})$ be injective. Then for each $\mathbb{R}^+$ -connected $U \in \text{Op}^c(X_{sa})$ $\text{R}\Gamma(U; R\rho_*\eta^{-1}F) \simeq \text{R}\Gamma(U; \eta^{-1}F) \simeq \text{R}\Gamma(\mathbb{R}^+U; F)$ is concentrated in degree zero. Hence $\eta^{-1}F$ is $R\rho_*$ -acyclic by (2.5) (i).

(ii) Let $F \in D_{\mathbb{R}^+}^b(k_X)$ and let F' be a complex of injective objects quasi-isomorphic to F . Then $\eta_{sa}^{-1}R\rho_{\mathbb{R}^+*}F \simeq \eta_{sa}^{-1}\rho_{\mathbb{R}^+*}F' \simeq \rho_*\eta^{-1}F' \simeq R\rho_*\eta^{-1}F$, where the second isomorphism follows from Lemma 2.3.2 and the third one follows from (i). $\square$

2.5 Operations

Let X be a real analytic manifold endowed with an analytic action of $\mathbb{R}^+$. In this section we study the operations in the category of conic sheaves on X_{sa} .

Proposition 2.5.1 *Let $F \in \text{Mod}_{\mathbb{R}^+}(k_X)$ and let $G \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$.*

(i) *we have $\rho_*F \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$,*

(ii) *we have $\rho^{-1}G \in \text{Mod}_{\mathbb{R}^+}(k_X)$.*

Proof. (i) Let $U \in \text{Op}^c(X_{sa})$ be $\mathbb{R}^+$ -connected. We have the chain of isomorphisms

$$\Gamma(U; \rho_*F) \simeq \Gamma(U; F) \simeq \Gamma(\mathbb{R}^+U; F) \simeq \Gamma(\mathbb{R}^+U; \rho_*F).$$

(ii) We have $G = \varinjlim_j \rho_*G_j$, with $G_j \in \text{Coh}(X_{sa, \mathbb{R}^+})$. Then

$$\rho^{-1}\varinjlim_j \rho_*G_j \simeq \varinjlim_j \rho^{-1}\rho_*G_j \simeq \varinjlim_j G_j$$

and $\varinjlim_j G_j$ belongs to $\text{Mod}_{\mathbb{R}^+}(k_X)$. $\square$

Proposition 2.5.2 *Let $F, G \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$. Then*

- (i) *we have $F \otimes G \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$,*
- (ii) *we have $\mathcal{H}om(F, G) \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$.*

Proof. We have $F = \varinjlim_i \rho_* F_i$ and $G = \varinjlim_j \rho_* G_j$, with $F_i, G_j \in \text{Coh}(X_{sa, \mathbb{R}^+})$.

- (i) we have $F \otimes G \simeq \varinjlim_{i,j} \rho_*(F_i \otimes G_j)$ and $F_i \otimes G_j$ belongs to $\text{Coh}(X_{sa, \mathbb{R}^+})$

for each i, j .

- (ii) we have $\mathcal{H}om(F, G) \simeq \varprojlim_i \varinjlim_j \rho_* \mathcal{H}om(F_i, G_j)$ and $\mathcal{H}om(F_i, G_j)$ is conic

for each i, j .

□

Let $f : X \rightarrow Y$ be a conic morphism of real analytic manifolds.

Proposition 2.5.3 *Let $F \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$ and let $G \in \text{Mod}_{\mathbb{R}^+}(k_{Y_{sa}})$.*

- (i) *we have $f_* F \in \text{Mod}_{\mathbb{R}^+}(k_{Y_{sa}})$,*
- (ii) *we have $f^{-1} G \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$.*

Proof. (i) Let $U \in \text{Op}^c(Y_{sa})$ be $\mathbb{R}^+$ -connected. Since f commutes with the action of $\mathbb{R}^+$, the set $f^{-1}(U)$ is $\mathbb{R}^+$ connected. We have the chain of isomorphisms

$$\Gamma(f^{-1}(U); F) \simeq \Gamma(\mathbb{R}^+ f^{-1}(U); F) \simeq \Gamma(f^{-1}(\mathbb{R}^+ U); F).$$

Hence $\Gamma(U; f_* F) \simeq \Gamma(\mathbb{R}^+ U; f_* F)$.

- (ii) We have $G = \varinjlim_j \rho_* G_j$, with $G_j \in \text{Coh}(Y_{sa, \mathbb{R}^+})$. Then

$$f^{-1} \varinjlim_j \rho_* G_j \simeq \varinjlim_j f^{-1} \rho_* G_j \simeq \varinjlim_j \rho_* f^{-1} G_j$$

and $f^{-1} G_j$ is conic for each j .

□

Now let us consider the operations in the derived category of conic sub-analytic sheaves.

Proposition 2.5.4 *Let $F \in \text{Coh}(X_{sa, \mathbb{R}^+})$ and let $G \in D_{\mathbb{R}^+}^b(k_{X_{sa}})$. Then $R\mathcal{H}om(F, G) \in D_{\mathbb{R}^+}^b(k_{X_{sa}})$.*

Proof. We may reduce to the case $G \in \text{Mod}_{\mathbb{R}^+}(k_{X_{sa}})$. Then $G = \varinjlim_j \rho_* G_j$, with $G_j \in \text{Coh}(X_{sa, \mathbb{R}^+})$. Then by Proposition 2.2.2 of [23]

$$R^k \mathcal{H}om(F, G) \simeq \varinjlim_i \rho_* R^k \mathcal{H}om(F, G_j)$$

for each $k \in \mathbb{Z}$ and the result follows since $R^k \mathcal{H}om(F, G_j)$ is conic for each $k \in \mathbb{Z}$ and for each j . $\square$

Proposition 2.5.5 *Let $F \in D_{\mathbb{R}^+}^b(k_{X_{sa}})$. Then $Rf_* F \in D_{\mathbb{R}^+}^b(k_{Y_{sa}})$.*

Proof. Remark that the functor η_{sa}^{-1} sends injective sheaves to f_* -acyclic sheaves. This is a consequence of the fact that η_{sa}^{-1} sends injective sheaves to $\Gamma(U; \cdot)$ -acyclic sheaves for each $\mathbb{R}^+$ -connected $U \in \text{Op}(X_{sa})$. There exists $F' \in D^b(k_{X_{sa, \mathbb{R}^+}})$ such that $F \simeq \eta_{sa}^{-1} F'$. Let $I^\bullet$ be a bounded injective resolution of F' . Then $\eta_{sa}^{-1} I^j$ is conic and f_* -acyclic for each j . We have $Rf_* F \simeq f_* \eta_{sa}^{-1} I^\bullet$ and $f_* \eta_{sa}^{-1} I^j$ is conic for each j . $\square$

Proposition 2.5.6 *Let $G \in D_{\mathbb{R}^+}^b(k_{Y_{sa}})$. Then $f^! G \in D_{\mathbb{R}^+}^b(k_{X_{sa}})$.*

Proof. We may reduce to the case $G \in \text{Mod}_{\mathbb{R}^+}(k_{Y_{sa}})$. Then $G = \varinjlim_j \rho_* G_j$, with $G_j \in \text{Coh}(Y_{sa, \mathbb{R}^+})$. By Proposition 2.4.5 of [23] we have $H^k f^! G \simeq \varinjlim_i \rho_* H^k f^! G_j$ for each $k \in \mathbb{Z}$ and the result follows since $H^k f^! G_j$ is conic for each $k \in \mathbb{Z}$ and for each j . $\square$

Remark 2.5.7 *The functor $f_{!!} : \text{Mod}(k_{X_{sa}}) \rightarrow \text{Mod}(k_{Y_{sa}})$ does not send conic sheaves to conic sheaves in general. In fact, let $p : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be the projection. It is a conic map with respect to the natural action of $\mathbb{R}^+$ on $\mathbb{R}^3$ and $\mathbb{R}^2$. Set*

$$\begin{aligned} U &= \{(x, y) \in \mathbb{R}^2; (x-1)^2 + y^2 < 1\}, \\ B_n &= \{(x, y) \in \mathbb{R}^2; x^2 + y^2 < n\}, \\ B_n^+ &= B_n \cap (\mathbb{R}^+ \times \mathbb{R}), \\ S &= \mathbb{R}^+(\partial U \times \{1\}). \end{aligned}$$

Let us consider the conic sheaf k_S . By definition of proper direct image we have $\Gamma(U; p_{!!}F) = \varinjlim_K \Gamma(p^{-1}(U); \Gamma_K F)$, where K ranges through the family of subanalytic compact subsets of $\mathbb{R}^3$. Since U is bounded we have

$$\Gamma(U; p_{!!}k_S) \simeq \varinjlim_K \Gamma(U \times \mathbb{R}; \Gamma_K k_S) \simeq \varinjlim_m \Gamma(U \times \mathbb{R}; \Gamma_{\mathbb{R}^2 \times [-m, m]} k_S) \simeq k,$$

where $m \in \mathbb{N}$. On the other hand we have

$$\Gamma(\mathbb{R}^+ U; p_{!!}k_S) \simeq \varprojlim_n \Gamma(B_n^+; p_{!!}k_S) \simeq \varprojlim_n \varinjlim_m \Gamma(B_n^+ \times \mathbb{R}; \Gamma_{\mathbb{R}^2 \times [-m, m]} k_S) = 0,$$

where $m, n \in \mathbb{N}$, since $\Gamma(B_n^+ \times \mathbb{R}; \Gamma_{\mathbb{R}^2 \times [-m, m]} k_S) = 0$ for each m, n .

Hence we shall need a new definition of proper direct image for conic sheaves.

Definition 2.5.8 We define functor $f_{\mathbb{R}+!!}$ of proper direct image for conic sheaves in the following way

$$\begin{aligned} f_{\mathbb{R}+!!} : \text{Mod}_{\mathbb{R}+}(k_{X_{sa}}) &\rightarrow \text{Mod}_{\mathbb{R}+}(k_{Y_{sa}}) \\ \varinjlim_i \rho_* F_i &\mapsto \varinjlim_i \rho_* f_! F_i, \end{aligned}$$

where $F_i \in \text{Coh}(X_{sa, \mathbb{R}+})$.

Note that if $F \in \text{Coh}(X_{sa, \mathbb{R}+})$ then $f_{\mathbb{R}+!!} \rho_* F \simeq \rho_* f_! F \not\simeq f_{!!} \rho_* F$. Moreover this definition is compatible with the classical one. In fact $f_{\mathbb{R}+!!}$ commutes with ρ^{-1} and we have the following commutative diagram

$$\begin{array}{ccc} \text{Mod}_{\mathbb{R}+}(k_X) & \xrightarrow{f_!} & \text{Mod}_{\mathbb{R}+}(k_Y) \\ \downarrow \rho_* & & \uparrow \rho^{-1} \\ \text{Mod}_{\mathbb{R}+}(k_{X_{sa}}) & \xrightarrow{f_{\mathbb{R}+!!}} & \text{Mod}_{\mathbb{R}+}(k_{Y_{sa}}). \end{array}$$

Remark 2.5.9 With the notation of Remark 2.5.7, we have

$$\Gamma(\mathbb{R}^+ U; p_{\mathbb{R}+!!} k_S) \simeq \Gamma(U; p_{\mathbb{R}+!!} k_S) \simeq k.$$

In fact the restriction of p to $S \cap \{(x, y, z) \in \mathbb{R}^3; x > 0\}$ is proper.

Let us see an explicit formula for the sections of $f_{\mathbb{R}+!!}$. Let $U \in \text{Op}^c(X_{sa, \mathbb{R}+})$ and let $F = \varinjlim_i \rho_* F_i$ with $F_i \in \text{Coh}(X_{sa, \mathbb{R}+})$. We have the chain of isomorphisms

$$\begin{aligned}
\Gamma(U; \varinjlim_i \rho_* f_! F_i) &\simeq \varinjlim_i \Gamma(U; f_! F_i) \\
&\simeq \varinjlim_{i, Z, K} \Gamma(f^{-1}(U); \Gamma_{Z \cap K} F_i) \\
&\simeq \varinjlim_{i, Z', K} \Gamma(f^{-1}(U); \Gamma_{Z' \cap K} F_i) \\
&\simeq \varinjlim_{Z', K} \Gamma(f^{-1}(U); \Gamma_{Z' \cap K} \varinjlim_i \rho_* F_i).
\end{aligned}$$

Here Z ranges into the family of closed subanalytic subsets of $f^{-1}(U)$ such that $f : Z \rightarrow U$ is proper, Z' ranges through the family of closed conic subanalytic subsets of $f^{-1}(U)$ such that $f^{-1}(y) \cap \mathbb{R}^+ x = \{\text{point}\}$ for any $y \in Y$, $x \in X$, and $K \subset\subset X_{\mathbb{R}+}$ are conic and closed. The first isomorphism follows since $U \in \text{Op}^c(X_{sa, \mathbb{R}+})$, the third since F_i is conic for each i and the last one since $f^{-1}(U) \cap K \subset\subset X_{\mathbb{R}+}$.

It is easy to prove that projection formula and base change formula for conic sheaves are satisfied. Moreover, $\mathbb{R}^+$ -quasi-injective objects are acyclic with respect to the functor $f_{\mathbb{R}+!!}$, since they are $\mathcal{H}om(G, \cdot)$ -injective for each $G \in \text{Coh}(X_{sa, \mathbb{R}+})$.

In order to find a right adjoint to $Rf_{\mathbb{R}+!!}$ we follow the method used to find a right adjoint to the functor proper direct image for subanalytic sheaves. We shall skip the details of the proof (which are an adaptation of the results of [23]). The subcategory $\mathcal{J}_{X_{sa, \mathbb{R}+}}$ of $\mathbb{R}^+$ -quasi-injective objects and the functor $f_{\mathbb{R}+!!}$ have the following properties:

$$\left\{ \begin{array}{l}
\text{(i) } \mathcal{J}_{X_{sa, \mathbb{R}+}} \text{ is cogenerating,} \\
\text{(ii) } \text{Mod}(k_{X_{sa, \mathbb{R}+}}) \text{ has finite quasi-injective dimension,} \\
\text{(iii) } \mathcal{J}_{X_{sa, \mathbb{R}+}} \text{ is } f_{\mathbb{R}+!!}\text{-injective,} \\
\text{(iv) } \mathcal{J}_{X_{sa, \mathbb{R}+}} \text{ is closed by small } \oplus, \\
\text{(v) } f_{\mathbb{R}+!!} \text{ commutes with small } \oplus.
\end{array} \right.$$

As a consequence of the Brown representability theorem (see [12], Corollary 14.3.7 for details) we find a right adjoint to the functor $Rf_{\mathbb{R}+!!}$, denoted by

$f_{\mathbb{R}^+}^!$. By adjunction $f_{\mathbb{R}^+}^!$ commutes with $R\rho_*$ and as in [23] one can prove that $H^k f_{\mathbb{R}^+}^!$ commutes with filtrant $\varinjlim$. Hence $f_{\mathbb{R}^+}^!$ coincides with the restriction of $f^!$ to $D_{\mathbb{R}^+}^b(k_{Y_{sa}})$.

3 Fourier-Sato transform for subanalytic sheaves.

In this section we construct the Fourier-Sato transform for subanalytic sheaves. Reference are made to [11] for the classical Fourier-Sato transform.

3.1 Conic sheaves on vector bundles

Let $E \xrightarrow{\tau} Z$ be a real vector bundle, with dimension n over a real analytic manifold Z . Then $\mathbb{R}^+$ acts naturally on E by multiplication on the fibers. We identify Z with the zero-section of E and denote by $i : Z \hookrightarrow E$ the embedding. We set $\dot{E} = E \setminus Z$ and $\dot{\tau} : \dot{E} \rightarrow Z$ denotes the projection.

Lemma 3.1.1 *The category $\text{Op}(E_{sa})$ satisfies (2.5).*

Proof. Let us prove (2.5) (i). Let $\{V_i\}_{i \in \mathbb{N}}$ be a locally finite covering of Z with $V_i \in \text{Op}^c(Z_{sa})$ such that $\tau^{-1}(V_i) \simeq \mathbb{R}^m \times \mathbb{R}^n$ and let $\{U_i\}$ be a refinement of $\{V_i\}$ with $U_i \in \text{Op}^c(Z_{sa})$ and $\overline{U_i} \subset V_i$ for each i . Then U is covered by a finite number of $\tau^{-1}(U_i)$ and $U \cap \tau^{-1}(U_i)$ is relatively compact in $\tau^{-1}(V_i)$ for each i . We may reduce to the case $E \simeq \mathbb{R}^m \times \mathbb{R}^n$. Let us consider the morphism of manifolds

$$\begin{aligned} \varphi : \mathbb{R}^m \times \mathbb{S}^{n-1} \times \mathbb{R} &\rightarrow \mathbb{R}^m \times \mathbb{R}^n \\ (z, \vartheta, r) &\mapsto (z, ri(\vartheta)), \end{aligned}$$

where $i : \mathbb{S}^{n-1} \hookrightarrow \mathbb{R}^n$ denotes the embedding. Then φ is proper and subanalytic. The subset $\varphi^{-1}(U)$ is subanalytic and relatively compact in $\mathbb{R}^m \times \mathbb{S}^{n-1} \times \mathbb{R}$.

(a) By Lemma A.1.11 $\varphi^{-1}(U \setminus Z)$ admits a finite cover $\{W_j\}_{j \in J}$ such that the intersections of each W_j with the fibers of $\pi : \mathbb{R}^m \times \mathbb{S}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}^m \times \mathbb{S}^{n-1}$ are contractible or empty. Then $\varphi(W_j)$ is an open subanalytic relatively compact $\mathbb{R}^+$ -connected subset of $\mathbb{R}^m \times \mathbb{R}^n$ for each j . In this way we obtain a finite covering of $U \setminus Z$ consisting of $\mathbb{R}^+$ -connected subanalytic open subsets.

(b) Let $p \in \pi(\varphi^{-1}(U \cap Z))$. Then $\pi^{-1}(p) \cap U$ is a disjoint union of intervals. Let us consider the interval $(m(p), M(p))$, $m(p) < M(p) \in \mathbb{R}$ containing 0. Set $W_Z = \{(p, r) \in U; m(p) < r < M(p)\}$. The set W_Z is open subanalytic

(it is a consequence of Proposition 1.2, Chapter 6 of [31]), contains $\varphi^{-1}(U \cap Z)$ and its intersections with the fibers of π are contractible. Then $\varphi(W_Z)$ is an open $\mathbb{R}^+$ -connected subanalytic neighborhood of $U \cap Z$ and it is contained in U .

By (a) there exists a finite covering $\{\varphi(W_j)\}_{j \in J}$ of $U \setminus Z$ consisting of $\mathbb{R}^+$ -connected subanalytic open subsets, and $\varphi(W_Z) \cup \bigcup_{j \in J} \varphi(W_j) = U$.

By Proposition 8.3.8 of [11] the category $\text{Op}(E_{sa})$ also satisfies (2.5) (ii). Moreover (2.5) (iii) and (iv) are clearly satisfied. $\square$

Now let us consider E endowed with the conic topology. In this situation, an object $U \in \text{Op}(E_{\mathbb{R}^+})$ is the union of $\dot{U} \in \text{Op}(\dot{E}_{\mathbb{R}^+})$ and $U_Z \in \text{Op}(Z)$ such that $\dot{\tau}^{-1}(U_Z) \subset \dot{U}$. If $U, V \in \text{Op}(E_{\mathbb{R}^+})$, then $U \subset\subset V$ if $U_Z \subset\subset V_Z$ in Z and $\dot{U} \subset\subset \dot{V}$ in $\dot{E}_{\mathbb{R}^+}$ (this means that $\pi(\dot{U}) \subset\subset \pi(\dot{V})$ in $\dot{E}/\mathbb{R}^+$, where $\pi : \dot{E} \rightarrow \dot{E}/\mathbb{R}^+$ denotes the projection).

Applying Theorem 2.4.9 we have the following

Theorem 3.1.2 *The categories $D_{\mathbb{R}^+}^b(k_{E_{sa}})$ and $D^b(k_{E_{sa, \mathbb{R}^+}})$ are equivalent.*

Consider the subcategory $\text{Mod}_{\mathbb{R}\text{-c}, \mathbb{R}^+}^{cb}(k_E)$ of $\text{Mod}_{\mathbb{R}\text{-c}, \mathbb{R}^+}(k_E)$ consisting of sheaves whose support is compact on the base (i.e. $\tau(\text{supp}(F))$ is compact in Z).

Let us consider the natural map $\eta : E \rightarrow E_{\mathbb{R}^+}$. The restriction of η^{-1} to $\text{Coh}(E_{sa, \mathbb{R}^+})$ gives rise to a functor

$$(3.1) \quad \overline{\eta}^{-1} : \text{Coh}(E_{sa, \mathbb{R}^+}) \rightarrow \text{Mod}_{\mathbb{R}\text{-c}, \mathbb{R}^+}^{cb}(k_E)$$

Since the functor η^{-1} is fully faithful and exact, we identify $\text{Coh}(E_{sa, \mathbb{R}^+})$ as a subcategory of $\text{Mod}_{\mathbb{R}\text{-c}, \mathbb{R}^+}^{cb}(k_E)$.

Theorem 3.1.3 *The functor $\overline{\eta}^{-1}$ in (3.1) is an equivalence of categories.*

Proof. (i) Let $F \in \text{Mod}_{\mathbb{R}\text{-c}, \mathbb{R}^+}^{cb}(k_E)$. Let us show that F is $E_{sa, \mathbb{R}^+}$ -finite. We may reduce to the case $E \simeq \mathbb{R}^m \times \mathbb{R}^n$ and $Z \simeq \mathbb{R}^m \times \{0\}$. It is well known that if X is a real analytic manifold and $G \in \text{Mod}_{\mathbb{R}\text{-c}}^c(k_X)$, then G is quasi-isomorphic to a bounded complex of finite sums $\oplus_W k_W$, where

$W \in \text{Op}_{sa}^c(X)$.

Let us consider the diagram $Z \xrightarrow{i} E \xrightarrow{\tau} Z$, where i is the embedding. We have $\tau_* F \in \text{Mod}_{\mathbb{R}-c}^c(k_Z)$. We have an exact sequence

$$(3.2) \quad \oplus_{i \in I} k_{\tau^{-1}(V_i)} \rightarrow \tau^{-1} \tau_* F \rightarrow 0,$$

where I is finite and $V_i \in \text{Op}_{sa}^c(Z)$.

Now let us consider the diagram $S \xrightarrow{j} \dot{E} \xrightarrow{\pi} S$, where $S = \dot{E}/\mathbb{R}^+ \simeq \mathbb{R}^m \times \mathbb{S}^{n-1}$ and π is the projection. We have $j^{-1} F|_{\dot{E}} \in \text{Mod}_{\mathbb{R}-c}^c(k_S)$. Since $F|_{\dot{E}}$ is conic $\pi^{-1} j^{-1} F|_{\dot{E}} \simeq F|_{\dot{E}}$. We have an exact sequence

$$(3.3) \quad \oplus_j k_{\pi^{-1}(U_j)} \rightarrow F|_{\dot{E}} \rightarrow 0.$$

where J is finite and $U_j \in \text{Op}_{sa}^c(S)$.

It is easy to check that the morphism $\tau^{-1} \tau_* F \oplus F|_{\dot{E}} \rightarrow F$ is an epimorphism and we obtain the result by (3.2) and (3.3).

(ii) Let us show that F is $E_{sa, \mathbb{R}^+}$ -pseudo-coherent. Let $G = \oplus_{i \in I} k_{W_i}$, with I finite and $W_i \in \text{Op}^c(E_{sa, \mathbb{R}^+})$, and consider a morphism $\psi : G \rightarrow F$. Since F and G are $\mathbb{R}$ -constructible and conic, then $\ker \psi$ belongs to $\text{Mod}_{\mathbb{R}-c, \mathbb{R}^+}(k_E)$, and its support is still compact on the base.

□

As a consequence of Theorems 2.3.5 and 3.1.3 one has the following

Theorem 3.1.4 *Let $F \in \text{Mod}_{\mathbb{R}^+}(k_{E_{sa}})$. Then there exists a small filtrant system $\{F_i\}$ in $\text{Mod}_{\mathbb{R}-c, \mathbb{R}^+}^{cb}(k_E)$ such that $F \simeq \varinjlim_i \rho_* F_i$.*

We end this section with the following result, which will be useful in § 3.2.

Lemma 3.1.5 *Let $F \in D_{\mathbb{R}^+}^b(k_{E_{sa}})$. Then:*

$$(i) \quad R\tau_* F \simeq i^{-1} F.$$

$$(ii) \quad R\tau_{!!} F \simeq i^! F.$$

Proof. (i) The adjunction morphism defines $R\tau_* F \simeq i^{-1} \tau^{-1} R\tau_* F \rightarrow i^{-1} F$. Let $V \in \text{Op}^c(Z_{sa})$. Then

$$\varinjlim_{U \supset V} R^k \Gamma(U; F) \simeq \varinjlim_{\substack{U \supset V \\ \tau(U)=V}} R^k \Gamma(U; F) \simeq R^k \Gamma(\tau^{-1}(V); F) \simeq R^k \Gamma(V; R\tau_* F),$$

where $U \in \text{Op}(X_{sa})$ and $\mathbb{R}^+$ -connected. The second isomorphism follows from Corollary 2.4.10.

(ii) The adjunction morphism defines $i^!F \rightarrow i^!\tau^!R\tau_!F \simeq R\tau_!F$. Let $V \in \text{Op}^c(Z_{sa})$, and let K be a compact subanalytic $\mathbb{R}^+$ -connected neighborhood of V in E . Then $\tau^{-1}(V) \setminus K$ is $\mathbb{R}^+$ -connected and subanalytic, and $\mathbb{R}^+(\tau^{-1}(V) \setminus K) = \tau^{-1}(V) \setminus Z$. By Corollary 2.4.10 we have the isomorphism $R\Gamma(\tau^{-1}(V); R\Gamma_Z F) \simeq R\Gamma(\tau^{-1}(V); R\Gamma_K F)$.

It follows from the definition of $R\tau_!$ that for any $k \in \mathbb{Z}$ and $V \in \text{Op}^c(Z_{sa})$ we have $R^k\Gamma(V; R\tau_!F) \simeq \varinjlim_K R^k\Gamma(\tau^{-1}(V); R\Gamma_K F)$, where K ranges through the family of compact subanalytic $\mathbb{R}^+$ -connected neighborhoods of V in E .

On the other hand for any $k \in \mathbb{Z}$ we have $R^k\Gamma(V; i^!F) \simeq R^k\text{Hom}(i_*k_V, F) \simeq R^k\text{Hom}(i_*i^{-1}\tau^{-1}k_V, F) \simeq R^k\Gamma(\tau^{-1}(V), R\Gamma_Z F)$ and the result follows. $\square$

3.2 Fourier-Sato transformation

Let $E \xrightarrow{\tau} Z$ be a real vector bundle, with dimension n over a real analytic manifold Z and $E^* \xrightarrow{\pi} Z$ its dual. We identify Z as the zero-section of E and denote $i : Z \hookrightarrow E$ the embedding, we define similarly $i : Z \hookrightarrow E^*$. We denote by p_1 and p_2 the projections from $E \times_Z E^*$:

$$\begin{array}{ccccc}
 & & E \times_Z E^* & & \\
 & \swarrow p_1 & & \searrow p_2 & \\
 E & & & & E^* \\
 & \searrow \tau & & \swarrow \pi & \\
 & & Z & &
 \end{array}$$

We set

$$P := \{(x, y) \in E \times_Z E^*; \langle x, y \rangle \geq 0\}$$

$$P' := \{(x, y) \in E \times_Z E^*; \langle x, y \rangle \leq 0\}$$

and we define the functors

$$\left\{ \begin{array}{l} \Psi_{P'} = Rp_{1*} \circ R\Gamma_{P'} \circ p_2^! : D_{\mathbb{R}+}^b(k_{E_{sa}^*}) \rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}}) \\ \Phi_{P'} = Rp_{2!!} \circ (\cdot)_{P'} \circ p_1^{-1} : D_{\mathbb{R}+}^b(k_{E_{sa}}) \rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}^*}) \\ \Psi_P = Rp_{2*} \circ R\Gamma_P \circ p_1^{-1} : D_{\mathbb{R}+}^b(k_{E_{sa}}) \rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}^*}) \\ \Phi_P = Rp_{1!!} \circ (\cdot)_P \circ p_2^! : D_{\mathbb{R}+}^b(k_{E_{sa}^*}) \rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}}) \end{array} \right.$$

Remark 3.2.1 *These functors are well defined, more generally they send subanalytic sheaves to conic subanalytic sheaves.*

Lemma 3.2.2 *Let $F \in D_{\mathbb{R}+}^b(k_{E_{sa}})$. Then $\text{supp}((R\Gamma_P(p_1^{-1}F))_{P'})$ is contained in $Z \times_Z E^*$.*

Proof. We may reduce to the case $F \in \text{Mod}_{\mathbb{R}+}(k_{E_{sa}})$. Then $F = \varinjlim_i \rho_* F_i$, with $F_i \in \text{Mod}_{\mathbb{R}-c, \mathbb{R}+}^{cb}(k_E)$. We have

$$\begin{aligned} H^k(R\Gamma_P(p_1^{-1} \varinjlim_i \rho_* F_i)_{P'}) &\simeq \varinjlim_i H^k(R\Gamma_P(p_1^{-1} \rho_* F_i)_{P'}) \\ &\simeq \varinjlim_i \rho_* H^k(R\Gamma_P(p_1^{-1} F_i)_{P'}) \\ &\simeq \varinjlim_i \rho_*(H^k(R\Gamma_P(p_1^{-1} F_i)_{P'}))_{Z \times_Z E^*}, \end{aligned}$$

where the last isomorphism follows from Lemma 3.7.6 of [11]. □

Lemma 3.2.3 *Let A and B be two closed subanalytic subsets of E such that $A \cup B = E$, and let $F \in D^b(k_{E_{sa}})$. Then $R\Gamma_A(F_B) \simeq (R\Gamma_A F)_B$.*

Proof. We have a natural arrow $(\Gamma_A F)_B \rightarrow \Gamma_A(F_B)$, and $R(\Gamma_A F)_B \simeq (R\Gamma_A F)_B$ since $(\cdot)_B$ is exact. Then we obtain a morphism $(R\Gamma_A F)_B \rightarrow R\Gamma_A(F_B)$. It is enough to prove that for any $k \in \mathbb{Z}$ and for any $F \in \text{Mod}(k_{E_{sa}})$ we have $(R^k \Gamma_A F)_B \xrightarrow{\sim} R^k \Gamma_A(F_B)$. Since both sides commute with filtrant $\varinjlim$, we may assume $F \in \text{Mod}_{\mathbb{R}-c}^c(k_E)$. Then the result follows from the corresponding one for classical sheaves. □

Proposition 3.2.4 *The two functors $\Phi_{P'}, \Psi_P : D_{\mathbb{R}+}^b(k_{E_{sa}}) \rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}^*})$ are isomorphic.*

Proof. We have the chain of isomorphisms:

$$\begin{aligned}\Phi_{P'} F &= Rp_{2!!}(p_1^{-1} F)_{P'} \\ &\simeq Rp_{2!!} R\Gamma_P((p_1^{-1} F)_{P'}) \\ &\simeq Rp_{2!!}(R\Gamma_P(p_1^{-1} F))_{P'} \\ &\simeq Rp_{2*}(R\Gamma_P(p_1^{-1} F))_{P'} \\ &\simeq Rp_{2*} R\Gamma_P(p_1^{-1} F).\end{aligned}$$

The first isomorphism follows from Lemma 3.1.5 (ii), the second one from Lemma 3.2.3, the third one from Lemma 3.2.2 and the last one from Lemma 3.1.5 (i). $\square$

Definition 3.2.5 *Let $F \in D_{\mathbb{R}+}^b(k_{E_{sa}})$.*

(i) *The Fourier-Sato transform is the functor*

$$\begin{aligned}(\cdot)^\wedge : D_{\mathbb{R}+}^b(k_{E_{sa}}) &\rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}^*}) \\ F^\wedge &= \Phi_{P'} F \simeq \Psi_P F.\end{aligned}$$

(ii) *The inverse Fourier-Sato transform is the functor*

$$\begin{aligned}(\cdot)^\vee : D_{\mathbb{R}+}^b(k_{E_{sa}^*}) &\rightarrow D_{\mathbb{R}+}^b(k_{E_{sa}}) \\ F^\vee &= \Psi_{P'} F \simeq \Phi_P F.\end{aligned}$$

It follows from definition that the functors $^\wedge$ and $^\vee$ commute with $R\rho_*$ and ρ^{-1} . We have quasi-commutative diagrams

$$\begin{array}{ccc} D_{\mathbb{R}+}^b(k_E) & \xrightleftharpoons[\vee]{\wedge} & D_{\mathbb{R}+}^b(k_{E^*}) \\ \downarrow R\rho_* & & \downarrow R\rho_* \\ D_{\mathbb{R}+}^b(k_{E_{sa}}) & \xrightleftharpoons[\vee]{\wedge} & D_{\mathbb{R}+}^b(k_{E_{sa}^*}) \end{array} \quad \begin{array}{ccc} D_{\mathbb{R}+}^b(k_E) & \xrightleftharpoons[\vee]{\wedge} & D_{\mathbb{R}+}^b(k_{E^*}) \\ \uparrow \rho^{-1} & & \uparrow \rho^{-1} \\ D_{\mathbb{R}+}^b(k_{E_{sa}}) & \xrightleftharpoons[\vee]{\wedge} & D_{\mathbb{R}+}^b(k_{E_{sa}^*}). \end{array}$$

This implies that these functors are the extension to conic subanalytic sheaves of the classical Fourier-Sato and inverse Fourier-Sato transforms.

Theorem 3.2.6 *The functors $^{\wedge}$ and $^{\vee}$ are equivalence of categories, inverse to each others. In particular we have*

$$\mathrm{Hom}_{D_{\mathbb{R}^+}^b(k_{E_{sa}})}(F, G) \simeq \mathrm{Hom}_{D_{\mathbb{R}^+}^b(k_{E_{sa}^*})}(F^{\wedge}, G^{\wedge}).$$

Proof. Let $F \in D_{\mathbb{R}^+}^b(k_{E_{sa}})$. The functors $^{\wedge}$ and $^{\vee}$ are adjoint functors, then we have a morphism $F \rightarrow F^{\wedge\vee}$. To show that it induces an isomorphism it is enough to check that $\mathrm{R}\Gamma(U; F) \rightarrow \mathrm{R}\Gamma(U; F^{\wedge\vee})$ is an isomorphism on a basis for the topology of E_{sa} . Hence we may assume that U is $\mathbb{R}^+$ -connected. By Proposition 2.4.10 we may suppose that U is an open subanalytic cone of E . we have the chain of isomorphisms:

$$\begin{aligned} \mathrm{RHom}(k_U, F^{\wedge\vee}) &= \mathrm{RHom}(k_U, \Psi_{P'} \Phi_{P'} F) \\ &\simeq \mathrm{RHom}(\Phi_{P'} k_U, \Phi_{P'} F) \\ &\simeq \mathrm{RHom}(\Phi_{P'} k_U, \Psi_P F) \\ &\simeq \mathrm{RHom}(\Phi_P \Phi_{P'} k_U, F) \\ &\simeq \mathrm{RHom}(k_U, F), \end{aligned}$$

where the last isomorphism follows from Theorem 3.7.9 of [11] and from the fact that the functors $^{\wedge}$ and $^{\vee}$ commute with $R\rho_*$. Similarly we can show that for $G \in D_{\mathbb{R}^+}^b(k_{E_{sa}^*})$ we have an isomorphism $G^{\vee\wedge} \xrightarrow{\sim} G$. $\square$

Remark 3.2.7 *We have seen that the functors $^{\vee}$ and $^{\wedge}$ commute with ρ_* and ρ^{-1} . They do not commute with $\rho_!$ in general since it does not send conic sheaves to conic sheaves. We have the following commutative diagram*

$$\begin{array}{ccc} D_{\mathbb{R}^+}^b(k_E) & \begin{array}{c} \xrightarrow{\wedge} \\ \xleftarrow{\vee} \end{array} & D_{\mathbb{R}^+}^b(k_{E^*}) \\ \downarrow \rho_{\mathbb{R}^+}! & & \downarrow \rho_{\mathbb{R}^+}! \\ D_{\mathbb{R}^+}^b(k_{E_{sa}}) & \begin{array}{c} \xrightarrow{\wedge} \\ \xleftarrow{\vee} \end{array} & D_{\mathbb{R}^+}^b(k_{E_{sa}^*}) \end{array}$$

Let us study some functorial properties of the Fourier-Sato transform. Let Z' be another real analytic manifold and let $f : Z' \rightarrow Z$ be a real analytic map. Set $E' = Z' \times_Z E$ and denote by f_{τ} (resp. f_{π}) the map from E' to E (resp. from E'^* to E^*).

Proposition 3.2.8 *Let $F \in D_{\mathbb{R}+}^b(k_{E'_{sa}})$. Then:*

$$\begin{aligned} (Rf_{\tau*}F)^\wedge &\simeq Rf_{\pi*}(F^\wedge) \\ (Rf_{\tau\mathbb{R}+!!}F)^\wedge &\simeq Rf_{\pi\mathbb{R}+!!}(F^\wedge). \end{aligned}$$

Let $G \in D_{\mathbb{R}+}^b(k_{E_{sa}})$. Then:

$$\begin{aligned} (f_\tau^!G)^\wedge &\simeq f_\pi^!(G^\wedge) \\ (f_\tau^{-1}G)^\wedge &\simeq f_\pi^{-1}(G^\wedge). \end{aligned}$$

Proof. The result follows adapting Proposition 3.7.13 of [11]. $\square$

Let $E_i, i = 1, 2$ be two real vector bundles over Z , $f : E_1 \rightarrow E_2$ a morphism of vector bundles. Set ${}^tf : E_2^* \rightarrow E_1^*$ the dual morphism.

Proposition 3.2.9 (i) *Let $F \in D_{\mathbb{R}+}^b(k_{E_{1sa}})$. Then:*

$$\begin{aligned} {}^tf^!(F^\vee) &\simeq (Rf_*F)^\vee \\ {}^tf^!(F^\wedge) &\simeq (Rf_*F)^\wedge \otimes {}^tf^!k_{E_1^*} \\ {}^tf^{-1}(F^\wedge) &\simeq (Rf_{\mathbb{R}+!!}F)^\wedge \\ {}^tf^{-1}(F^\vee) &\simeq (Rf_{\mathbb{R}+!!}F)^\vee \otimes {}^tf^!k_{E_1^*}. \end{aligned}$$

(ii) *Let $G \in D_{\mathbb{R}+}^b(k_{E_{2sa}})$. Then:*

$$\begin{aligned} (f^!G)^\wedge &\simeq (R{}^tf_*G^\wedge) \\ (f^!k_{E_2} \otimes f^!G)^\vee &\simeq (R{}^tf_*G^\vee) \\ (f^{-1}G)^\vee &\simeq R{}^tf_{\mathbb{R}+!!}G^\vee \\ (f^!k_{E_2} \otimes f^{-1}G)^\wedge &\simeq R{}^tf_{\mathbb{R}+!!}G^\wedge. \end{aligned}$$

Proof. The result follows adapting Proposition 3.7.14 of [11]. $\square$

Let $E_i, i = 1, 2$ be two vector bundles over a real analytic manifold Z . We set for short $E = E_1 \times_Z E_2$ and $E^* = E_1^* \times_Z E_2^*$. We denote by $\wedge$ the Fourier-Sato transform on $E_i, i = 1, 2$ and E .

Proposition 3.2.10 *Let $F_i \in D_{\mathbb{R}+}^b(k_{E_{isa}})$, $i = 1, 2$. There is an isomorphism*

$$F_1^\wedge \boxtimes_Z F_2^\wedge \simeq \left(F_1 \boxtimes_Z F_2 \right)^\wedge.$$

Proof. Let p_i^j and p_i the i -th projection defined on $E_j \times_Z E_j^*$, $j = 1, 2$ and $E \times E^*$ respectively. Let $P'_j = \{\langle x_j, y_j \rangle \leq 0\} \subset E_j \times_Z E_j^*$, $j = 1, 2$ and $P' = \{\langle (x_1, x_2), (y_1, y_2) \rangle \leq 0\} \subset E \times_Z E^*$. The Künneth formula gives rise to the isomorphisms:

$$\begin{aligned} F_1^\wedge \boxtimes_Z F_2^\wedge &\simeq Rp_{2!!} \left(p_1^{1-1} F_1 \boxtimes_Z p_1^{2-1} F_2 \right)_{P'_1 \times_Z P'_2}, \\ \left(F_1 \boxtimes_Z F_2 \right)^\wedge &\simeq Rp_{2!!} \left(p_1^{1-1} F_1 \boxtimes_Z p_1^{2-1} F_2 \right)_{P'}. \end{aligned}$$

It is enough to show that for any sheaf $F \in D^+(k_{(E \times_Z E^*)_{sa}})$ conic with respect to the actions of $\mathbb{R}^+$ on E_j and E_j^* , $j = 1, 2$, the morphism $Rp_{2!!} F_{P'} \rightarrow Rp_{2!!} F_{P'_1 \times_Z P'_2}$ induces an isomorphism $R^k p_{2!!} F_{P'} \xrightarrow{\sim} R^k p_{2!!} F_{P'_1 \times_Z P'_2}$ for any $k \in \mathbb{Z}$. We may reduce to the case F concentrated in degree zero. Then as in § 2.3 one can show that $F = \varinjlim_i \rho_* F_i$, with F_i conic with respect to the actions of $\mathbb{R}^+$ on E_j and E_j^* , $j = 1, 2$, $\mathbb{R}$ -constructible and with compact support on the base for each i . We have the chain of isomorphisms

$$\begin{aligned} R^k p_{2!!} (\varinjlim_i \rho_* F_i)_{P'} &\simeq \varinjlim_i \rho_* R^k p_{2!!} (F_i)_{P'} \\ &\simeq \varinjlim_i \rho_* R^k p_{2!!} (F_i)_{P'_1 \times_Z P'_2} \\ &\simeq R^k p_{2!!} (\varinjlim_i \rho_* F_i)_{P'_1 \times_Z P'_2}. \end{aligned}$$

$R^k p_{2!!}$ commutes with ρ_* by Lemma 3.1.5 and the second isomorphism follows from Proposition 3.7.15 of [11]. $\square$

4 Specialization of subanalytic sheaves

In this section we define specialization for subanalytic sheaves. We refer to [11] for the classical theory of specialization.

4.1 Review on normal deformation

Let X be a real n -dimensional analytic manifold and let M be a closed submanifold of codimension ℓ . As usual we denote by $T_M X \xrightarrow{\tau} M$ the normal bundle.

We follow the notations of [11]. We consider the normal deformation of X , i.e. an analytic manifold $\tilde{X}_M$, an application $(p, t) : \tilde{X}_M \rightarrow X \times \mathbb{R}$, and an action of $\mathbb{R} \setminus \{0\}$ on $\tilde{X}_M$ $(\tilde{x}, r) \mapsto \tilde{x} \cdot r$ satisfying

$$\begin{cases} p^{-1}(X \setminus M) & \text{isomorphic to } (X \setminus M) \times (\mathbb{R} \setminus \{0\}), \\ t^{-1}(c) & \text{isomorphic to } X \text{ for each } c \neq 0, \\ t^{-1}(0) & \text{isomorphic to } T_M X. \end{cases}$$

Let $s : T_M X \hookrightarrow \tilde{X}_M$ be the inclusion, Ω the open subset of $\tilde{X}_M$ defined by $\{t > 0\}$, $i_\Omega : \Omega \hookrightarrow \tilde{X}_M$ and $\tilde{p} = p \circ i_\Omega$. We get a commutative diagram

$$\begin{array}{ccccc} T_M X & \xrightarrow{s} & \tilde{X}_M & \xleftarrow{i_\Omega} & \Omega \\ \downarrow \tau & & \downarrow p & \swarrow \tilde{p} & \\ M & \xrightarrow{i_M} & X & & \end{array}$$

The morphism $\tilde{p}$ is smooth and Ω is homeomorphic to $X \times \mathbb{R}^+$ by the map $(\tilde{p}, t)$.

Definition 4.1.1 *Let S be a subset of X . The normal cone to S along M , denoted by $C_M(S)$, is the closed conic subset of $T_M X$ defined by*

$$C_M(S) = T_M X \cap \overline{\tilde{p}^{-1}(S)}.$$

Let us recall the following result of [11].

Lemma 4.1.2 *Let V be a conic open subset of $T_M X$.*

- (i) *Let W be an open neighborhood of V in $\tilde{X}_M$ and let $U = \tilde{p}(W \cap \Omega)$. Then $V \cap C_M(X \setminus U) = \emptyset$.*
- (ii) *Let U be an open subset of X such that $V \cap C_M(X \setminus U) = \emptyset$. Then $\tilde{p}(U) \cup V$ is an open neighborhood of V in $\Omega \cup T_M X$.*

Lemma 4.1.3 *Let $V \in \text{Op}(T_M X_{sa})$ be conic. Each subanalytic neighborhood W of V in $\tilde{X}_M$ contains a subanalytic neighborhood $\tilde{W}$ satisfying*

$$(4.1) \quad \begin{cases} (i) \text{ the fibers of the map } \tilde{p} : \tilde{W} \cap \Omega \rightarrow X \text{ are connected,} \\ (ii) \tilde{p}(\tilde{W} \cap \Omega) \text{ is subanalytic in } X. \end{cases}$$

Proof. (a) Let $W \in \text{Op}(\tilde{X}_{Msa})$ be a neighborhood of V . Up to shrink W we may suppose $V = W \cap T_M X$. Set $X' = \tilde{X}_M \setminus (M \times \mathbb{R})$, $S = X'/\mathbb{R}^+$. Then $\alpha : X' \rightarrow S$ is an $\mathbb{R}^+$ -bundle and $\varphi : S \rightarrow X$ is proper. Consider a continuous subanalytic section of $T_M X \rightarrow T_M X/\mathbb{R}^+$, extend it to a continuous subanalytic section σ of $X' \rightarrow S$ and set

$$W' = \bigcup_{x \in \alpha(W)} W'_x,$$

where W'_x denotes the connected component of $\alpha^{-1}(x) \cap W$ containing $\sigma(x)$. By construction, the fibers of $W' \rightarrow S$ are connected and W' is an open neighborhood of $V \setminus M$. Let

$$W'' = \bigcup_{x \in M \cap W} W''_x,$$

where W''_x denotes the connected component of $(\{x\} \times \mathbb{R}) \cap W$ intersecting M . Up to shrink W'' , $\widetilde{W} = W' \cup W'' \cup (W \setminus \Omega)$ is an open neighborhood of V and satisfies (i).

(b) Let us see that $\widetilde{W}$ is subanalytic and satisfies (ii). We may reduce to the case $X = \mathbb{R}^n$, $M = \{0\} \times \mathbb{R}^{n-\ell} \subset X$, $\tilde{X}_M = \mathbb{R}^{n+1}$, $T_M X = \mathbb{R}^n \times \{0\} \subset \tilde{X}_M$, $\Omega = \mathbb{R}^n \times \mathbb{R}^+ \subset \tilde{X}_M$. So that $(x', x'', t) \cdot c \mapsto (cx', x'', c^{-1}t)$ is the action of $\mathbb{R}^+$ on $\tilde{X}_M$ and $\tilde{p} : \Omega \rightarrow X$ is the map $(x', x'', t) \mapsto (tx', x'')$. In this situation $X' = \mathbb{R}^{n+1} \setminus (M \times \mathbb{R}) \simeq \mathbb{S}^{\ell-1} \times \mathbb{R}^+ \times \mathbb{R}^{n-\ell} \times \mathbb{R}$. Moreover $X'/\mathbb{R}^+ \simeq \mathbb{S}^{\ell-1} \times \mathbb{R}^{n-\ell} \times \mathbb{R}$, indeed

$$X' = S \times \mathbb{R}^+ = \{(ci(\vartheta), x'', sc^{-1}), (\vartheta, x'', s) \in S, c \in \mathbb{R}^+\},$$

where $i : \mathbb{S}^{n-1} \hookrightarrow \mathbb{R}^n$ denotes the embedding. The section $\sigma : S \xrightarrow{\sim} S \times \{1\} \subset X'$ is a globally subanalytic subset of X' . Let us consider the globally subanalytic (even semialgebraic) homeomorphism $\psi : \Omega \rightarrow \Omega$ defined by $\psi(x', x'', t) = (tx', x'', t^{-1})$. Then $\pi \circ \psi = \tilde{p}$, where $\pi : \mathbb{R}^n \times \mathbb{R}^+ \rightarrow \mathbb{R}^n$ is the projection. The set $\psi(W \cap \Omega)$ is still subanalytic. Let $p \in \mathbb{R}^n$. Then $\pi^{-1}(p) \cap \psi(W \cap X' \cap \Omega)$ is a disjoint union of intervals. Let us consider the interval $(m(p), M(p))$, $m(p) < M(p) \in \mathbb{R} \cup \{\pm\infty\}$ intersecting $\psi(\sigma \cap \Omega)$. Then $\psi(W') = \{(p, r) \in \psi(W \cap X' \cap \Omega); m(p) < r < M(p)\}$. The set $\psi(W' \cap \Omega)$ is open subanalytic (it is a consequence of Proposition 1.2, Chapter 6 of [31]). Moreover, up to shrink W we may suppose that $\tilde{p}(W' \cap \Omega) = \tilde{p}(\sigma \cap W \cap \Omega)$ which is subanalytic. Indeed, since we are working in a local chart, we may assume that $W \cap \sigma$ is globally subanalytic. Let $x'' \in \mathbb{R}^{n-\ell}$. Then $(\{x''\} \times \mathbb{R}) \cap W''$ is a disjoint union of (bounded, up to shrink W) intervals.

Let us consider the interval $(m(x''), M(x''))$, $m(x'') < M(x'') \in \mathbb{R}$ containing 0. Then $W'' = \{r \in W \cap (M \times \mathbb{R}); m(x'') < r < M(x'')\}$. The set W'' is subanalytic (it is a consequence of Proposition 1.2, Chapter 6 of [31]). Moreover (up to shrink W'') $\tilde{p}(W'' \cap \Omega) = W'' \cap M$, which is subanalytic. $\square$

4.2 Specialization of subanalytic sheaves

Definition 4.2.1 *The specialization along M is the functor*

$$\begin{aligned} \nu_M^{sa} : D^b(k_{X_{sa}}) &\rightarrow D^b(k_{T_M X_{sa}}) \\ F &\mapsto s^{-1} \mathrm{R}\Gamma_{\Omega} p^{-1} F. \end{aligned}$$

Theorem 4.2.2 *Let $F \in D^b(k_{X_{sa}})$.*

- (i) $\nu_M^{sa} F \in D_{\mathbb{R}+}^b(k_{T_M X_{sa}})$.
- (ii) *Let V be a conic subanalytic open subset of $T_M X$. Then:*

$$H^j(V; \nu_M^{sa} F) \simeq \varinjlim_U H^j(U; F),$$

where U ranges through the family of $\mathrm{Op}(X_{sa})$ such that $C_M(X \setminus U) \cap V = \emptyset$.

- (iii) *One has the isomorphisms*

$$\begin{aligned} (\nu_M^{sa} F)|_M &\simeq R\tau_*(\nu_M^{sa} F) \simeq F|_M, \\ (\mathrm{R}\Gamma_M \nu_M^{sa} F)|_M &\simeq R\tau_{!!} \nu_M^{sa} F \simeq (\mathrm{R}\Gamma_M F)|_M. \end{aligned}$$

Proof. (i) We may reduce to the case $F \in \mathrm{Mod}(k_{X_{sa}})$. Hence $F = \varinjlim_i \rho_* F_i$ with $F_i \in \mathrm{Mod}_{\mathbb{R}\text{-c}}(k_{X_{sa}})$ for each i . We have $p^{-1} \varinjlim_i \rho_* F_i \simeq \varinjlim_i \rho_* p^{-1} F_i$ and $p^{-1} F_i$ is $\mathbb{R}$ -constructible and conic for each i . Hence $p^{-1} F$ is conic. Since the functors $\mathrm{R}\Gamma_{\Omega}$ and s^{-1} send conic sheaves to conic sheaves we obtain $s^{-1} \mathrm{R}\Gamma_{\Omega} p^{-1} F = \nu_M^{sa} F \in D_{\mathbb{R}+}^b(k_{T_M X_{sa}})$.

(ii) Let $U \in \mathrm{Op}(X_{sa})$ such that $V \cap C_M(X \setminus U) = \emptyset$. We have the chain of morphisms

$$\begin{aligned} \mathrm{R}\Gamma(U; F) &\rightarrow \mathrm{R}\Gamma(p^{-1}(U); p^{-1} F) \\ &\rightarrow \mathrm{R}\Gamma(p^{-1}(U) \cap \Omega; p^{-1} F) \\ &\rightarrow \mathrm{R}\Gamma(\tilde{p}^{-1}(U) \cup V; \mathrm{R}\Gamma_{\Omega} p^{-1} F) \\ &\rightarrow \mathrm{R}\Gamma(V; \nu_M^{sa} F) \end{aligned}$$

where the third arrow exists since $\tilde{p}^{-1}(U) \cup V$ is a neighborhood of V in $\overline{\Omega}$ by Lemma 4.1.2 (ii). Let us show that it is an isomorphism. Let V be a conic open subanalytic subset of $T_M X$. We have

$$\begin{aligned} H^k(V; \nu_M^{sa} F) &\simeq \varinjlim_W H^k(W; R\Gamma_{\Omega} p^{-1} F) \\ &\simeq \varinjlim_W H^k(W \cap \Omega; p^{-1} F), \end{aligned}$$

where W ranges through the family of subanalytic open neighborhoods of V in X . By Lemma 4.1.3 we may assume that W satisfies (4.1). Since $p^{-1} F$ is conic, we have

$$\begin{aligned} H^k(W \cap \Omega; p^{-1} F) &\simeq H^k(p^{-1}(p(W \cap \Omega)); p^{-1} F) \\ &\simeq H^k(p(W \cap \Omega) \times \{1\}; p^{-1} F) \\ &\simeq H^k(p(W \cap \Omega); F), \end{aligned}$$

where the second isomorphism follows since every subanalytic neighborhood of $p(W \cap \Omega) \times \{1\}$ contains an $\mathbb{R}^+$ -connected subanalytic neighborhood (the proof is similar to that of Lemma 4.1.3). By Lemma 4.1.2 (i) we have that $p(W \cap \Omega)$ ranges through the family of subanalytic open subsets U of X such that $V \cap C_M(X \setminus U) = \emptyset$ and we obtain the result.

(iii) The result follows adapting Theorem 4.2.3 (iv) of [11]. □

Proposition 4.2.3 *Let $F \in \text{Mod}(k_{X_{sa}})$ be quasi-injective. Then $\nu_M^{sa} F$ is concentrated in degree zero. Moreover $\nu_M^{sa} F$ is $\mathbb{R}^+$ -quasi-injective.*

Proof. Since $\nu_M^{sa} F$ is conic, it is enough to prove that $H^j(V; \nu_M^{sa} F) = 0$, $j \neq 0$, when V is a conic subanalytic open subset of $T_M X$. By Theorem 4.2.2 we have $H^j(V; \nu_M^{sa} F) \simeq \varinjlim_U H^j(U; F)$, where U ranges through the family of

$\text{Op}(X_{sa})$ such that $C_M(X \setminus U) \cap V = \emptyset$, and $H^j(U; F) = 0$ if $j \neq 0$ since F is quasi-injective. Moreover if $V_1 \subset V_2$ are conic subanalytic open subset of $T_M X$ the morphism

$$\Gamma(V_2; \nu_M^{sa} F) \simeq \varinjlim_{U_1} \Gamma(U_2; F) \rightarrow \varinjlim_{U_2} \Gamma(U_1; F) \simeq \Gamma(V_1; \nu_M^{sa} F),$$

where U_i ranges through the family of $\text{Op}(X_{sa})$ such that $C_M(X \setminus U_i) \cap V_i = \emptyset$, $i = 1, 2$, is surjective. □

Let us study the relation with the classical functor of specialization ν_M .

Proposition 4.2.4 *Let $F \in D^b(k_X)$. Then $\rho^{-1}\nu_M^{sa}R\rho_*F \simeq \nu_M F$.*

Proof. We have to show that for each $x \in T_M X$ we have $H^k(\rho^{-1}\nu_M^{sa}R\rho_*F)_x \simeq H^k(\nu_M F)_x$. Hence we have to prove the isomorphism $\varinjlim_{x \in V} H^k(V; \nu_M^{sa}R\rho_*F) \simeq \varinjlim_{x \in V} H^k(V; \nu_M F)$, where V ranges through the family of open $\mathbb{R}^+$ -connected relatively compact subanalytic subset of $T_M X$. This is a consequence of Theorem 4.2.2 and Theorem 4.2.3 of [11]. $\square$

Remark 4.2.5 *Remark that the functor of specialization does not commute with $R\rho_*$ and ρ^{-1} in general.*

*In fact let $V \in \text{Op}_{sa}^c(T_M X)$ be $\mathbb{R}^+$ -connected and let $F \in \text{Mod}(k_X)$. Then $H^k(V; \nu_M^{sa}R\rho_*F) \simeq \varinjlim_U H^k(U; F)$, where U ranges through the family of $\text{Op}(X_{sa})$ such that $C_M(X \setminus U) \cap V = \emptyset$, which is not cofinal to the family of $\text{Op}(X)$ such that $C_M(X \setminus U) \cap V = \emptyset$.*

Now let $V \in \text{Op}_{sa}^c(T_M X)$ be $\mathbb{R}^+$ -connected and let $G \in \text{Mod}(k_{X_{sa}})$. Then $H^k(V; \nu_M \rho^{-1}G) \simeq \varinjlim_U \varprojlim_{W \subset \subset U} H^k(W; G)$, where U ranges through the family of $\text{Op}(X_{sa})$ such that $C_M(X \setminus U) \cap V = \emptyset$ and $W \in \text{Op}(X_{sa})$. Then $H^k(\rho^{-1}\nu_M^{sa}G)_x \neq H^k(\nu_M \rho^{-1}G)_x$.

Let $f : X \rightarrow Y$ be a morphism of manifolds, let N be a closed submanifold of Y of codimension k and assume $f(M) \subset N$. We denote by f' the map from TX to $X \times_Y TY$ associated to f and by $f_\tau : X \times_Y TY \rightarrow TY$ the base change. We denote by Tf the composite map. Similarly, replacing X, Y, TX, TY by $M, N, T_M X, T_N Y$ we get the morphisms $f'_M, f_{M\tau}, T_M f$.

We have a commuative diagram, where all the squares are cartesian

$$\begin{array}{ccccc} T_M X & \xrightarrow{s_X} & \tilde{X}_M & \xleftarrow{i_{\Omega_X}} & \Omega_X & \xrightarrow{\tilde{p}_X} & X \\ \downarrow T_M f & & \downarrow \tilde{f}' & & \downarrow \tilde{f} & & \downarrow f \\ T_N Y & \xrightarrow{s_Y} & \tilde{Y}_N & \xleftarrow{i_{\Omega_Y}} & \Omega_Y & \xrightarrow{\tilde{p}_Y} & Y \end{array}$$

Recall that the following diagram is not cartesian in general

$$(4.2) \quad \begin{array}{ccc} \tilde{X}_M & \xrightarrow{p_X} & X \\ \downarrow \tilde{f}' & & \downarrow f \\ \tilde{Y}_N & \xrightarrow{p_Y} & Y. \end{array}$$

Definition 4.2.6 (i) One says that f is clean with respect to N if $f^{-1}(N)$ is a submanifold M of X and the map ${}^t f'_M : M \times_N T_N^* Y \rightarrow T_M^* X$ is surjective.

(ii) One says that f is transversal to N if the map ${}^t f'|_{X \times_Y T_N^* Y} : X \times_Y T_N^* Y \rightarrow T^* Y$ is injective.

If f is transversal to N and $f^{-1}(N) = M$, then the square (4.2) is cartesian.

We will not prove the following results, which can be easily recovered adapting §IV.4.2 of [11], using the construction we did for subanalytic sheaves.

Proposition 4.2.7 Let $F \in D^b(k_{X_{sa}})$.

(i) There exists a commutative diagram of canonical morphisms

$$\begin{array}{ccc} R(T_M f)_{\mathbb{R}+!!} \nu_M F & \longrightarrow & \nu_N Rf_{\mathbb{R}+!!} F \\ \downarrow & & \downarrow \\ R(T_M f)_* \nu_M F & \longleftarrow & \nu_N Rf_* F. \end{array}$$

(ii) Moreover if $\text{supp} F \rightarrow Y$ and $C_M(\text{supp} F) \rightarrow T_N Y$ are proper, and if $\text{supp} F \cap f^{-1}(N) \subset M$, then the above morphisms are isomorphisms.

In particular f is clean with respect to N and $f^{-1}(N) = M$, then the above morphisms are isomorphisms.

Proposition 4.2.8 Let $G \in D^b(k_{Y_{sa}})$.

(i) There exists a commutative diagram of canonical morphisms

$$\begin{array}{ccc} \omega_{T_M X/T_N Y} \otimes (T_M f)^{-1} \nu_N G & \longrightarrow & \nu_M (\omega_{X/Y} \otimes f^{-1} G) \\ \downarrow & & \downarrow \\ (T_M f)^! \nu_N G & \longleftarrow & \nu_M f^! G \end{array}$$

- (ii) Moreover if $f : X \rightarrow Y$ and $f|_M : M \rightarrow N$ are smooth the above morphisms are isomorphisms.

Let X and Y be two real analytic manifolds and let M, N be two closed submanifolds of X and Y respectively.

Proposition 4.2.9 *Let $F \in D^b(k_{X_{sa}})$ and $G \in D^b(k_{Y_{sa}})$. There is a natural morphism*

$$\nu_M F \boxtimes \nu_N G \rightarrow \nu_{M \times N}(F \boxtimes G).$$

Corollary 4.2.10 *Let $F, G \in D^b(k_{X_{sa}})$. There is a natural morphism*

$$\nu_M F \otimes \nu_M G \rightarrow \nu_M(F \otimes G).$$

5 Microlocalization of subanalytic sheaves

With the construction of the Fourier-Sato transform and the specialization we have all the tools to define the functor of microlocalization in the framework of subanalytic sites. See [11] for the classical theory of microlocalization. Then we introduce the functor μhom^{sa} for subanalytic sheaves, we study the relations with the notion of microsupport of [15] and with the functor of ind-microlocalization of [16].

5.1 Microlocalization of subanalytic sheaves

Let us denote by $T_M^* X$ the conormal bundle to M in X , i.e. the kernel of the map $M \times_X T^* X \rightarrow T^* M$. We denote by π the projection $T_M^* X \rightarrow M$.

Definition 5.1.1 *Let $F \in D^b(k_{X_{sa}})$. The microlocalization of F along M is the Fourier-Sato transform of the specialization, i.e.*

$$\mu_M^{sa} F = (\nu_M^{sa} F)^\wedge.$$

Theorem 5.1.2 *Let $F \in D^b(k_{X_{sa}})$.*

- (i) $\mu_M^{sa} F \in D_{\mathbb{R}^+}^b(k_{T_M^* X_{sa}})$.
- (ii) Let V be an open convex subanalytic cone of $T_M^* X$. Then:

$$H^j(V; \mu_M^{sa} F) \simeq \varinjlim_{U, Z} H_Z^j(U; F),$$

where U ranges through the family of $\text{Op}(X_{sa})$ such that $U \cap M = \pi(V)$ and Z through the family of closed subanalytic subsets such that $C_M(Z) \subset V^\circ$, where V° denotes the polar cone.

(iii) One has the isomorphisms

$$\begin{aligned} (\mu_M^{sa} F)|_M &\simeq R\pi_*(\mu_M^{sa} F) \simeq i_M^! F, \\ (R\Gamma_M \mu_M^{sa} F)|_M &\simeq R\pi_{!!} \mu_M^{sa} F \simeq i_M^{-1} F \otimes i_M^! k_X. \end{aligned}$$

Proof. The result follows from the functorial properties of the Fourier-Sato transform and Theorem 4.2.2. $\square$

As in classical sheaf theory, we get the Sato's triangle for subanalytic sheaves:

$$F|_M \otimes \omega_{M|X} \rightarrow R\Gamma_M F|_M \rightarrow R\dot{\pi}_* \mu_M^{sa} F \xrightarrow{+}$$

where $\dot{\pi}$ is the restriction of π to $T_M^* X \setminus M$.

Proposition 5.1.3 *Let $F \in \text{Mod}(k_{X_{sa}})$ be quasi-injective. Then $\rho^{-1} \mu_M^{sa} F$ is concentrated in degree zero.*

Proof. The result follows from Theorem 5.1.2 (ii). $\square$

Remark 5.1.4 *Remark that the functor of microlocalization does not commute with $R\rho_*$ and ρ^{-1} since specialization does not. If $F \in D^b(k_X)$ we have $\rho^{-1} \mu_M^{sa} R\rho_* F \simeq \mu_M F$ since the Fourier-Sato transform commutes with ρ^{-1} and $\rho^{-1} \circ \nu_M^{sa} \circ R\rho_* \simeq \nu_M$.*

Let $f : X \rightarrow Y$ be a morphism of manifolds, let N be a closed submanifold of Y of codimension k and assume $f(M) \subset N$. The map Tf defines the maps

$$T^* X \xleftarrow{tf'} X \times_Y T^* Y \xrightarrow{f_\pi} T^* Y$$

and similarly one can define the maps ${}^{tf'}_M$ and $f_{M\pi}$.

Applying the Fourier-Sato transform to the morphisms of § 4.2 we get the following results (see also [11] § IV.4.3 for the classical case)

Proposition 5.1.5 *Let $F \in D^b(k_{X_{sa}})$.*

(i) There exists a commutative diagram of canonical morphisms

$$\begin{array}{ccc}
Rf_{M\pi\mathbb{R}+!!} {}^{tf'_M-1} \mu_M F & \longrightarrow & \mu_N Rf_{!!} F \\
\downarrow & & \downarrow \\
Rf_{M\pi*} ({}^{tf'_M} \omega_{X/Y} \otimes \omega_{M/N}^{\otimes -1}) & \longleftarrow & \mu_N Rf_* F.
\end{array}$$

(ii) Moreover if $\text{supp} F \rightarrow Y$ and $C_M(\text{supp} F) \rightarrow T_N Y$ are proper, and if $\text{supp} F \cap f^{-1}(N) \subset M$, then the above morphisms are isomorphisms.

In particular f is clean with respect to N and $f^{-1}(N) = M$, then the above morphisms are isomorphisms.

Proposition 5.1.6 Let $G \in D^b(k_{Y_{sa}})$.

(i) There exists a commutative diagram of canonical morphisms

$$\begin{array}{ccc}
Rf'_{M\mathbb{R}+!!} (\omega_{M/N} \otimes f_{M\pi}^{-1} \mu_N G) & \longrightarrow & \mu_M (\omega_{X/Y} \otimes f^{-1} G) \\
\downarrow & & \downarrow \\
Rf'_{M*} f'_{M\pi} \mu_N G & \longleftarrow & \mu_M f^! G
\end{array}$$

(ii) Moreover if $f : X \rightarrow Y$ and $f|_M : M \rightarrow N$ are smooth the above morphisms are isomorphisms.

Let X and Y be two real analytic manifolds and let M, N be two closed submanifolds of X and Y respectively.

Proposition 5.1.7 Let $F \in D^b(k_{X_{sa}})$ and $G \in D^b(k_{Y_{sa}})$. There is a natural morphism

$$\mu_M F \boxtimes \mu_N G \rightarrow \mu_{M \times N} (F \boxtimes G).$$

Corollary 5.1.8 Let M be a closed submanifold of X and let $\gamma : T_M^* X \times_M T_M^* X \rightarrow T_M^* X$ be the morphism given by the addition. There is a natural morphism

$$R\gamma_{\mathbb{R}+!!} \left(\mu_M F \boxtimes_M \mu_M G \right) \rightarrow \mu_M (F \boxtimes G) \otimes \omega_{M/X}.$$

5.2 The functor μhom^{sa}

We denote by Δ the diagonal of $X \times X$, and we denote by δ the diagonal embedding. The normal deformation of the diagonal in $X \times X$ can be visualized by the following diagram

$$(5.1) \quad \begin{array}{ccccc} TX & \xrightarrow{\sim} & T_\Delta(X \times X) & \xrightarrow{s} & \widetilde{X \times X} & \xleftarrow{i_\Omega} & \Omega \\ & & \downarrow \tau_X & & \downarrow p & \nearrow \tilde{p} & \\ & & \Delta & \xrightarrow{\delta} & X \times X & & \\ & & \searrow \sim & & \downarrow q_1 \downarrow q_2 & & \\ & & & & X & & \end{array}$$

Set $p_i = q_i \circ p$, $i = 1, 2$. While $\tilde{p}$ and p_i , $i = 1, 2$, are smooth, p is not, and moreover the square is not cartesian.

Definition 5.2.1 Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and $G \in D^b(k_{X_{sa}})$. We set

$$\mu\text{hom}^{sa}(F, G) := \mu_\Delta^{sa} R\mathcal{H}om(q_2^{-1}F, q_1^!G) = (\nu_\Delta^{sa} R\mathcal{H}om(q_2^{-1}F, q_1^!G))^\wedge.$$

Let π denote the projection from $T_\Delta^*(X \times X)$ to $\Delta \simeq X$.

Proposition 5.2.2 Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and $G \in D^b(k_{X_{sa}})$. There is a canonical isomorphism $\pi_* \mu\text{hom}^{sa}(F, G) \simeq R\mathcal{H}om(F, G)$.

Proof. The result follows adapting Proposition 4.4.2 of [11]. □

Remark 5.2.3 The functor μhom^{sa} is well defined also if $F \in D^b(k_{X_{sa}})$. In this case we do not know if $\mu\text{hom}^{sa}(F, G)$ has bounded cohomology or not.

Remark 5.2.4 Adapting the results of § 4.2 and § 5.1 one gets the functorial properties of μhom for subanalytic sheaves. Since the proofs are essentially the same as the classical ones we will skip them and refer to [11].

Let $\pi : T^*X \rightarrow X$ be the projection and consider the canonical 1-form ω , the restriction to the diagonal of the map ${}^t\pi' : T^*X \times_X T^*X \rightarrow T^*T^*X$. We have a diagram

$$\begin{array}{ccccc} T^*T^*X & \xleftarrow{{}^t\pi'} & T^*X \times_X T^*X & \xrightarrow{\pi\pi} & T^*X \\ & \searrow \omega & \uparrow \delta_{T^*X} & \nearrow \sim & \\ & & \Delta_{T^*X} & & \end{array}$$

Lemma 5.2.5 *Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and $G \in D^b(k_{X_{sa}})$. We have*

$$\omega^{-1} \mu hom^{sa}(\pi^{-1}F, \pi^{-1}G) \simeq \mu hom^{sa}(F, G).$$

Proof. We have the isomorphism ${}^t\pi'_! \pi_\pi^{-1} \mu hom^{sa}(F, G) \xrightarrow{\sim} \mu hom^{sa}(\pi^{-1}F, \pi^{-1}G)$. Hence we get the isomorphisms

$$\begin{aligned} \omega^{-1} \mu hom^{sa}(\pi^{-1}F, \pi^{-1}G) &\simeq \omega^{-1} {}^t\pi'_! \pi_\pi^{-1} \mu hom^{sa}(F, G) \\ &\simeq \delta_{T^*X}^{-1} \pi_\pi^{-1} \mu hom^{sa}(F, G) \\ &\simeq \mu hom^{sa}(F, G). \end{aligned}$$

□

5.3 Microlocalization and microsupport

In [11] the authors prove that the support of $\mu hom(F, G)$ is contained in the product of the microsupports of F and G . We extend this result to the functor μhom^{sa} . Let X be a real analytic manifold and let $T^*X \xrightarrow{\pi} X$ be the cotangent bundle. We recall the following two equivalent definitions of microsupport of a subanalytic sheaves of [15]. For the notion of microsupport for classical sheaves we refer to [11]. For the functorial properties of the microsupport of subanalytic sheaves we refer to [21].

Definition 5.3.1 *The microsupport of $F \in D^b(k_{X_{sa}})$, denoted by $SS(F)$ is the subset of T^*X defined as follows. Let $p \in T^*X$, then $p \notin SS(F)$ if one of the following equivalent conditions is satisfied.*

(i) *There exists a conic neighborhood U of p and a small filtrant system $\{F_i\}$ in $C^{[a,b]}(\text{Mod}_{\mathbb{R}\text{-c}}(k_X))$ with $SS(F_i) \cap U = \emptyset$ such that F is quasi-isomorphic to $\varinjlim_i \rho_* F_i$ in a neighborhood of $\pi(p)$.*

(ii) *There exists a conic neighborhood U of p such that for any $G \in D_{\mathbb{R}\text{-c}}^b(k_X)$ with $\text{supp}(G) \subset \subset \pi(U)$ and such that $SS(G) \subset U \cup T_X^*X$, one has $\text{Hom}_{D^b(k_{X_{sa}})}(G, F) = 0$.*

Remark 5.3.2 *In [15] microsupport was defined for ind-sheaves. The above definition follows from the equivalence between subanalytic sheaves and ind- $\mathbb{R}$ -constructible sheaves (see [21] for details).*

We need the following result of [21].

Lemma 5.3.3 *Let X, Y be two real analytic manifolds and let q_1, q_2 be the projections from $X \times Y$ to X and Y respectively. Let $G \in D_{\mathbb{R}\text{-c}}^b(k_Y)$ and $F \in D^b(k_{X_{sa}})$. Then*

$$(5.2) \quad SS(R\mathcal{H}om(q_1^{-1}G, q_2^!F)) \subseteq SS(F) \times SS(G)^a.$$

Let M be a real closed submanifold of X .

Proposition 5.3.4 *Let $F \in D^b(k_{X_{sa}})$. Then $\text{supp}(\mu_M^{sa}F) \subseteq SS(F) \cap T_M^*X$.*

Proof. Let $F \in D^b(k_{X_{sa}})$ and let $p \notin SS(F)$. There exists conic neighborhood U of p and a small filtrant system $\{F_i\}$ in $C^{[a,b]}(\text{Mod}_{\mathbb{R}\text{-c}}(k_X))$ with $SS(F_i) \cap \overline{U} = \emptyset$ such that there exists $W \in \text{Op}(X_{sa})$ with $U \subseteq \pi^{-1}(W)$ and $F_W \simeq \varinjlim_i \rho_* F_i$. We have $H^k \mu_M^{sa} F_W \simeq \varinjlim_i \rho_* H^k \mu_M F_{iW}$, hence $(\mu_M^{sa} F)|_U = 0$ since $\text{supp}(\mu_M F_i) \subseteq SS(F_i)$. $\square$

Corollary 5.3.5 *Let $G \in D_{\mathbb{R}\text{-c}}^b(k_X)$, $F \in D^b(k_{X_{sa}})$. Then*

$$\text{supp}(\mu_{\text{hom}}^{sa}(F, G)) \subseteq SS(F) \cap SS(G).$$

The result follows from Proposition 5.3.4 and (5.2). $\square$

Let $f : X \rightarrow Y$ be a morphism of real analytic manifolds and denote by $f_\pi : X \times_Y T^*Y \rightarrow T^*Y$ the base change map.

Definition 5.3.6 *Let $f : X \rightarrow Y$ be a morphism of real analytic manifolds and let $F \in D^b(k_{Y_{sa}})$. One says that f is non characteristic for $SS(F)$ if*

$$f_\pi^{-1}(SS(F)) \cap T_X^*Y \subseteq X \times_Y T_Y^*Y.$$

If f is a closed embedding X is said to be non characteristic.

Proposition 5.3.7 *Let $f : X \rightarrow Y$ be a morphism of real analytic manifolds and let $F \in D^b(k_{Y_{sa}})$. Assume that f is non characteristic for $SS(F)$. Then the natural morphism*

$$f^{-1}F \otimes \omega_{X|Y} \rightarrow f^!F$$

is an isomorphism.

Proof. We may reduce to the case f closed embedding, hence we have to prove the isomorphism $F|_X \otimes \omega_{X|Y} \simeq R\Gamma_X F|_X$ when $SS(F) \cap T_X^* Y \subseteq T_Y^* Y$. Consider the Sato's triangle

$$F|_X \otimes \omega_{X|Y} \rightarrow R\Gamma_X F|_X \rightarrow R\dot{\pi}_* \mu_X^{sa} F \xrightarrow{+}.$$

Since $SS(F) \cap T_X^* Y \subseteq T_Y^* Y$ we have $R\dot{\pi}_* \mu_X^{sa} F = 0$ by Proposition 5.3.4 and the result follows. $\square$

Lemma 5.3.8 *Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and let $G \in D^b(k_{X_{sa}})$. Then*

$$D'F \boxtimes G \xrightarrow{\sim} R\mathcal{H}om(q_1^{-1}F, q_2^{-1}G).$$

Proof. We may reduce to the case $F = k_U$ with $D'k_U \simeq k_{\overline{U}}$ and G quasi-injective. Set $G = \varinjlim_i \rho_* G_i$ with $G_i \in \text{Mod}_{\mathbb{R}\text{-c}}(k_X)$, we have the chain of isomorphisms

$$(q_2^{-1}G)_{q_1^{-1}(\overline{U})} \simeq \varinjlim_i \rho_*(q_2^{-1}G_i)_{q_1^{-1}(\overline{U})} \simeq \varinjlim_i \rho_* \Gamma_{q_1^{-1}(U)}(q_2^{-1}G_i) \simeq \Gamma_{q_1^{-1}(U)}(q_2^{-1}G)$$

where the second isomorphism follows from Proposition 3.4.4 of [11]. $\square$

Proposition 5.3.9 *Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and let $G \in D^b(k_{X_{sa}})$. Suppose that $SS(F) \cap SS(G) \subseteq T_X^* X$. Then*

$$D'F \otimes G \xrightarrow{\sim} R\mathcal{H}om(F, G).$$

Proof. Let $\delta : \Delta \rightarrow X \times X$ be the embedding and let us consider the Sato's triangle

$$\begin{aligned} \delta^{-1} R\mathcal{H}om(q_1^{-1}F, q_2^!G) \otimes \omega_{\Delta|X \times X} &\rightarrow \delta^! R\mathcal{H}om(q_1^{-1}F, q_2^!G) \\ &\rightarrow R\dot{\pi}_* \mu hom^{sa}(F, G) \xrightarrow{+}. \end{aligned}$$

We have $\delta^! R\mathcal{H}om(q_1^{-1}F, q_2^!G) \simeq R\mathcal{H}om(F, G)$. Moreover

$$\begin{aligned} \delta^{-1} R\mathcal{H}om(q_1^{-1}F, q_2^!G) \otimes \omega_{\Delta|X \times X} &\simeq \delta^{-1} R\mathcal{H}om(q_1^{-1}F, q_2^{-1}G) \\ &\simeq \delta^{-1}(D'F \boxtimes G) \\ &\simeq D'F \otimes G \end{aligned}$$

where the second isomorphism follows from Lemma 5.3.8. Then we obtain a distinguished triangle

$$D'F \otimes G \rightarrow R\mathcal{H}om(F, G) \rightarrow R\dot{\pi}_* \mu hom^{sa}(F, G) \xrightarrow{+}$$

and the result follows since $R\dot{\pi}_* \mu hom^{sa}(F, G) = 0$ by Corollary 5.3.5. $\square$

5.4 The link with the functor μ of microlocalization

We will study the relation between microlocalization for subanalytic sheaves and the functor μ of [16]. We will need first some results. Let X be a real analytic manifold and consider the normal deformation of Δ in $X \times X$ visualized by the diagram (5.1).

Lemma 5.4.1 *Let $G \in D^b(k_{(X \times X)_{sa}})$ and let $F \in D_{\mathbb{R}\text{-c}}^b(k_{(X \times X)_{sa}})$. We have $R\Gamma_{\Omega} p^{-1} R\mathcal{H}om(F, G) \simeq R\Gamma_{\Omega} R\mathcal{H}om(p^{-1}F, p^{-1}G)$.*

Proof. We may reduce to the case $F \in \text{Mod}_{\mathbb{R}\text{-c}}(k_{X \times X})$ and $G \in \text{Mod}(k_{(X \times X)_{sa}})$. Hence $G = \varinjlim_i \rho_* G_i$ with $G_i \in \text{Mod}_{\mathbb{R}\text{-c}}(k_{(X \times X)_{sa}})$ for each i . Let $\widetilde{\Omega \times X_{sa}}$ be

the site induced by $\widetilde{X \times X_{sa}}$ on Ω . We have

$$\begin{aligned} H^k i_{\Omega \times \widetilde{X_{sa}}}^{-1} p^{-1} R\mathcal{H}om(F, \varinjlim_i \rho_* G_i) &\simeq \varinjlim_i \rho_* H^k i_{\Omega}^{-1} p^{-1} R\mathcal{H}om(F, G_i) \\ &\simeq \varinjlim_i \rho_* H^k R\mathcal{H}om(i_{\Omega}^{-1} p^{-1} F, i_{\Omega}^{-1} p^{-1} G_i) \\ &\simeq \varinjlim_i \rho_* H^k i_{\Omega}^{-1} R\mathcal{H}om(p^{-1} F, p^{-1} G_i) \\ &\simeq H^k i_{\Omega \times \widetilde{X_{sa}}}^{-1} R\mathcal{H}om(p^{-1} F, \varinjlim_i \rho_* p^{-1} G_i). \end{aligned}$$

The second isomorphism follows because $p \circ i_{\Omega}$ is smooth. Composing with $Ri_{\Omega \times \widetilde{X_{sa}}}^*$ we obtain the result. $\square$

Lemma 5.4.2 *Let $G \in D^b(k_{X_{sa}})$, then for $i = 1, 2$*

$$(5.3) \quad k_{\overline{\Omega}} \otimes p_i^{-1} G \simeq R\mathcal{H}om(k_{\Omega}, p_i^{-1} G)$$

$$(5.4) \quad k_{\Omega} \otimes p_i^{-1} G \simeq R\mathcal{H}om(k_{\overline{\Omega}}, p_i^{-1} G).$$

Proof. Let us prove (5.3). Since for $i = 1, 2$ p_i is smooth, Proposition 3.16 of [21] implies that $SS(p_i^{-1}G) \cap SS(k_\Omega)$ is contained on the zero section of $T^*(\widetilde{X \times X})$. Then the result is a consequence of Proposition 5.3.9, and the fact that $D'k_\Omega \simeq k_{\overline{\Omega}}$. The proof of (5.4) is similar. $\square$

Let σ be a section of $T^*X \rightarrow X$ and consider the following commutative diagram with cartesian square

$$(5.5) \quad \begin{array}{ccccc} & & TX & & \\ & \pi_2 \uparrow & \swarrow \text{id} & & \\ T^*X \times_X TX & \xleftarrow{\sigma'} & TX & & \\ \pi_1 \downarrow & & \downarrow \tau_X & & \\ T^*X & \xleftarrow{\sigma} & X & & \end{array}$$

We set

$$\begin{aligned} P &:= \{((x, \xi), (x, v)) \in T^*X \times_X TX; \langle \xi, v \rangle \geq 0\} \\ P' &:= \{((x, \xi), (x, v)) \in T^*X \times_X TX; \langle \xi, v \rangle \leq 0\} \\ P_\sigma &:= \{(x, v) \in TX; \langle \sigma(x), v \rangle \geq 0\} = \sigma'^{-1}(P) \\ P'_\sigma &:= \{(x, v) \in TX; \langle \sigma(x), v \rangle \leq 0\} = \sigma'^{-1}(P') \end{aligned}$$

The kernel K_σ is defined as follows

$$(5.6) \quad K_\sigma := Rp_{!!}(k_{\overline{\Omega}} \otimes \rho_! k_{P_\sigma}) \otimes \rho_! \delta_* \omega_{\Delta|X \times X}^{\otimes -1}.$$

Proposition 5.4.3 (i) Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and $G \in D^b(k_{X_{sa}})$. There is a natural arrow

$$\varphi : R\mathcal{H}om(F, K_\sigma \circ G) \rightarrow \sigma^{-1} \mu hom^{sa}(F, G),$$

where $K_\sigma \circ G$ means $Rq_{1!!}(q_2^{-1}G \otimes K_\sigma)$.

(ii) Let $\rho : X \rightarrow X_{sa}$ be the natural functor of sites. Then $\rho^{-1}(\varphi)$ is an isomorphism.

Proof. (i)_a We have the chain of isomorphisms

$$\begin{aligned} \sigma^{-1} \mu hom^{sa}(F, G) &\simeq \sigma^{-1} R\pi_{1!!}(\pi_2^{-1} \nu_\Delta^{sa} R\mathcal{H}om(q_2^{-1}F, q_1^!G) \otimes k_{P'}) \\ &\simeq R\tau_{X!!} \sigma'^{-1}(\pi_2^{-1} \nu_\Delta^{sa} R\mathcal{H}om(q_2^{-1}F, q_1^!G) \otimes k_{P'}) \\ &\simeq R\tau_{X!!}(\nu_\Delta^{sa} R\mathcal{H}om(q_2^{-1}F, q_1^!G) \otimes k_{P'_\sigma}). \end{aligned}$$

Consider the normal deformation of Δ in $X \times X$ visualized by the diagram (5.1). We have

$$\begin{aligned}
\nu_{\Delta}^{sa} R\mathcal{H}om(q_2^{-1}F, q_1^!G) &\simeq s^{-1}R\Gamma_{\Omega}p^{-1}R\mathcal{H}om(q_2^{-1}F, q_1^!G) \\
&\simeq s^{-1}R\mathcal{H}om(p_2^{-1}F, R\Gamma_{\Omega}p^{-1}q_1^!G) \\
&\simeq s^{-1}R\mathcal{H}om(p_2^{-1}F, R\Gamma_{\Omega}p_1^{-1}G) \otimes \tau_X^{-1}\omega_X \\
&\simeq s^{-1}R\mathcal{H}om(p_2^{-1}F, R\Gamma_{\Omega}p_1^{-1}G) \otimes \tau_X^{-1}\omega_{\Delta|X \times X}^{\otimes -1} \\
&\simeq s^{-1}R\mathcal{H}om(p_2^{-1}F, p_1^{-1}G \otimes k_{\overline{\Omega}}) \otimes \tau_X^{-1}\omega_{\Delta|X \times X}^{\otimes -1},
\end{aligned}$$

where the second isomorphism follows from Lemma 5.4.1 and the last one follows from Lemma 5.4.2. Hence we get

$$\begin{aligned}
\sigma^{-1}\mu hom^{sa}(F, G) &\simeq R\tau_{X!!}(s^{-1}R\mathcal{H}om(p_2^{-1}F, p_1^{-1}G \otimes k_{\overline{\Omega}}) \otimes \tau_X^{-1}\omega_{\Delta|X \times X}^{\otimes -1} \otimes k_{P'_\sigma}) \\
&\simeq Rp_{2!!}s!!(s^{-1}R\mathcal{H}om(p_2^{-1}F, p_1^{-1}G \otimes k_{\overline{\Omega}}) \otimes \tau_X^{-1}\omega_{\Delta|X \times X}^{\otimes -1} \otimes k_{P'_\sigma}) \\
&\simeq Rp_{2!!}(R\mathcal{H}om(p_2^{-1}F, p_1^{-1}G \otimes k_{\overline{\Omega}}) \otimes p^{-1}\delta_*\omega_{\Delta|X \times X}^{\otimes -1} \otimes k_{P'_\sigma}).
\end{aligned}$$

(i)_b On the other hand we have the chain of isomorphisms

$$\begin{aligned}
K_{\sigma} \circ G &\simeq Rq_{1!!}(q_2^{-1}G \otimes Rp_{1!!}(k_{\overline{\Omega}} \otimes \rho_!k_{P_{\sigma}}) \otimes \rho_!\delta_*\omega_{\Delta|X \times X}^{\otimes -1}) \\
&\simeq Rp_{1!!}(p_2^{-1}G \otimes k_{\overline{\Omega}} \otimes \rho_!k_{P_{\sigma}} \otimes p^{-1}\rho_!\delta_*\omega_{\Delta|X \times X}^{\otimes -1}) \\
&\simeq Rp_{2!!}(p_1^{-1}G \otimes k_{\overline{\Omega}} \otimes \rho_!k_{P'_\sigma} \otimes p^{-1}\rho_!\delta_*\omega_{\Delta|X \times X}^{\otimes -1}) \\
&\simeq Rp_{2!!}(p_1^{-1}G \otimes k_{\overline{\Omega}} \otimes \rho_!(k_{P'_\sigma} \otimes p^{-1}\delta_*\omega_{\Delta|X \times X}^{\otimes -1})).
\end{aligned}$$

Hence we get

$$\begin{aligned}
R\mathcal{H}om(F, K_{\sigma} \circ G) &\simeq R\mathcal{H}om(F, Rp_{2!!}(p_1^{-1}G \otimes k_{\overline{\Omega}} \otimes \rho_!(k_{P'_\sigma} \otimes p^{-1}\delta_*\omega_{\Delta|X \times X}^{\otimes -1}))) \\
&\simeq Rp_{2!!}R\mathcal{H}om(p_2^{-1}F, p_1^{-1}G \otimes k_{\overline{\Omega}} \otimes \rho_!(k_{P'_\sigma} \otimes p^{-1}\delta_*\omega_{\Delta|X \times X}^{\otimes -1})) \\
&\simeq Rp_{2!!}(R\mathcal{H}om(p_2^{-1}F, p_1^{-1}G \otimes k_{\overline{\Omega}}) \otimes \rho_!(k_{P'_\sigma} \otimes p^{-1}\delta_*\omega_{\Delta|X \times X}^{\otimes -1})).
\end{aligned}$$

(i)_c The adjunction morphism defines a morphism $\rho_! \rightarrow \rho_*$. It induces the morphism

$$\varphi : R\mathcal{H}om(F, K_{\sigma} \circ G) \rightarrow \sigma^{-1}\mu hom^{sa}(F, G).$$

(ii) Composing with ρ^{-1} we get $\rho^{-1} \circ \rho_! \xrightarrow{\sim} \rho^{-1} \circ \rho_* \simeq \text{id}$. Hence we get

$$\rho^{-1}(\varphi) : \rho^{-1}R\mathcal{H}om(F, K_{\sigma} \circ G) \xrightarrow{\sim} \rho^{-1}\sigma^{-1}\mu hom^{sa}(F, G).$$

□

Let $\pi : T^*X \rightarrow X$ be the projection and consider the canonical 1-form ω , the restriction to the diagonal of the map ${}^t\pi'$. Replace X with T^*X and σ with ω in (5.6) and consider the microlocal kernel

$$K_{T^*X} = Rp_{!!}(k_{\overline{\Omega}} \otimes \rho_! k_{P_\omega}) \otimes \rho_! \delta_* \omega_{\Delta_{T^*X}|T^*X \times T^*X}^{\otimes -1}$$

Definition 5.4.4 *The functor of microlocalization of [16] is defined as*

$$\begin{aligned} \mu : D^b(k_{X_{sa}}) &\rightarrow D^b(k_{T^*X_{sa}}) \\ F &\mapsto \mu F = K_{T^*X} \circ \pi^{-1} F \end{aligned}$$

Theorem 5.4.5 (i) *Let $F \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and $G \in D^b(k_{X_{sa}})$. There is a natural arrow*

$$(5.7) \quad \varphi : R\mathcal{H}om(\pi^{-1}F, \mu G) \rightarrow \mu hom^{sa}(F, G).$$

(ii) *Let $\rho : T^*X \rightarrow T^*X_{sa}$ be the natural functor of sites. Then $\rho^{-1}(\varphi)$ is an isomorphism.*

Proof. (i) By Lemma 5.2.5 and Proposition 5.4.3 (i) we get the morphisms

$$\mu hom^{sa}(F, G) \simeq \omega^{-1} \mu hom^{sa}(\pi^{-1}F, \pi^{-1}G) \leftarrow R\mathcal{H}om(\pi^{-1}F, \mu G).$$

(ii) The result follows from Proposition 5.4.3 (ii). □

Example 5.4.6 *The morphism (5.7) is not an isomorphism in general. For example let $F \in \text{Mod}(k_{X_{sa}})$. Then*

$$R\mathcal{H}om(\pi^{-1}k_X, \mu F) \simeq R\mathcal{H}om(k_{T^*X}, \mu F) \simeq \mu F,$$

on the other hand we have

$$\mu hom^{sa}(k_X, F) \simeq j_* \mu_X F \simeq j_* F,$$

*where $j : T_X^*X \hookrightarrow T^*X$ denotes the embedding of the zero section.*

6 Holomorphic functions with growth conditions

In this section we show how the functors we defined before generalize classical constructions.

In § 6.3 we show the relation between specialization of Whitney holomorphic functions with the functor of formal specialization of [6], and the sheaf of asymptotically developable functions of [20] and [29].

In § 6.4 we study the microlocalization of tempered and Whitney holomorphic functions. We establish a relation with the functors of tempered and formal microlocalization introduced by Andronikof in [1] and Colin in [5].

6.1 Review on temperate and formal cohomology

From now on, the base field is $\mathbb{C}$. Let M be a real analytic manifold. One denotes by $\mathcal{D}b_M$ the sheaf of Schwartz's distributions, by $\mathcal{C}_M^\infty$ the sheaf of $\mathcal{C}^\infty$ -functions, and by $\mathcal{D}_M$ the sheaf of finite order differential operators with analytic coefficients. As usual, for $F \in \text{Mod}(\mathbb{C}_M)$, we set $D'F = R\mathcal{H}om(F, \mathbb{C}_M)$.

In [9] the author defined the functor

$$T\mathcal{H}om(\cdot, \mathcal{D}b_M) : \text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_M) \rightarrow \text{Mod}(\mathcal{D}_M)$$

in the following way: let U be a subanalytic subset of M and $Z = M \setminus U$. Then the sheaf $T\mathcal{H}om(\mathbb{C}_U, \mathcal{D}b_M)$ is defined by the exact sequence

$$0 \rightarrow \Gamma_Z \mathcal{D}b_M \rightarrow \mathcal{D}b_M \rightarrow T\mathcal{H}om(\mathbb{C}_U, \mathcal{D}b_M) \rightarrow 0.$$

This functor is exact and extends as functor in the derived category, from $D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$ to $D^b(\mathcal{D}_M)$. Moreover the sheaf $T\mathcal{H}om(F, \mathcal{D}b_M)$ is soft for any $\mathbb{R}$ -constructible sheaf F .

Definition 6.1.1 *Let Z be a closed subset of M . We denote by $\mathcal{I}_{M,Z}^\infty$ the sheaf of $\mathcal{C}^\infty$ -functions on M vanishing up to infinite order on Z .*

Definition 6.1.2 *A Whitney function on a closed subset Z of M is an indexed family $F = (F^k)_{k \in \mathbb{N}^n}$ consisting of continuous functions on Z such that $\forall m \in \mathbb{N}, \forall k \in \mathbb{N}^n, |k| \leq m, \forall x \in Z, \forall \varepsilon > 0$ there exists a neighborhood U of x such that $\forall y, z \in U \cap Z$*

$$\left| F^k(z) - \sum_{|j+k| \leq m} \frac{(z-y)^j}{j!} F^{j+k}(y) \right| \leq \varepsilon d(y, z)^{m-|k|}.$$

We denote by $W_{M,Z}^\infty$ the space of Whitney $\mathcal{C}^\infty$ -functions on Z . We denote by $\mathcal{W}_{M,Z}^\infty$ the sheaf $U \mapsto W_{U,U \cap Z}^\infty$.

In [13] the authors defined the functor

$$\cdot \otimes_M^w \mathcal{C}_M^\infty : \text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_M) \rightarrow \text{Mod}(\mathcal{D}_M)$$

in the following way: let U be a subanalytic open subset of M and $Z = M \setminus U$. Then $\mathbb{C}_U \otimes_M^w \mathcal{C}_M^\infty = \mathcal{I}_{M,Z}^\infty$, and $\mathbb{C}_Z \otimes_M^w \mathcal{C}_M^\infty = \mathcal{W}_{M,Z}^\infty$. This functor is exact and extends as a functor in the derived category, from $D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$ to $D^b(\mathcal{D}_M)$. Moreover the sheaf $F \otimes_M^w \mathcal{C}_M^\infty$ is soft for any $\mathbb{R}$ -constructible sheaf F .

Now let X be a complex manifold, $X_{\mathbb{R}}$ the underlying real analytic manifold and $\overline{X}$ the complex conjugate manifold. The product $X \times \overline{X}$ is a complexification of $X_{\mathbb{R}}$ by the diagonal embedding $X_{\mathbb{R}} \hookrightarrow X \times \overline{X}$. One denotes by $\mathcal{O}_X$ the sheaf of holomorphic functions and by $\mathcal{D}_X$ the sheaf of finite order differential operators with holomorphic coefficients. For $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ one sets

$$\begin{aligned} T\mathcal{H}om(F, \mathcal{O}_X) &= R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, T\mathcal{H}om(F, \mathcal{D}b_{X_{\mathbb{R}}})) , \\ F \otimes_X^w \mathcal{O}_X &= R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, F \otimes_{X_{\mathbb{R}}}^w \mathcal{C}_{X_{\mathbb{R}}}^\infty), \end{aligned}$$

and these functors are called the functors of temperate and formal cohomology respectively.

6.2 Tempered and Whitney holomorphic functions

Definition 6.2.1 *One denotes by $\mathcal{D}b_M^t$ the presheaf of tempered distributions on M_{sa} defined as follows:*

$$U \mapsto \Gamma(M; \mathcal{D}b_M) / \Gamma_{M \setminus U}(M; \mathcal{D}b_M).$$

As a consequence of the Łojasiewicz's inequalities [17], for $U, V \in \text{Op}(M_{sa})$ the sequence

$$0 \rightarrow \mathcal{D}b_M^t(U \cup V) \rightarrow \mathcal{D}b_M^t(U) \oplus \mathcal{D}b_M^t(V) \rightarrow \mathcal{D}b_M^t(U \cap V) \rightarrow 0$$

is exact. Then $\mathcal{D}b_M^t$ is a sheaf on M_{sa} . Moreover, by definition $\mathcal{D}b_M^t$ is quasi-injective.

Definition 6.2.2 One denotes by $\mathcal{C}_M^{\infty, w}$ the presheaf of Whitney $\mathcal{C}^\infty$ -functions on M_{sa} defined as follows:

$$U \mapsto \Gamma(M; H^0 D' \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_M^\infty).$$

As a consequence of a result of [19], for $U, V \in \text{Op}(M_{sa})$ the sequence

$$0 \rightarrow \mathcal{C}_M^{\infty, w}(U \cup V) \rightarrow \mathcal{C}_M^{\infty, w}(U) \oplus \mathcal{C}_M^{\infty, w}(V) \rightarrow \mathcal{C}_M^{\infty, w}(U \cap V)$$

is exact. Then $\mathcal{C}_M^{\infty, w}$ is a sheaf on M_{sa} . Moreover if $U \in \text{Op}(M_{sa})$ is locally cohomologically trivial (l.c.t. for short), i.e. if $D' \mathbb{C}_U \simeq \mathbb{C}_{\overline{U}}$, the morphism $\Gamma(X; \mathcal{C}_M^{\infty, w}) \rightarrow \Gamma(U; \mathcal{C}_M^{\infty, w})$ is surjective and $R\Gamma(U; \mathcal{C}_M^{\infty, w})$ is concentrated in degree zero.

We have the following result (see [14], [23]).

Proposition 6.2.3 For each $F \in \text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_M)$ one has the isomorphisms

$$\begin{aligned} \rho^{-1} R\mathcal{H}om(F, \mathcal{D}b_M^t) &\simeq T\mathcal{H}om(F, \mathcal{D}b_M), \\ \rho^{-1} R\mathcal{H}om(F, \mathcal{C}_M^{\infty, w}) &\simeq D' F \overset{w}{\otimes} \mathcal{C}_M^\infty. \end{aligned}$$

Now let X be a complex manifold, $X_{\mathbb{R}}$ the underlying real analytic manifold and $\overline{X}$ the complex conjugate manifold. One denotes by $\mathcal{O}_X^t$ and $\mathcal{O}_X^w$ the sheaves of tempered and Whitney holomorphic functions defined as follows:

$$\begin{aligned} \mathcal{O}_X^t &:= R\mathcal{H}om_{\rho! \mathcal{D}_{\overline{X}}}(\rho! \mathcal{O}_{\overline{X}}, \mathcal{D}b_{X_{\mathbb{R}}}^t), \\ \mathcal{O}_X^w &:= R\mathcal{H}om_{\rho! \mathcal{D}_{\overline{X}}}(\rho! \mathcal{O}_{\overline{X}}, \mathcal{C}_{X_{\mathbb{R}}}^{\infty, w}). \end{aligned}$$

The relation with the functors of temperate and formal cohomology are given by the following result (see [14], [23])

Proposition 6.2.4 For each $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ one has the isomorphisms

$$\begin{aligned} T\mathcal{H}om(F, \mathcal{O}_X) &\simeq \rho^{-1} R\mathcal{H}om(F, \mathcal{O}_X^t), \\ D' F \overset{w}{\otimes} \mathcal{O}_X &\simeq \rho^{-1} R\mathcal{H}om(F, \mathcal{O}_X^w). \end{aligned}$$

6.3 Asymptotic expansions

Let M be a real analytic manifold. We consider a slight generalization of the sheaf of Whitney $\mathcal{C}^\infty$ -functions of [14].

Definition 6.3.1 Let $F \in \text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_M)$ and let $U \in \text{Op}(M_{sa})$. We define the presheaf $\mathcal{C}_{M|F}^{\infty, w}$ as follows:

$$U \mapsto \Gamma(M; H^0 D' \mathbb{C}_U \otimes F \overset{w}{\otimes} \mathcal{C}_M^\infty).$$

Let $U, V \in \text{Op}(M_{sa})$, and consider the exact sequence

$$0 \rightarrow \mathbb{C}_{U \cap V} \rightarrow \mathbb{C}_U \oplus \mathbb{C}_V \rightarrow \mathbb{C}_{U \cup V} \rightarrow 0,$$

applying the functor $\mathcal{H}om(\cdot, \mathbb{C}_M) = H^0 D'(\cdot)$ we obtain

$$0 \rightarrow H^0 D' \mathbb{C}_{U \cap V} \rightarrow H^0 D' \mathbb{C}_U \oplus H^0 D' \mathbb{C}_V \rightarrow H^0 D' \mathbb{C}_{U \cup V},$$

applying the exact functors $\cdot \otimes F, \cdot \overset{w}{\otimes} \mathcal{C}_M^\infty$ and taking global sections we obtain

$$0 \rightarrow \mathcal{C}_{M|F}^{\infty, w}(U \cup V) \rightarrow \mathcal{C}_{M|F}^{\infty, w}(U) \oplus \mathcal{C}_{M|F}^{\infty, w}(V) \rightarrow \mathcal{C}_{M|F}^{\infty, w}(U \cap V).$$

This implies that $\mathcal{C}_{M|F}^{\infty, w}$ is a sheaf on M_{sa} . Moreover if $U \in \text{Op}(M_{sa})$ is l.c.t., the morphism $\Gamma(X; \mathcal{C}_M^{\infty, w}) \rightarrow \Gamma(U; \mathcal{C}_{M|F}^{\infty, w})$ is surjective and $R\Gamma(U; \mathcal{C}_{M|F}^{\infty, w})$ is concentrated in degree zero. Let $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$ be an exact sequence in $\text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_M)$, we obtain an exact sequence in $\text{Mod}(\mathbb{C}_{M_{sa}})$

$$(6.1) \quad 0 \rightarrow \mathcal{C}_{M|F}^{\infty, w} \rightarrow \mathcal{C}_{M|G}^{\infty, w} \rightarrow \mathcal{C}_{M|H}^{\infty, w} \rightarrow 0.$$

We can easily extend the sheaf $\mathcal{C}_{M|F}^{\infty, w}$ to the case of $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$, taking a finite resolution of F consisting of locally finite sums $\oplus \mathbb{C}_V$, with V l.c.t. in $\text{Op}^c(M_{sa})$. In fact, the sheaves $\mathcal{C}_{M| \oplus \mathbb{C}_V}^{\infty, w}$ form a complex quasi-isomorphic to $\mathcal{C}_{M|F}^{\infty, w}$ consisting of acyclic objects with respect to $\Gamma(U; \cdot)$, where U is l.c.t. in $\text{Op}^c(M_{sa})$.

As in the case of Whitney $\mathcal{C}^\infty$ -functions one can prove that, if $G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$ one has

$$\rho^{-1} R\mathcal{H}om(G, \mathcal{C}_{F|M}^{\infty, w}) \simeq D'G \otimes F \overset{w}{\otimes} \mathcal{C}_M^\infty.$$

Example 6.3.2 Setting $F = \mathbb{C}_M$ we obtain the sheaf of Whitney $\mathcal{C}^\infty$ -functions. Let N be a closed analytic submanifold of M . Then $\mathcal{C}_{M|\mathbb{C}_{M \setminus N}}^{\infty, w}$ is the sheaf of Whitney $\mathcal{C}^\infty$ -functions vanishing on N with all their derivatives.

Notations 6.3.3 Let Z be a locally closed subanalytic subset of M . We set for short $\mathcal{C}_{M|Z}^{\infty, w}$ instead of $\mathcal{C}_{M|\mathbb{C}_Z}^{\infty, w}$.

Let N be a closed analytic submanifold of M , let $T_N M \xrightarrow{\tau} N$ be the normal vector bundle and consider the normal deformation $\widetilde{M}_N$ as in § 4.1.

Set $F = \mathbb{C}_{M \setminus N}$, $G = \mathbb{C}_M$, $H = \mathbb{C}_N$ in (6.1). The exact sequence

$$0 \rightarrow \mathcal{C}_{M|M \setminus N}^{\infty, w} \rightarrow \mathcal{C}_M^{\infty, w} \rightarrow \mathcal{C}_{M|N}^{\infty, w} \rightarrow 0$$

induces an exact sequence

$$0 \rightarrow \nu_N^{sa} \mathcal{C}_{M|M \setminus N}^{\infty, w} \rightarrow \nu_N^{sa} \mathcal{C}_M^{\infty, w} \rightarrow \nu_N^{sa} \mathcal{C}_{M|N}^{\infty, w} \rightarrow 0,$$

in fact let V be a l.c.t. conic subanalytic subset of $\widetilde{M}_N$ and $U \in \text{Op}(M_{sa})$ such that $C_M(X \setminus U) \cap V = \emptyset$, then we can find a l.c.t. $U' \subset U$ satisfying the same property. Moreover it is easy to see that $\nu_N^{sa} \mathcal{C}_{M|N}^{\infty, w} \simeq \tau^{-1} \mathcal{C}_{M|N}^{\infty, w}$, hence we get the exact sequence

$$(6.2) \quad 0 \rightarrow \nu_N^{sa} \mathcal{C}_{M|M \setminus N}^{\infty, w} \rightarrow \nu_N^{sa} \mathcal{C}_M^{\infty, w} \rightarrow \tau^{-1} \mathcal{C}_{M|N}^{\infty, w} \rightarrow 0.$$

Now let us study the relation with the constructions of [6]. In that work the author defined the functors of Whitney specialization as follows: let $F \in D_{\mathbb{R}\text{-c}}^b(k_M)$, then

$$w\nu_N(F, \mathcal{C}_M^\infty) = s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X}}}(\mathcal{D}_{\widetilde{M}_N \rightarrow M}, (p^{-1}F)_\Omega \otimes^w \mathcal{C}_M^\infty).$$

The stalks are given by the following formula: let $v \in T_N M$. Then

$$H^k(w\nu_N(F, \mathcal{C}_M^\infty))_v \simeq \varinjlim_U H^k(X; F_U \otimes^w \mathcal{C}_M^\infty),$$

where $U \in \text{Op}(M_{sa})$ l.c.t. such that $v \notin C_M(X \setminus U)$. By Theorem 4.2.2 (ii) it turns out that

$$w\nu_N(F, \mathcal{C}_M^\infty) \simeq \rho^{-1} \nu_N^{sa} \mathcal{C}_{M|F}^{\infty, w},$$

this means that Whitney specialization is obtained by specializing the sheaf $\mathcal{C}_{M|F}^{\infty, w}$.

Assume that $M \simeq \{(x, y) \in \mathbb{R}^\ell \times \mathbb{R}^{n-\ell}\}$ and $N \simeq \{0\} \times \mathbb{R}^{n-\ell}$. A sector S of M is a subanalytic open subset $S = U \times V$ with $U \in \text{Op}(\mathbb{R}_{sa}^{n-\ell})$ and $V = W \cap B(0, \varepsilon)$, where $W \in \text{Op}(\mathbb{R}_{sa, \mathbb{R}^+}^\ell)$ and $B(0, \varepsilon)$ is the open ball of center 0 and radius ε . We say that S' is a subsector of S if $\overline{S'} \setminus N \subset S$ or, with the notations of Definition 2.1.7 if $\mathbb{R}^+ S' \subset \subset \mathbb{R}^+ S$ with the conic topology. We write for short $S' < S$.

Definition 6.3.4 Let S be an open sector of M and let $f \in \mathcal{C}_M^\infty$. One says that f is asymptotically developable along M , if there exists a formal series $\sum_{k \in \mathbb{N}^\ell} a_k(x) y^k$ with $\mathcal{C}^\infty$ coefficients a_k such that, for all $S' < S$, $m \in \mathbb{N}$, there exists $C > 0$ such that

$$\forall (x, y) \in S', \quad \left| f(x, y) - \sum_{|k| \leq m} a_k(x) y^k \right| \leq C \|y\|^m.$$

One denotes by $\sigma_N(S) \subset \mathcal{C}_M^\infty$ the space of functions asymptotically developable along M , and by $\sigma_M^\infty = \{f \in \mathcal{C}_N^\infty(S), \forall k \in \mathbb{N}^\ell, D^k f \in \sigma_M(S)\}$.

We have the following result (see Proposition 2.10 of [6]).

Proposition 6.3.5 Let S be a sector of M . Then $\Gamma(\mathbb{R}^+ S; \rho^{-1} \nu_N^{sa} \mathcal{C}_M^{\infty, w}) \simeq \sigma_N^\infty(S)$ and $\Gamma(\mathbb{R}^+ S; \rho^{-1} \nu_N^{sa} \mathcal{C}_{M|M \setminus N}^{\infty, w})$ is the subspace of functions asymptotically developable to the identically zero series.

Applying the functor ρ^{-1} to the exact sequence (6.2) we obtain the exact sequence

$$0 \rightarrow \rho^{-1} \nu_N^{sa} \mathcal{C}_{M|M \setminus N}^{\infty, w} \rightarrow \rho^{-1} \nu_N^{sa} \mathcal{C}_M^{\infty, w} \rightarrow \rho^{-1} \tau^{-1} \mathcal{C}_{M|N}^{\infty, w} \rightarrow 0,$$

where the surjective arrow is the map which associates to a function its asymptotic expansion.

Let X be a complex manifold and let Z be a complex submanifold of X . Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We denote by $\mathcal{O}_{X|F}^w$ the sheaf defined as follows:

$$\mathcal{O}_{X|F}^w := R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mathcal{C}_{X_{\mathbb{R}}|F}^{\infty, w}).$$

Let $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$ be an exact sequence in $\text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_X)$. Then the exact sequence (6.1) gives rise to the distinguished triangle

$$(6.3) \quad \mathcal{O}_{X|F}^w \rightarrow \mathcal{O}_{X|G}^w \rightarrow \mathcal{O}_{X|H}^w \xrightarrow{+}.$$

If we consider the functor of specialization of formal cohomology of [6]

$$w\nu_Z(F, \mathcal{O}_X) = R\mathcal{H}om_{\tau^{-1} \mathcal{D}_{\overline{X}}}(\tau^{-1} \mathcal{O}_{\overline{X}}, w\nu_Z(F, \mathcal{C}_{X_{\mathbb{R}}}^\infty)),$$

we have the isomorphism

$$w\nu_Z(F, \mathcal{O}_X) \simeq \rho^{-1} \nu_Z^{sa} \mathcal{O}_{X|F}^w.$$

Setting $F = \mathbb{C}_{X \setminus Z}$, $G = \mathbb{C}_X$, $H = \mathbb{C}_Z$ in (6.3) and applying the functor of specialization, we have the distinguished triangle

$$(6.4) \quad \rho^{-1} \nu_Z^{sa} \mathcal{O}_{X|X \setminus Z}^w \rightarrow \rho^{-1} \nu_Z^{sa} \mathcal{O}_X^w \rightarrow \rho^{-1} \tau^{-1} \mathcal{O}_{X|Z} \xrightarrow{+}.$$

The sheaves $\rho^{-1} \nu_Z^{sa} \mathcal{O}_X^w$ and $\rho^{-1} \tau^{-1} \mathcal{O}_{X|Z}$ are concentrated in degree zero. This follows from the following result of [7]: if $U \in \text{Op}(X_{sa})$ is convex, then $\text{R}\Gamma(X; \mathbb{C}_{\overline{U}}^w \otimes \mathcal{O}_X)$ is concentrated in degree zero. Moreover (see [13]) the sheaf $\rho^{-1} \mathcal{O}_{X|Z}^w$ is isomorphic to the sheaf $\widehat{\mathcal{O}_X|_Z}$, the formal completion of $\mathcal{O}_X$ along Z . We have an exact sequence

$$0 \rightarrow \rho^{-1} H^0 \nu_Z^{sa} \mathcal{O}_{X|X \setminus Z}^w \rightarrow \rho^{-1} \nu_Z^{sa} \mathcal{O}_X \rightarrow \tau^{-1} \widehat{\mathcal{O}_X|_Z} \rightarrow \rho^{-1} H^1 \nu_Z^{sa} \mathcal{O}_{X|X \setminus Z}^w \rightarrow 0.$$

Let $\sigma_Z^{hol}(S)$ be the space of holomorphic function asymptotically developable in S , having an asymptotic expansion with holomorphic coefficients. We have the following results of [6].

Proposition 6.3.6 *Let S be a sector of M . Then $\Gamma(\mathbb{R}^+ S; \rho^{-1} \nu_Z^{sa} \mathcal{O}_X^w) \simeq \sigma_Z^{hol}(S)$ and $\Gamma(\mathbb{R}^+ S; \rho^{-1} \nu_Z^{sa} \mathcal{O}_{X|X \setminus Z}^w)$ is the subspace of functions asymptotically developable to the identically zero series.*

Proposition 6.3.7 *The distinguished triangle (6.4) induces an exact sequence outside the zero section*

$$(6.5) \quad 0 \rightarrow \rho^{-1} H^0 \nu_Z^{sa} \mathcal{O}_{X|Z}^w|_{\dot{T}_Z X} \rightarrow \rho^{-1} \nu_Z^{sa} \mathcal{O}_X^w|_{\dot{T}_Z X} \rightarrow \dot{\tau}^{-1} \widehat{\mathcal{O}_X|_Z} \rightarrow 0.$$

On the zero section we have the exact sequence

$$(6.6) \quad 0 \rightarrow \mathcal{O}_X|_Z \rightarrow \widehat{\mathcal{O}_X|_Z} \rightarrow \rho^{-1} H^1 \nu_Z^{sa} \mathcal{O}_{X|X \setminus Z}^w|_Z \rightarrow 0.$$

Remark that on the exact sequence (6.6) we used Theorem 4.2.2 (iii) and the fact that $\rho^{-1} \mathcal{O}_X^w \simeq \mathcal{O}_X$.

Example 6.3.8 *Set $X = \mathbb{C}$ and $Z = \{0\}$. Outside the zero section the sheaves $\rho^{-1} \nu_Z^{sa} \mathcal{O}_{X|X \setminus Z}^w$ and $\rho^{-1} \nu_Z^{sa} \mathcal{O}_X^w$ are the well known sheaves $\mathcal{A}_0$ and $\mathcal{A}$ of Malgrange [20] and Sibuya [29]. These sheaves were defined in the real blow up of the origin of $\mathbb{C}$ identified with $\mathbb{S}^1 \times (\mathbb{R}^+ \cup \{0\})$. Let π be the projection on $\mathbb{C}$. The sequence (6.5) is a generalization of the exact sequence in $\text{Mod}(\mathbb{C}_{\mathbb{S}^1})$*

$$0 \rightarrow \mathcal{A}_0 \rightarrow \mathcal{A} \rightarrow \pi^{-1} \mathbb{C}[[z]] \rightarrow 0,$$

and the sequence (6.6) is a generalization of the exact sequence

$$0 \rightarrow \mathbb{C}\{z\} \rightarrow \mathbb{C}[[z]] \rightarrow H^1(\mathbb{S}^1; \mathcal{A}_0) \rightarrow 0.$$

6.4 Microlocalization of $\mathcal{O}_X^t$ and $\mathcal{O}_X^w$

Let $f : X \rightarrow Y$ be a smooth morphism of real analytic manifolds. We have the following results (see the Appendix):

$$\begin{aligned} R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{D}b_X^t) &\simeq f^{-1} \mathcal{D}b_Y^t, \\ R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{C}_X^{\infty, w}) &\simeq f^{-1} \mathcal{C}_Y^{\infty, w}. \end{aligned}$$

Let us consider the normal deformation of the diagonal in $X \times X$ of diagram (5.1). Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We recall the definitions of the Andronikof's functor of microlocalization of tempered distributions

$$t\mu hom(F, \mathcal{D}b_X) := (s^{-1}(\mathcal{D}_{\widetilde{X \times X} \xrightarrow{p_1} X} \otimes_{\mathcal{D}_{\widetilde{X \times X}}} T\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{D}b_{\widetilde{X \times X}})[-1]))^\wedge$$

and Colin's microlocalization of the Whitney tensor product

$$F \otimes_\mu^w \mathcal{C}_X^\infty := (s^{-1} R\mathcal{H}om(\mathcal{D}_{\widetilde{X \times X} \xrightarrow{p_1} X}, (p_2^{-1}F)_{\overline{\Omega}} \otimes_\mu^w \mathcal{C}_{\widetilde{X \times X}}^\infty))^\vee.$$

Theorem 6.4.1 *Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We have the isomorphisms*

$$(6.7) \quad \rho^{-1} \mu hom^{sa}(F, \mathcal{D}b_X^t) \simeq t\mu hom(F, \mathcal{D}b_X),$$

$$(6.8) \quad \rho^{-1} \mu hom^{sa}(F, \mathcal{C}_X^{\infty, w}) \simeq (D'F \otimes_\mu^w \mathcal{C}_X^\infty)^a,$$

where $(\cdot)^a$ denotes the direct image of the antipodal map.

Proof. Let $G \in D(\mathbb{C}_{X_{sa}})$. We have

$$\begin{aligned} \mu hom^{sa}(F, G) &\simeq (s^{-1} R\Gamma_\Omega p^{-1} R\mathcal{H}om(q_2^{-1}F, q_1^!G))^\wedge \\ &\simeq (s^{-1} R\Gamma_\Omega R\mathcal{H}om(p^{-1}q_2^{-1}F, p^{-1}q_1^!G))^\wedge \\ &\simeq (s^{-1} R\mathcal{H}om((p^{-1}q_2^{-1}F)_\Omega, p^{-1}q_1^!G))^\wedge \\ &\simeq (s^{-1} (R\mathcal{H}om((p^{-1}q_2^{-1}F)_\Omega, p^{-1}q_1^{-1}G) \otimes p^{-1}q_1^!\mathbb{C}_X))^\wedge \\ &\simeq (s^{-1} R\mathcal{H}om((p_2^{-1}F)_\Omega, p_1^{-1}G) \otimes s^{-1} p^{-1}q_1^!\mathbb{C}_X)^\wedge, \end{aligned}$$

where the second isomorphism follows from Lemma 5.4.1.

(i) Let us prove (6.7). Setting $G = \mathcal{D}b_X^t$ and composing with ρ^{-1} we have

$$\begin{aligned}
& \rho^{-1} \mu hom^{sa}(F, \mathcal{D}b_X^t) \\
& \simeq (s^{-1} \rho^{-1} R\mathcal{H}om((p_2^{-1}F)_\Omega, p_1^{-1} \mathcal{D}b_X^t) \otimes s^{-1} p^{-1} q_1^! \mathbb{C}_X)^\wedge \\
& \simeq (s^{-1} \rho^{-1} R\mathcal{H}om_{\rho_! \mathcal{D}_{\widetilde{X \times X}}}(\rho_! \mathcal{D}_{\widetilde{X \times X} \rightarrow X}, R\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{D}b_{\widetilde{X \times X}}^t)) \otimes s^{-1} p^{-1} q_1^! \mathbb{C}_X)^\wedge \\
& \simeq (s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, \rho^{-1} R\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{D}b_{\widetilde{X \times X}}^t)) \otimes s^{-1} p^{-1} q_1^! \mathbb{C}_X)^\wedge \\
& \simeq (s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, T\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{D}b_{\widetilde{X \times X}})) \otimes s^{-1} p^{-1} q_1^! \mathbb{C}_X)^\wedge \\
& \simeq (s^{-1} (\mathcal{D}_{\widetilde{X \times X} \rightarrow X} \otimes_{\mathcal{D}_{\widetilde{X \times X}}} T\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{D}b_{\widetilde{X \times X}})[-1]))^\wedge \\
& \simeq t\mu hom(F, \mathcal{D}b_X).
\end{aligned}$$

(ii) Let us prove (6.8). Setting $G = \mathcal{C}_X^{\infty, w}$ and composing with ρ^{-1} we have

$$\begin{aligned}
& \rho^{-1} \mu hom^{sa}(F, \mathcal{C}_X^{\infty, w}) \\
& \simeq (s^{-1} \rho^{-1} R\mathcal{H}om((p_2^{-1}F)_\Omega, p_1^{-1} \mathcal{C}_X^{\infty, w}) \otimes s^{-1} p^{-1} q_1^! \mathbb{C}_X)^\wedge \\
& \simeq (s^{-1} \rho^{-1} R\mathcal{H}om((p_2^{-1}F)_\Omega, p_1^{-1} \mathcal{C}_X^{\infty, w}))^{\vee a} \\
& \simeq (s^{-1} \rho^{-1} R\mathcal{H}om_{\rho_! \mathcal{D}_{\widetilde{X \times X}}}(\rho_! \mathcal{D}_{\widetilde{X \times X} \rightarrow X}, R\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{C}_{\widetilde{X \times X}}^{\infty, w})))^{\vee a} \\
& \simeq (s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, \rho^{-1} R\mathcal{H}om((p_2^{-1}F)_\Omega, \mathcal{C}_{\widetilde{X \times X}}^{\infty, w})))^{\vee a} \\
& \simeq (s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, D'(p_2^{-1}F)_\Omega \overset{w}{\otimes} \mathcal{C}_{\widetilde{X \times X}}^\infty))^{\vee a} \\
& \simeq (s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, (p_2^{-1} D'F)_{\overline{\Omega}} \overset{w}{\otimes} \mathcal{C}_{\widetilde{X \times X}}^\infty))^{\vee a}
\end{aligned}$$

where the last isomorphism follows since

$$D'((p_2^{-1}F)_\Omega) \simeq R\Gamma_\Omega D'(p_2^{-1}F) \simeq R\Gamma_\Omega p_2^{-1} D'F \simeq (p_2^{-1} D'F)_{\overline{\Omega}}.$$

Here we used Lemma 5.4.2 and the fact that p_2 is smooth. We have

$$s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, (p_2^{-1} D'F)_{\overline{\Omega}} \overset{w}{\otimes} \mathcal{C}_{\widetilde{X \times X}}^\infty)^{\vee a} = (D'F \overset{w}{\otimes}_\mu \mathcal{C}_X^\infty)^a$$

and the result follows. $\square$

Let X be a complex manifold and let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. In [1] and [5] the authors constructed the functors $t\mu hom(F, \mathcal{O}_X)$ of tempered microlocalization and $F \overset{w}{\otimes}_\mu \mathcal{O}_X$ of formal microlocalization taking the Dolbeaut resolutions of the real ones.

Theorem 6.4.2 *Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We have the isomorphisms*

$$(6.9) \quad \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^t) \simeq t \mu hom(F, \mathcal{O}_X),$$

$$(6.10) \quad \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^w) \simeq (D' F \underset{\mu}{\overset{w}{\otimes}} \mathcal{O}_X)^a,$$

where $(\cdot)^a$ denotes the direct image of the antipodal map.

Proof. The result follows by taking Dolbeaut resolutions on the left and the right-hand side of (6.7) and (6.8). Let us see the proof of (6.9). Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We have the chain of isomorphisms

$$\begin{aligned} \mu hom^{sa}(F, \mathcal{O}_X^t) &\simeq (s^{-1} R\mathcal{H}om((p_2^{-1} F)_\Omega, p_1^! R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mathcal{D}b_{X_{\mathbb{R}}}^t)))^\wedge[-1] \\ &\simeq (s^{-1} R\mathcal{H}om((p_2^{-1} F)_\Omega, R\mathcal{H}om_{p_1^{-1} \rho_! \mathcal{D}_{\overline{X}}}(p_1^{-1} \rho_! \mathcal{O}_{\overline{X}}, p_1^! \mathcal{D}b_{X_{\mathbb{R}}}^t)))^\wedge[-1] \\ &\simeq (s^{-1} R\mathcal{H}om_{p_1^{-1} \rho_! \mathcal{D}_{\overline{X}}}(p_1^{-1} \rho_! \mathcal{O}_{\overline{X}}, R\mathcal{H}om((p_2^{-1} F)_\Omega, p_1^! \mathcal{D}b_{X_{\mathbb{R}}}^t)))^\wedge[-1] \\ &\simeq (R\mathcal{H}om_{s^{-1} p_1^{-1} \rho_! \mathcal{D}_{\overline{X}}}(s^{-1} p_1^{-1} \rho_! \mathcal{O}_{\overline{X}}, s^{-1} R\mathcal{H}om((p_2^{-1} F)_\Omega, p_1^! \mathcal{D}b_{X_{\mathbb{R}}}^t)))^\wedge[-1] \\ &\simeq (R\mathcal{H}om_{\tau_X^{-1} \rho_! \mathcal{D}_{\overline{X}}}(\tau_X^{-1} \rho_! \mathcal{O}_{\overline{X}}, s^{-1} R\mathcal{H}om((p_2^{-1} F)_\Omega, p_1^! \mathcal{D}b_{X_{\mathbb{R}}}^t)))^\wedge[-1] \\ &\simeq R\mathcal{H}om_{\pi^{-1} \rho_! \mathcal{D}_{\overline{X}}}(\pi^{-1} \rho_! \mathcal{O}_{\overline{X}}, (s^{-1} R\mathcal{H}om((p_2^{-1} F)_\Omega, p_1^! \mathcal{D}b_{X_{\mathbb{R}}}^t)))^\wedge[-1] \\ &\simeq R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mu hom^{sa}(F, \mathcal{D}b_{X_{\mathbb{R}}}^t)). \end{aligned}$$

Applying the functor ρ^{-1} we have

$$\begin{aligned} \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^t) &\simeq \rho^{-1} R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mu hom^{sa}(F, \mathcal{D}b_{X_{\mathbb{R}}}^t)) \\ &\simeq R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \rho^{-1} \mu hom^{sa}(F, \mathcal{D}b_{X_{\mathbb{R}}}^t)) \\ &\simeq R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, t \mu hom(F, \mathcal{D}b_{X_{\mathbb{R}}})) \\ &\simeq t \mu hom(F, \mathcal{O}_X). \end{aligned}$$

The proof of (6.10) is similar. □

7 Integral transforms

In this section we give some applications related to the microlocalization of subanalytic sheaves. We show the existence of a natural action of microlocal operators on tempered and formal microlocalization. We show also the invariance under contact transformation of tempered and formal microlocalization.

7.1 $\mathcal{E}_X$ -modules

Let X be a complex manifold of complex dimension d_X . Following the notations of [11] one sets $\mathcal{E}_X^{\mathbb{R}} = H^{d_X}(\mu_{\Delta}\mathcal{O}_{X \times X}^{(0,d_X)})$. It is a sheaf of rings over T^*X and for each $F \in D^b(\mathbb{C}_X)$, $j \in \mathbb{Z}$ the sheaf $H^j \mu hom(F, \mathcal{O}_X)$ is naturally endowed with a structure of left $\mathcal{E}_X^{\mathbb{R}}$ -module. The sheaf $\mathcal{E}_X^{\mathbb{R}}$ is called the ring of microlocal operators on X . It contains a subring, denoted by $\mathcal{E}_X$ and called the ring of (finite-order) microdifferential operators. We will not recall all the properties of this sheaf and refer to [26] for a detailed study.

In [1] the author introduced the sheaf $\mathcal{E}_X^{\mathbb{R},f}$ of tempered microdifferential operators as

$$\mathcal{E}_X^{\mathbb{R},f} := H^{d_X}(t\mu hom(\mathbb{C}_{\Delta}, \mathcal{O}_{X \times X}) \bigotimes_{\mathcal{O}_{X \times X}}^L \mathcal{O}_{X \times X}^{(0,d_X)}).$$

Proposition 7.1.1 *One has $\mathcal{E}_X^{\mathbb{R},f} \simeq \rho^{-1} H^{d_X} \mu_{\Delta}^{sa} \mathcal{O}_{X \times X}^{t(0,d_X)}$.*

Proof. There is a natural morphism

$$\mu hom^{sa}(\mathbb{C}_{\Delta}, \mathcal{O}_{X \times X}) \bigotimes_{\rho! \mathcal{O}_{X \times X}}^L \rho! \mathcal{O}_{X \times X}^{(0,d_X)} \rightarrow \mu hom^{sa}(\mathbb{C}_{\Delta}, \mathcal{O}_{X \times X}^{t(0,d_X)}).$$

Applying the exact functor ρ^{-1} we obtain the morphism

$$\mathcal{E}_X^{\mathbb{R},f} \rightarrow \rho^{-1} H^{d_X} \mu hom^{sa}(\mathbb{C}_{\Delta}, \mathcal{O}_{X \times X}^{t(0,d_X)}) \simeq \rho^{-1} H^{d_X} \mu_{\Delta}^{sa} \mathcal{O}_{X \times X}^{t(0,d_X)},$$

which is an isomorphism on the fibers by Theorem 5.1.2. □

Let us recall the following results:

- the sheaf $t\mu hom(\mathbb{C}_{\Delta}, \mathcal{O}_{X \times X})$ is concentrated in degree d_X ,
- one has the ring inclusions $\mathcal{E}_X \subset \mathcal{E}_X^{\mathbb{R},f} \subset \mathcal{E}_X^{\mathbb{R}}$.

7.2 Integral transforms

Let X, Y, Z be three manifolds. Let q_{ij} be the (i, j) -th projection defined on $X \times Y \times Z$ and let p_{ij} be the (i, j) -th projection defined on $T^*X \times T^*Y \times T^*Z$. Let p_{ij}^a be the composition of p_{ij} with the antipodal map a and let $j :$

$X \times Y \times Y \times Z \rightarrow X \times Y \times Z$ be the diagonal embedding. Consider the following diagram

$$(7.1) \quad \begin{array}{ccccc} T^*(X \times Y) \times T^*(Y \times Z) & \xleftarrow{p_{12}^a \times p_{23}^a} & T^*X \times T^*Y \times T^*Z & \xrightarrow{\quad} & \\ \delta_\pi \uparrow & & \downarrow \text{id} \times p_2 \times a & & \\ T^*(X \times Y) \times_Y T^*(Y \times Z) & \xleftarrow{\quad} & T^*X \times T^*_{\Delta_Y}(Y \times Y) \times T^*Z & & \\ \downarrow \iota_{\delta'} & \square & \downarrow & & \\ T^*(X \times Y \times Z) & \xleftarrow{t_{q'_{13}}} & T^*X \times Y \times T^*Z & & \\ & & \downarrow q_{13\pi} & & \\ & & T^*X \times T^*Z & \xleftarrow{\quad} & \end{array} \quad \begin{array}{l} p_{13} \\ \end{array}$$

For $F_1 \in D^b(k_{(X \times Y)_{sa}})$ and $F_2 \in D^b(k_{(Y \times Z)_{sa}})$ set $F_1 \circ F_2 = Rq_{13!!}(q_{12}^{-1}F_1 \otimes q_{23}^{-1}F_2)$ and for $G_1 \in D^b(k_{(T^*X \times T^*Y)_{sa}})$ and $G_2 \in D^b(k_{(T^*Y \times T^*Z)_{sa}})$ set $G_1 \circ G_2 = Rp_{13\mathbb{R}+!!}^a(p_{12}^{a-1}G_1 \otimes p_{23}^{a-1}G_2)$. We need this proposition which follows from the functorial properties of μhom^{sa} (it is an adaptation of Proposition 4.4.11 of [11]).

Proposition 7.2.1 *Let us consider the sheaves $K_1 \in D_{\mathbb{R}\text{-c}}^b(k_{X \times Y})$, $F_1 \in D^b(k_{(X \times Y)_{sa}})$, $K_2 \in D_{\mathbb{R}\text{-c}}^b(k_{Y \times Z})$ and $F_2 \in D^b(k_{(Y \times Z)_{sa}})$. Suppose that q_{13} is proper on $\text{supp}(q_{12}^{-1}K_1 \otimes q_{23}^{-1}K_2)$. There is a morphism*

$$(7.2) \quad \mu hom^{sa}(K_1, F_1) \overset{a}{\circ} \mu hom^{sa}(K_2, F_2) \rightarrow \mu hom^{sa}(K_1 \circ K_2, F_1 \circ F_2).$$

Proposition 7.2.2 *Let $\lambda = \emptyset, t$. Let $K_1 \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_{X \times Y})$ and $K_2 \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_{Y \times Z})$. Suppose that q_{13} is proper on $\text{supp}(q_{12}^{-1}K_1 \otimes q_{23}^{-1}K_2)$. Morphism (7.2) defines a morphism*

$$(7.3) \quad \begin{aligned} \mu hom^{sa}(K_1, \mathcal{O}_{X \times Y}^{\lambda(0, d_Y)}) \overset{a}{\circ} \mu hom^{sa}(K_2, \mathcal{O}_{Y \times Z}^{\lambda(0, d_Z)}) \\ \rightarrow \mu hom^{sa}(K_1 \circ K_2, \mathcal{O}_{X \times Z}^{\lambda(0, d_Z)})[-d_Y]. \end{aligned}$$

Proof. It follows from (7.2) setting $F_1 = \mathcal{O}_{X \times Y}^{\lambda(0, d_Y)}$, $F_2 = \mathcal{O}_{Y \times Z}^{\lambda(0, d_Z)}$ and using the integration morphism $\mathcal{O}_{X \times Y}^{\lambda(0, d_Y)} \circ \mathcal{O}_{Y \times Z}^{\lambda(0, d_Z)} \rightarrow \mathcal{O}_{X \times Z}^{\lambda(0, d_Z)}[-d_Y]$. $\square$

Corollary 7.2.3 *Morphism (7.2) induces the ring structures on $\mathcal{E}_X^{\mathbb{R}}$ and $\mathcal{E}_X^{\mathbb{R}, f}$.*

Proof. It follows applying ρ^{-1} to (7.3) with $X = Y = Z$ and $K_1 = K_2 = \mathbb{C}_\Delta[-d_X]$. □

Proposition 7.2.4 *Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. Morphism (7.2) defines a morphism*

$$(7.4) \quad \mu_\Delta^{sa} \mathcal{O}_{X \times X}^{\lambda(0, d_X)}[d_X] \otimes \mu hom^{sa}(F, \mathcal{O}_X^\lambda) \rightarrow \mu hom^{sa}(F, \mathcal{O}_X^\lambda).$$

Proof. We apply Proposition 7.2.1 with $X = Y$, $Z = \{\text{point}\}$. We set $(K_1, K_2, F_1, F_2) = (\mathbb{C}_\Delta[-d_X], F, \mathcal{O}_{X \times X}^{\lambda(0, d_X)}, \mathcal{O}_X^\lambda)$. In this case we have $\mathbb{C}_\Delta \circ F \simeq F$. We obtain the desired morphism using the integration morphism $\mathcal{O}_{X \times X}^{\lambda(0, d_X)} \circ \mathcal{O}_X^\lambda \rightarrow \mathcal{O}_X^\lambda[-d_X]$. □

Applying the functor ρ^{-1} to (7.4), we find the morphisms of [1] and [11] (recall that $\rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^t) \simeq t \mu hom(F, \mathcal{O}_X)$).

Corollary 7.2.5 *Morphism (7.4) induces morphisms*

$$(7.5) \quad \mathcal{E}_X^{\mathbb{R}, f} \otimes \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^t) \rightarrow \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^t)$$

$$(7.6) \quad \mathcal{E}_X^{\mathbb{R}} \otimes \mu hom(F, \mathcal{O}_X) \rightarrow \mu hom(F, \mathcal{O}_X)$$

which induces a structure of $\mathcal{E}_X^{\mathbb{R}, f}$ -module (resp. $\mathcal{E}_X^{\mathbb{R}}$ -module) on the sheaves $H^k \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^t)$ (resp. $H^k \mu hom(F, \mathcal{O}_X)$), for each $k \in \mathbb{Z}$.

Now we will study the action of $\mathcal{E}_X^{\mathbb{R}, f}$ on formal microlocalization. We first need to introduce the sheaf of tempered $\mathcal{C}^\infty$ -functions.

Definition 7.2.6 *Let X be a real analytic manifold and let $U \in \text{Op}(X)$. Let $f \in \Gamma(U; \mathcal{C}_X^\infty)$. One says that f has polynomial growth at $p \in X$ if for a local coordinate system $(x_1, \dots, x_n)$ around p , there exists a compact neighborhood K of p and $N \in \mathbb{N}$ such that*

$$\sup_{x \in K \cap U} (d(x, K \setminus U))^N |f(x)| < \infty.$$

One says that f is tempered at p if all its derivatives have polynomial growth at p . One says that f is tempered if it is tempered at any point.

Definition 7.2.7 One denotes by $\mathcal{C}_X^{\infty,t}$ the presheaf of tempered $\mathcal{C}^\infty$ -functions on X_{sa} defined as follows:

$$U \mapsto \{f \in \Gamma(U; \mathcal{C}_X^\infty), f \text{ is tempered}\}.$$

As a consequence of a result of [13], for $U, V \in \text{Op}(X_{sa})$ the sequence

$$0 \rightarrow \mathcal{C}_X^{\infty,t}(U \cup V) \rightarrow \mathcal{C}_X^{\infty,t}(U) \oplus \mathcal{C}_X^{\infty,t}(V) \rightarrow \mathcal{C}_X^{\infty,t}(U \cap V)$$

is exact. Then $\mathcal{C}_X^{\infty,t}$ is a sheaf on X_{sa} . Moreover $R\Gamma(U; \mathcal{C}_X^{\infty,t})$ is concentrated in degree zero for any $U \in \text{Op}(X_{sa})$.

We have the following results (see [14]).

Proposition 7.2.8 For each $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ one has the isomorphism

$$\rho^{-1}R\mathcal{H}om(F, \mathcal{C}_X^{\infty,t}) \simeq T\mathcal{H}om(F, \mathcal{C}_X^\infty),$$

where $T\mathcal{H}om(F, \mathcal{C}_X^\infty)$ is the sheaf of [13]. When $F = \mathbb{C}_U$, $U \in \text{Op}(X_{sa})$ it is defined by $V \mapsto \mathcal{C}_V^{\infty,t}(U \cap V)$.

Proposition 7.2.9 Let X be a complex manifold, $X_{\mathbb{R}}$ the underlying real analytic manifold and $\overline{X}$ the conjugate manifold. Then

$$\mathcal{O}_X^t \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mathcal{C}_{X_{\mathbb{R}}}^{\infty,t}).$$

We prove the following result.

Lemma 7.2.10 Let $f : X \rightarrow Y$ be a smooth morphism of real analytic manifolds. Then we have the isomorphism

$$f^{-1}\mathcal{C}_Y^{\infty,t} \xrightarrow{\sim} R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{C}_X^{\infty,t}).$$

Proof. We may reduce to the case of a projection $\pi : Y \times \mathbb{R} \rightarrow Y$. We shall prove that the morphism

$$\partial_t : \mathcal{C}_{Y \times \mathbb{R}}^{\infty,t} \rightarrow \mathcal{C}_{Y \times \mathbb{R}}^{\infty,t},$$

where t denotes the variable in $\mathbb{R}$, is surjective. Let $U \in \text{Op}^c((Y \times \mathbb{R})_{sa})$, then by Lemma A.1.11 it admits a finite covering $\{U_i\}_{i=1}^N$ such that each U_i is simply connected and the intersections of each U_i with the fibers of π are contractible (or empty). Hence we may reduce to the case that the intersections of U with the fibers of π are contractible (or empty). Moreover we can assume that

$$U = \{(x, t) \in Y \times \mathbb{R}; f(x) < t < g(x)\},$$

where $f, g : \pi(U) \rightarrow \mathbb{R}$ are continuous subanalytic maps and $\pi(U)$ is simply connected. Let us consider $h, k : \pi(U) \rightarrow \mathbb{R}$ subanalytic and $\varphi \in \Gamma(\pi(U); \mathcal{C}_Y^\infty)$ such that $f < h < \varphi < k < g$.

Let $s \in \Gamma(U; \mathcal{C}_{Y \times \mathbb{R}}^{\infty, t})$ and define

$$\tilde{s}(x, t) = \int_{(x, \varphi(x))}^{(x, t)} s(x, \tau) d\tau.$$

Then $\tilde{s} \in \Gamma(U; \mathcal{C}_{Y \times \mathbb{R}}^\infty)$ and $\partial_t \tilde{s} = s$. Moreover

$$|\tilde{s}(x, t)| \leq |\varphi(x) - t| \sup_{(x, \tau) \in \{x\} \times [\varphi(x), t]} |s(x, \tau)|.$$

Since U is bounded, there exists $M > 0$ such that $|\varphi(x) - t| < M$ for each $(x, t) \in U$. Since s is tempered, for each $x \in \pi(U)$ and each $\tau \in [\varphi(x), t]$ there exist $c_1, r_1 > 0$ such that

$$\begin{aligned} |s(x, \tau)| &\leq c_1 \frac{1}{d((x, \tau), \partial U)^{r_1}} \\ &\leq c_1 \frac{1}{\min(d((x, t), \partial U), d((x, h(x)), \partial U), d((x, k(x)), \partial U))^{r_1}}. \end{aligned}$$

As a consequence of Łojaciewicz's inequality (see Theorem 6.4 of [3]) there exist $c_2, r_2 > 0$ such that

$$d((x, h(x)), \partial U), d((x, k(x)), \partial U) \geq c_2 d(x, \partial(\pi(U)))^{r_2} \geq c_2 d((x, t), \partial U)^{r_2}.$$

Hence there exists $c, r > 0$ such that

$$\tilde{s}(x, t) \leq c \frac{1}{d((x, t), \partial U)^r}$$

and the result follows. □

Lemma 7.2.11 *Let $f : X \rightarrow Y$ be a smooth morphism of real analytic manifolds. Let $\mathcal{M}, \mathcal{N} \in \mathcal{D}^b(\mathcal{D}_X)$. There is a natural morphism*

$$R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{M}) \underset{f^{-1}\mathcal{A}_Y}{\overset{L}{\otimes}} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{N}) \rightarrow R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{M} \overset{L}{\otimes}_{\mathcal{A}_X} \mathcal{N}).$$

Proof. By Lemma 4.9 of [10] we have

$$(7.7) \quad \mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\mathcal{A}_X} \mathcal{D}_{X \rightarrow Y} \simeq \mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{f^{-1}\mathcal{A}_Y} f^{-1}\mathcal{D}_Y.$$

Then if $\mathcal{M}$ is a $\mathcal{D}_X$ -module

$$\begin{aligned} (\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\mathcal{D}_X} \mathcal{M}) \overset{L}{\otimes}_{f^{-1}\mathcal{A}_Y} f^{-1}\mathcal{D}_Y &\simeq (\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{f^{-1}\mathcal{A}_Y} f^{-1}\mathcal{D}_Y) \overset{L}{\otimes}_{\mathcal{D}_X} \mathcal{M} \\ &\simeq (\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\mathcal{A}_X} \mathcal{D}_{X \rightarrow Y}) \overset{L}{\otimes}_{\mathcal{D}_X} \mathcal{M} \\ &\simeq \mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\mathcal{D}_X} (\mathcal{M} \overset{L}{\otimes}_{\mathcal{A}_X} \mathcal{D}_{X \rightarrow Y}). \end{aligned}$$

Now when f is smooth $\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\mathcal{D}_X} \cdot \simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \cdot)[d_X - d_Y]$. Then if $\mathcal{N}$ is another $\mathcal{D}_X$ -module

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{M}) \overset{L}{\otimes}_{f^{-1}\mathcal{A}_Y} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{N}) \\ \simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{M} \overset{L}{\otimes}_{\mathcal{A}_X} \mathcal{D}_{X \rightarrow Y} \overset{L}{\otimes}_{f^{-1}\mathcal{D}_Y} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{N})) \\ \rightarrow R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow Y}, \mathcal{M} \overset{L}{\otimes}_{\mathcal{A}_X} \mathcal{N}). \end{aligned}$$

□

Lemma 7.2.12 *Let X be a real analytic manifold. Let $F, G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ and let S be a closed subanalytic subset of X . There is a morphism*

$$\begin{aligned} \rho^{-1}R\mathcal{H}om(F, (\mathcal{C}_X^{\infty, t})_S) \otimes_{\mathcal{A}_X} \rho^{-1}R\mathcal{H}om(D'((F \otimes G)_S), \mathcal{C}_X^{\infty, w}) \\ \rightarrow \rho^{-1}R\mathcal{H}om(D'(G_S), \mathcal{C}_X^{\infty, w}). \end{aligned}$$

Proof. (i) Let $V_1, V_2 \in \text{Op}(X_{sa})$. The sheaf $\rho^{-1}\Gamma_{V_1}(\mathcal{C}_X^{\infty, t})_S$ is concentrated in degree zero since $\mathcal{C}_X^{\infty, t}$ is $\Gamma(U; \cdot)$ -acyclic for each $U \in \text{Op}(X_{sa})$. Moreover the sheaves $\rho^{-1}R\mathcal{H}om(D'\mathbb{C}_{V_1 \cap V_2 \cap S}, \mathcal{C}_X^{\infty, w}) \simeq \mathbb{C}_{V_1 \cap V_2 \cap S}^w \otimes \mathcal{C}_X^{\infty}$ and $\rho^{-1}R\mathcal{H}om(D'\mathbb{C}_{V_2 \cap S}, \mathcal{C}_X^{\infty, w}) \simeq \mathbb{C}_{V_2 \cap S}^w \otimes \mathcal{C}_X^{\infty}$ are also concentrated in degree zero. There is a morphism

$$(7.8) \quad \rho^{-1}\Gamma_{V_1}(\mathcal{C}_X^{\infty, t})_S \otimes_{\mathcal{A}_X} \mathbb{C}_{V_1 \cap V_2 \cap S}^w \otimes \mathcal{C}_X^{\infty} \rightarrow \mathbb{C}_{V_2 \cap S}^w \otimes \mathcal{C}_X^{\infty}.$$

This follows since the multiplication of a function tempered on V_1 by a function vanishing with all its derivatives outside V_1 is a function vanishing with all its derivatives outside V_1 .

(ii) By Theorem 1.1 of [13] the morphism (7.8) extends to a morphism

$$(7.9) \quad \rho^{-1}\Gamma_{V_1}(\mathcal{C}_X^{\infty,t})_S \otimes_{\mathcal{A}_X} G_{V_1 \cap S} \overset{w}{\otimes} \mathcal{C}_X^\infty \rightarrow G_S \overset{w}{\otimes} \mathcal{C}_X^\infty,$$

functorial in $G \in \text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_X)$. By adjunction this gives a morphism

$$(7.10) \quad \rho^{-1}\Gamma_{V_1}(\mathcal{C}_X^{\infty,t})_S \rightarrow \mathcal{H}om_{\mathcal{A}_X}(G_{V_1 \cap S} \overset{w}{\otimes} \mathcal{C}_X^\infty, G_S \overset{w}{\otimes} \mathcal{C}_X^\infty).$$

By Theorem 1.1 of [13] the morphism (7.10) extends to a morphism

$$(7.11) \quad \rho^{-1}\mathcal{H}om(F, (\mathcal{C}_X^{\infty,t})_S) \rightarrow \mathcal{H}om_{\mathcal{A}_X}((F \otimes G)_S \overset{w}{\otimes} \mathcal{C}_X^\infty, G_S \overset{w}{\otimes} \mathcal{C}_X^\infty).$$

functorial in $F \in \text{Mod}_{\mathbb{R}\text{-c}}(\mathbb{C}_X)$.

(iii) Let $F, G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We have the following chain of morphisms

$$\begin{aligned} \rho^{-1}R\mathcal{H}om(F, (\mathcal{C}_X^{\infty,t})_S) &\xrightarrow{\sim} R(\rho^{-1}\mathcal{H}om(F, (\mathcal{C}_X^{\infty,t})_S)) \\ &\rightarrow R(\mathcal{H}om_{\mathcal{A}_X}((F \otimes G)_S \overset{w}{\otimes} \mathcal{C}_X^\infty, G_S \overset{w}{\otimes} \mathcal{C}_X^\infty)) \\ &\rightarrow R\mathcal{H}om_{\mathcal{A}_X}((F \otimes G)_S \overset{w}{\otimes} \mathcal{C}_X^\infty, G_S \overset{w}{\otimes} \mathcal{C}_X^\infty), \end{aligned}$$

where the first isomorphism follows since ρ^{-1} is exact and $(\cdot)_S$ sends quasi-injective objects to quasi-injective objects, the second arrow follows from (7.11) and the third one is a canonical morphism of derived functors (see [12], Proposition 13.3.13).

By adjunction we obtain the desired morphism. $\square$

Lemma 7.2.13 *Let us consider the normal deformation of the diagonal in $X \times X$ of diagram (5.1). Let $F, G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. There is a morphism*

$$\begin{aligned} \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}F, q_2^{-1}\mathcal{C}_X^{\infty,t}) \otimes_{\mathcal{A}_X} \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}D'(F \otimes G), q_2^{-1}\mathcal{C}_X^{\infty,w}) \\ \rightarrow \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}D'G, q_2^{-1}\mathcal{C}_X^{\infty,w}). \end{aligned}$$

Proof. (i) As in the proof of Theorem 6.4.1, if X is a real analytic manifold, $K \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$, $\lambda = t, w$, we have

$$\begin{aligned} \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}K, q_2^{-1}\mathcal{C}_X^{\infty,\lambda}) \\ \simeq \rho^{-1}s^{-1}R\mathcal{H}om((p_1^{-1}K)_{\Omega}, p_2^{-1}\mathcal{C}_X^{\infty,\lambda}) \\ \simeq \rho^{-1}s^{-1}R\mathcal{H}om((p_1^{-1}K)_{\Omega}, R\mathcal{H}om_{\rho_!\mathcal{D}_{\widetilde{X \times X}}}(\rho_!\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, \mathcal{C}_{\widetilde{X \times X}}^{\infty,\lambda})) \\ \simeq s^{-1}R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \rightarrow X}, \rho^{-1}R\mathcal{H}om((p_1^{-1}K)_{\Omega}, \mathcal{C}_{\widetilde{X \times X}}^{\infty,\lambda})), \end{aligned}$$

where the second isomorphism follows from Lemma 7.2.10.

(ii) By Lemma 5.4.2 for $H \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ we have $(p_1^{-1}D'H)_\Omega \simeq D'((p_1^{-1}H)_{\overline{\Omega}})$ and $R\Gamma_\Omega p_2^{-1}\mathcal{C}_X^{\infty,t} \simeq (p_2^{-1}\mathcal{C}_X^{\infty,t})_{\overline{\Omega}}$.

(iii) By Lemma 7.2.11 with $(X, Y) = (\widetilde{X \times X}, X)$, $\mathcal{M} = R\mathcal{H}om(p_1^{-1}F, (\mathcal{C}_{\widetilde{X \times X}}^{\infty,t})_{\overline{\Omega}})$, $\mathcal{N} = R\mathcal{H}om(D'((p_1^{-1}(F \otimes G))_{\overline{\Omega}}), \mathcal{C}_{\widetilde{X \times X}}^{\infty,w})$, we are reduced to find a morphism

$$\begin{aligned} \rho^{-1}R\mathcal{H}om(p_1^{-1}F, (\mathcal{C}_{\widetilde{X \times X}}^{\infty,t})_{\overline{\Omega}}) \otimes_{\mathcal{A}_{\widetilde{X \times X}}} \rho^{-1}R\mathcal{H}om(D'((p_1^{-1}(F \otimes G))_{\overline{\Omega}}), \mathcal{C}_{\widetilde{X \times X}}^{\infty,w}) \\ \rightarrow \rho^{-1}R\mathcal{H}om(D'((p_1^{-1}G)_{\overline{\Omega}}), \mathcal{C}_{\widetilde{X \times X}}^{\infty,w}) \end{aligned}$$

which follows replacing (X, S, F, G) with $(\widetilde{X \times X}, \overline{\Omega}, p_1^{-1}F, p_1^{-1}G)$ in Lemma 7.2.12. $\square$

Let us consider the complex case. Let X be a complex manifold.

Lemma 7.2.14 *Let $\mathcal{L}, \mathcal{H} \in D^b(\mathcal{D}_{X_{\mathbb{R}}})$. There is a natural morphism*

$$R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L}) \overset{L}{\otimes}_{\mathcal{O}_X} R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{H}) \rightarrow R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H}).$$

Proof. By definition we have $\mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H} = \mathcal{D}_{X_{\mathbb{R}} \rightarrow X_{\mathbb{R}} \times X_{\mathbb{R}}} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}} \times X_{\mathbb{R}}}} (\mathcal{L} \overset{\mathcal{D}}{\boxtimes} \mathcal{M})$.

Hence we get

$$\mathcal{L} \overset{\mathcal{D}}{\boxtimes} \mathcal{H} \rightarrow R\mathcal{H}om_{\mathcal{D}_{X_{\mathbb{R}}}}(\mathcal{D}_{X_{\mathbb{R}} \rightarrow X_{\mathbb{R}} \times X_{\mathbb{R}}}, \mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H}).$$

There is a chain of morphisms

$$\begin{aligned}
& R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L}) \overset{L}{\otimes}_{\mathcal{O}_X} R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{H}) \\
& \simeq \mathcal{D}_{X \rightarrow X \times X} \overset{L}{\otimes}_{\mathcal{D}_{X \times X}} (R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L}) \overset{\mathcal{D}}{\boxtimes} R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{H})) \\
& \rightarrow \mathcal{D}_{X \rightarrow X \times X} \overset{L}{\otimes}_{\mathcal{D}_{X \times X}} R\mathcal{H}om_{\mathcal{D}_{\overline{X}} \boxtimes \mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}} \boxtimes \mathcal{O}_{\overline{X}}, \mathcal{D}_{X \times X} \overset{L}{\otimes}_{\mathcal{D}_X \boxtimes \mathcal{D}_X} (\mathcal{L} \boxtimes \mathcal{H})) \\
& \rightarrow \mathcal{D}_{X \rightarrow X \times X} \overset{L}{\otimes}_{\mathcal{D}_{X \times X}} R\mathcal{H}om_{\mathcal{D}_{\overline{X} \times \overline{X}}}(\mathcal{O}_{\overline{X} \times \overline{X}}, \mathcal{D}_{\overline{X} \times \overline{X}} \overset{L}{\otimes}_{\mathcal{D}_{\overline{X}} \boxtimes \mathcal{D}_{\overline{X}}} \mathcal{D}_{X \times X} \overset{L}{\otimes}_{\mathcal{D}_X \boxtimes \mathcal{D}_X} (\mathcal{L} \boxtimes \mathcal{H})) \\
& \rightarrow \mathcal{D}_{X \rightarrow X \times X} \overset{L}{\otimes}_{\mathcal{D}_{X \times X}} R\mathcal{H}om_{\mathcal{D}_{\overline{X} \times \overline{X}}}(\mathcal{O}_{\overline{X} \times \overline{X}}, \mathcal{L} \overset{\mathcal{D}}{\boxtimes} \mathcal{H}) \\
& \rightarrow \mathcal{D}_{X \rightarrow X \times X} \overset{L}{\otimes}_{\mathcal{D}_{X \times X}} R\mathcal{H}om_{\mathcal{D}_{\overline{X} \times \overline{X}}}(\mathcal{O}_{\overline{X} \times \overline{X}}, R\mathcal{H}om_{\mathcal{D}_{X_{\mathbb{R}}}}(\mathcal{D}_{X_{\mathbb{R}} \rightarrow X_{\mathbb{R}} \times X_{\mathbb{R}}}, \mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H})) \\
& \simeq \mathcal{D}_{X \rightarrow X \times X} \overset{L}{\otimes}_{\mathcal{D}_{X \times X}} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{D}_{X \rightarrow X \times X}, R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H})) \\
& \rightarrow R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H}).
\end{aligned}$$

□

Lemma 7.2.15 *Let us consider the normal deformation of the diagonal in $X \times X$ of diagram (5.1). Let $F, G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. There is a morphism*

$$\begin{aligned}
& \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}F, q_2^{-1}\mathcal{O}_X^t) \otimes_{\mathcal{O}_X} \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(F \otimes G), q_2^{-1}\mathcal{O}_X^w) \\
& \rightarrow \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'G, q_2^{-1}\mathcal{O}_X^w).
\end{aligned}$$

Proof. If X is a complex manifold, $K \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$, $\lambda = t, w$, we have

$$\begin{aligned}
& \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}K, q_2^{-1}\mathcal{O}_X^{\lambda}) \\
& \simeq \rho^{-1}s^{-1} R\mathcal{H}om((p_1^{-1}K)_{\Omega}, p_2^{-1}\mathcal{O}_X^{\lambda}) \\
& \simeq \rho^{-1}s^{-1} R\mathcal{H}om((p_1^{-1}K)_{\Omega}, p_2^{-1} R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mathcal{C}_{X_{\mathbb{R}}}^{\infty, \lambda})) \\
& \simeq R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \rho^{-1}s^{-1} R\mathcal{H}om((p_1^{-1}K)_{\Omega}, p_2^{-1}\mathcal{C}_{X_{\mathbb{R}}}^{\infty, \lambda})) \\
& \simeq R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}K, q_2^{-1}\mathcal{C}_{X_{\mathbb{R}}}^{\infty, \lambda})).
\end{aligned}$$

Set

$$\begin{aligned}
\mathcal{L} &= \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}F, q_2^{-1}\mathcal{C}_{X_{\mathbb{R}}}^{\infty, t}), \\
\mathcal{H} &= \rho^{-1}\nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(F \otimes G), q_2^{-1}\mathcal{C}_{X_{\mathbb{R}}}^{\infty, w}).
\end{aligned}$$

By Lemma 7.2.14 there is a natural morphism

$$R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L}) \overset{L}{\otimes}_{\mathcal{O}_X} R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{H}) \rightarrow R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\mathcal{O}_{\overline{X}}, \mathcal{L} \overset{L}{\otimes}_{\mathcal{A}_{X_{\mathbb{R}}}} \mathcal{H}).$$

Then the result follows from Lemma 7.2.13. $\square$

Lemma 7.2.16 *Let $f : X \rightarrow Y$ be a smooth morphism of complex manifolds. Then there is a natural morphism*

$$Rf_{!!}\Omega_X^w[d_X] \rightarrow \Omega_Y[d_Y].$$

Proof. By Theorem A.4.4 we have the isomorphism

$$f^!\mathcal{O}_Y^w[2d_Y] \xrightarrow{\sim} R\mathcal{H}om_{\rho_!\mathcal{D}_X}(\rho_!\mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w)[2d_X].$$

We have $R\mathcal{H}om_{\rho_!\mathcal{D}_X}(\rho_!\mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w) \simeq \rho_!\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\rho_!\mathcal{D}_X} \mathcal{O}_X^w[d_Y - d_X]$. Hence we get

$$\rho_!\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\rho_!\mathcal{D}_X} \mathcal{O}_X^w \simeq f^!\mathcal{O}_Y^w[d_Y - d_X].$$

By adjunction we get

$$Rf_{!!}(\rho_!\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\rho_!\mathcal{D}_X} \mathcal{O}_X^w) \rightarrow \mathcal{O}_Y^w[d_Y - d_X].$$

From this we can deduce

$$Rf_{!!}\Omega_X^w \rightarrow Rf_{!!}(\Omega_X^w \overset{L}{\otimes}_{\rho_!\mathcal{D}_X} \rho_!\mathcal{D}_{X \rightarrow Y}) \rightarrow \Omega_Y^w[d_Y - d_X].$$

$\square$

Let us consider the diagram (7.1) with $Z = \{\text{point}\}$. Set $p_X : T^*X \times T^*Y \rightarrow T^*X$, $p_Y : T^*X \times T^*Y \rightarrow T^*Y$, $q_X : X \times Y \rightarrow X$, $q_Y : X \times Y \rightarrow Y$.

Proposition 7.2.17 *Let $G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ and $K \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_{X \times Y})$ such that q_Y is proper on $\text{supp}(q_X^{-1}G) \cap \text{supp}(K)$. Then we have a morphism*

$$(7.12) \quad \begin{aligned} & \rho^{-1}\mu\text{hom}^{sa}(K, \mathcal{O}_{X \times Y}^{t(0, d_Y)})[d_Y] \overset{a}{\circ} \rho^{-1}\mu\text{hom}^{sa}(D'(K \circ G), \mathcal{O}_Y^w) \\ & \rightarrow \rho^{-1}\mu\text{hom}^{sa}(D'G, \mathcal{O}_X^w). \end{aligned}$$

Proof. We will prove the assertion in several steps. Set

$$\begin{aligned}
H_1 &= \rho^{-1} \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}K, q_2^! \mathcal{O}_{X \times Y}^{t(0, d_Y)}) \\
&\simeq \rho^{-1} \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}K, q_2^{-1} \mathcal{O}_{X \times Y}^{t(0, d_Y)})[2d_{X \times Y}] \\
H_2 &= \rho^{-1} \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(K \circ G), q_2^! \mathcal{O}_Y^w) \\
&\simeq \rho^{-1} \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(K \circ G), q_2^{-1} \mathcal{O}_Y^w)[2d_Y].
\end{aligned}$$

Since the Fourier-Sato transform commutes with ρ^{-1} we have

$$H_1^{\wedge} \simeq \rho^{-1} \mu hom^{sa}(K, \mathcal{O}_{X \times Y}^{t(0, d_Y)}), \quad H_2^{\wedge} \simeq \rho^{-1} \mu hom^{sa}(D'(K \circ G), \mathcal{O}_Y^w).$$

(i) By the commutativity of the diagram (7.1) we have an isomorphism

$$Rp_{X!}^a((H_1^{\wedge})^a \otimes p_Y^{a-1} H_2^{\wedge}) \simeq Rq_{X\pi!}^t q_X'^{-1} R^t \delta'_! \delta_{\pi}^{-1} (H_1^{\wedge} \boxtimes H_2^{\wedge}).$$

(ii) By Proposition 3.7.15 of [11] we have an isomorphism

$$(H_1)^{\wedge} \boxtimes (H_2)^{\wedge} \xrightarrow{\sim} (H_1 \boxtimes H_2)^{\wedge},$$

(iii) Denote by $Tq_Y : T(X \times Y) \rightarrow TY$ the tangent map. By Propositions 3.7.13 and 3.7.14 of [11] we have the isomorphism

$$R^t \delta'_! \delta_{\pi}^{-1} (H_1 \boxtimes H_2)^{\wedge} \simeq (H_1 \otimes Tq_Y^{-1} H_2)^{\wedge} [-2d_Y].$$

(iv) We have the chain of morphisms

$$\begin{aligned}
&Tq_Y^{-1} \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(K \circ G), q_2^{-1} \mathcal{O}_Y^w) \\
&\simeq \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}q_Y^{-1}D'(K \circ G), q_2^{-1}q_Y^{-1} \mathcal{O}_Y^w) \\
&\simeq \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(q_Y^{-1}q_{Y*}(K \otimes q_X^{-1}G)), q_2^{-1}q_Y^{-1} \mathcal{O}_Y^w) \\
&\rightarrow \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(K \otimes q_X^{-1}G), q_2^{-1}q_Y^{-1} \mathcal{O}_Y^w) \\
&\simeq \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}D'(K \otimes q_X^{-1}G), q_2^{-1} R\mathcal{H}om_{\rho! \mathcal{D}_{X \times Y}}(\rho! \mathcal{D}_{X \times Y \rightarrow Y}, \mathcal{O}_{X \times Y}^w)),
\end{aligned}$$

where the first isomorphism follows since q_Y is smooth, the second one since $\text{supp}(q_X^{-1}G) \cap \text{supp}(K)$ is proper over Y and the last one is given by inverse image formula for Whitney holomorphic functions.

(v) We have a morphism

$$(H_1 \otimes Tq_Y^{-1} H_2)^{\wedge} [-2d_Y] \rightarrow \rho^{-1} \mu hom^{sa}(q_X^{-1}D'G, \mathcal{O}_{X \times Y}^{w(0, d_Y)}).$$

To prove the existence of this morphism we use (iv) to prove the morphism

$$H_1 \otimes Tq_Y^{-1} H_2 [-2d_Y] \rightarrow \nu_{\Delta}^{sa} R\mathcal{H}om(q_1^{-1}q_X^{-1}D'G, q_2^{-1} \mathcal{O}_{X \times Y}^{w(0, d_Y)})[2d_{X \times Y}].$$

Hence we may reduce to the case of the morphism

$$\begin{aligned} \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}K, q_2^{-1}\mathcal{O}_{X \times Y}^t) \otimes \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}D'(K \otimes q_X^{-1}G), q_2^{-1}\mathcal{O}_{X \times Y}^w) \\ \rightarrow \rho^{-1}\nu_{\Delta}^{sa}R\mathcal{H}om(q_1^{-1}D'(q_X^{-1}G), q_2^{-1}\mathcal{O}_{X \times Y}^w). \end{aligned}$$

This is a consequence of Lemma 7.2.15 with (X, F, G) replaced by $(X \times Y, K, q_X^{-1}G)$.

(vi) We have the chain of morphisms

$$\begin{aligned} Rq_{X\pi!}{}^t q_X'^{-1} \rho^{-1} \mu hom^{sa}(q_X^{-1} D' G, \mathcal{O}_{X \times Y}^{w(0, d_Y)}) \\ \rightarrow \rho^{-1} \mu hom^{sa}(Rq_{X*} q_X^{-1} D' G, Rq_{X!!} \mathcal{O}_{X \times Y}^{w(0, d_Y)}) \\ \rightarrow \rho^{-1} \mu hom^{sa}(D' G, \mathcal{O}_X^w)[-d_Y], \end{aligned}$$

where the second morphism is a consequence of the integration morphism $Rq_{X!!} \mathcal{O}_{X \times Y}^{w(0, d_Y)} \rightarrow \mathcal{O}_X^w[-d_Y]$ defined in Lemma 7.2.16 (see also Remark 3.4 of [13]) and the fact that $Rq_{X*} q_X^{-1} \simeq \text{id}$. Composing morphisms (i)-(vi) we get the desired morphism. $\square$

Corollary 7.2.18 *Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. Morphism (7.12) defines a morphism*

$$(7.13) \quad \mathcal{E}_X^{\mathbb{R}, f} \otimes \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^w) \rightarrow \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^w)$$

which induces a structure of $\mathcal{E}_X^{\mathbb{R}, f}$ -module on $H^k \rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^w)$ for each $k \in \mathbb{Z}$.

Proof. We apply Proposition 7.2.17 setting $X = Y$ and $(G, K) = (D'F, \mathbb{C}_{\Delta})$. In this case we have $D'(\mathbb{C}_{\Delta} \circ D'F) \simeq D'D'F \simeq F$. $\square$

In this way we find the morphism of [5] (recall that $\rho^{-1} \mu hom^{sa}(F, \mathcal{O}_X^w) \simeq (D'F \underset{\mu}{\overset{w}{\otimes}} \mathcal{O}_X)^a$)

$$(\mathcal{E}_X^{\mathbb{R}, f})^a \otimes F \underset{\mu}{\overset{w}{\otimes}} \mathcal{O}_X \rightarrow F \underset{\mu}{\overset{w}{\otimes}} \mathcal{O}_X.$$

Remark 7.2.19 *We would like to see the compatibility between this morphism and the one of Andronikof ([1], Proposition 3.3.10). Steps (i) to (iii) of Proposition 7.2.17 are the same. We need the compatibility between the multiplications. We will see the compatibility between*

$$\rho^{-1} R\Gamma_Z \mathcal{O}_X^t \otimes \rho^{-1} R\mathcal{H}om(F, \mathcal{O}_X^t) \rightarrow \rho^{-1} R\mathcal{H}om(F, \mathcal{O}_X^t)$$

and

$$\rho^{-1}R\Gamma_Z\mathcal{O}_X^t \otimes \rho^{-1}R\mathcal{H}om(F, \mathcal{O}_X^w) \rightarrow \rho^{-1}R\mathcal{H}om(F, \mathcal{O}_X^w)$$

when $Z \subset X$ is closed subanalytic and $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$.

We reduce to the case of a real analytic manifold and we use the fact that $\rho^{-1}R\mathcal{H}om(G, \mathcal{C}_X^{\infty, t}) \simeq T\mathcal{H}om(G, \mathcal{C}_X^{\infty})$ and $\rho^{-1}R\mathcal{H}om(G, \mathcal{C}_X^{\infty, w}) \simeq D'G \overset{w}{\otimes} \mathcal{C}_X^{\infty}$ for $G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. Define $F \overset{w}{\otimes} T\mathcal{H}om(G, \mathcal{C}_X^{\infty}) = T\mathcal{H}om(G, F \overset{w}{\otimes} \mathcal{C}_X^{\infty})$ saying that, if U, V are open subanalytic $\mathbb{C}_U \overset{w}{\otimes} T\mathcal{H}om(\mathbb{C}_V, \mathcal{C}_X^{\infty}) = T\mathcal{H}om(\mathbb{C}_V, \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_X^{\infty})$ are $\mathcal{C}^{\infty}$ -functions tempered on V and vanishing up to infinity outside U . Then we have

$$\begin{aligned} T\mathcal{H}om(\mathbb{C}_Z, \mathcal{C}_X^{\infty}) \otimes T\mathcal{H}om(F, \mathcal{C}_X^{\infty}) &\rightarrow T\mathcal{H}om(\mathbb{C}_Z, \mathcal{C}_X^{\infty}) \otimes T\mathcal{H}om(F, \mathbb{C}_Z \overset{w}{\otimes} \mathcal{C}_X^{\infty}) \\ &\rightarrow T\mathcal{H}om(F_Z, \mathcal{C}_X^{\infty}) \\ &\rightarrow T\mathcal{H}om(F, \mathcal{C}_X^{\infty}) \end{aligned}$$

and

$$\begin{aligned} T\mathcal{H}om(\mathbb{C}_Z, \mathcal{C}_X^{\infty}) \otimes D'F \overset{w}{\otimes} \mathcal{C}_X^{\infty} &\rightarrow T\mathcal{H}om(\mathbb{C}_Z, \mathcal{C}_X^{\infty}) \otimes (D'F)_Z \overset{w}{\otimes} \mathcal{C}_X^{\infty} \\ &\rightarrow T\mathcal{H}om(\mathbb{C}_Z, D'F \overset{w}{\otimes} \mathcal{C}_X^{\infty}) \\ &\rightarrow D'F \overset{w}{\otimes} \mathcal{C}_X^{\infty}. \end{aligned}$$

The first and the third arrows of the two diagrams are clearly compatible. Let us see the compatibility between the second arrows. Note that $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ plays no role in these arrows (it denotes a growth conditions which is preserved after the multiplication), so in order to better understand how they are constructed we set $F = \mathbb{C}_X$. Let $U = X \setminus Z$. Then $T\mathcal{H}om(\mathbb{C}_Z, \mathcal{C}_X^{\infty})$ and $\mathbb{C}_Z \overset{w}{\otimes} \mathcal{C}_X^{\infty}$ are represented by the complexes

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathcal{C}_X^{\infty} & \rightarrow & T\mathcal{H}om(\mathbb{C}_U, \mathcal{C}_X^{\infty}) & & \\ \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_X^{\infty} & \rightarrow & \mathcal{C}_X^{\infty} & \rightarrow & 0 & & \end{array}$$

where in both cases $\mathcal{C}_X^{\infty}$ is the degree zero of the complex. The morphism is induced by the following diagram, where the vertical arrows are given by multiplication

$$\begin{array}{ccccc} \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_X^{\infty} \otimes \mathcal{C}_X^{\infty} & \longrightarrow & \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_X^{\infty} \otimes T\mathcal{H}om(\mathbb{C}_U, \mathcal{C}_X^{\infty}) \oplus \mathcal{C}_X^{\infty} \otimes \mathcal{C}_X^{\infty} & \longrightarrow & \mathcal{C}_X^{\infty} \otimes T\mathcal{H}om(\mathbb{C}_U, \mathcal{C}_X^{\infty}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_X^{\infty} & \longrightarrow & \mathbb{C}_U \overset{w}{\otimes} \mathcal{C}_X^{\infty} \oplus \mathcal{C}_X^{\infty} & \longrightarrow & T\mathcal{H}om(\mathbb{C}_U, \mathcal{C}_X^{\infty}). \end{array}$$

In the complex in the second line the first arrow is given by $s \mapsto (s, s)$ and the second one by $(u, v) \mapsto u - v$. Computing the cohomology, it is quasi-isomorphic to $T\mathcal{H}om(\mathbb{C}_Z, \mathcal{C}_X^\infty)$.

7.3 Microlocal integral transformations

In the case of contact transformation the hypothesis of properness of the previous section are not satisfied. Hence we are going to define microlocal operations on $\mu hom^{sa}(\cdot, \mathcal{O}_X^\lambda)$ extending those of [11] and [1].

Let $\Omega \subset T^*X$. Denote by $D^b(X_{sa}, \Omega)$ (resp. $D^b(X, \Omega)$, resp. $D_{\mathbb{R}\text{-c}}^b(X, \Omega)$) the category $D^b(\mathbb{C}_{X_{sa}})/\mathcal{N}_\Omega$ (resp. $D^b(\mathbb{C}_X)/\mathcal{N}_\Omega$, resp. $D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)/\mathcal{N}_\Omega$), where $\mathcal{N}_\Omega = \{F \in D^b(\mathbb{C}_{X_{sa}}) ; SS(F) \cap \Omega = \emptyset\}$ (resp. $F \in D^b(\mathbb{C}_X)$, resp. $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$). It follows from Corollary 5.3.5 that the functor

$$\rho^{-1} \mu hom^{sa} : D^b(X_{sa}, \Omega)^{op} \times D^b(X_{sa}, \Omega) \rightarrow D^b(\Omega)$$

is well defined.

Notations 7.3.1 *If there is no risk of confusion we will write for short $\mu hom(\cdot, \mathcal{O}_X^\lambda)$ instead of $\rho^{-1} \mu hom^{sa}(\cdot, \mathcal{O}_X^\lambda)$.*

Denote by $\circ_\mu$ the microlocal composition of kernels of [11] (and [1] for $\mathbb{R}$ -constructible sheaves). As usual, given $K \in D^b(\mathbb{C}_{(X \times Y)_{sa}})$ and $F \in D^b(\mathbb{C}_{Y_{sa}})$ we set $\Phi_K^\mu F = K \circ_\mu F$.

Proposition 7.3.2 *(i) Let X, Y be two complex analytic manifolds, let $K \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_{X \times Y})$, $p_X \in T^*X$, $p_Y \in T^*Y$ such that $SS(K) \cap (\{p_X\} \times T^*Y) \subseteq (p_X, p_Y^a)$ in a neighborhood of this point. Then for each $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_Y)$ and $G \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ there are morphisms*

$$(7.14) \quad \begin{aligned} \mu hom(K, \mathcal{O}_{X \times Y}^{t(0, d_Y)})_{(p_X, p_Y^a)}[d_Y] &\otimes \mu hom(F, \mathcal{O}_Y^t)_{p_Y} \\ &\rightarrow \mu hom(\Phi_K^\mu F, \mathcal{O}_X^t)_{p_X}. \end{aligned}$$

$$(7.15) \quad \begin{aligned} \mu hom(K, \mathcal{O}_{X \times Y}^{t(0, d_Y)})_{(p_X, p_Y^a)}[d_Y] &\otimes \mu hom(D'(\Phi_K^\mu G), \mathcal{O}_Y^w)_{p_Y} \\ &\rightarrow \mu hom(D'G, \mathcal{O}_X^w)_{p_X}. \end{aligned}$$

*(ii) Let Z be another complex analytic manifold, let $K_1 \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_{X \times Y})$ and $K_2 \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_{Y \times Z})$ be microlocally composable at $(p_X, p_Y, p_Z) \in T^*X \times T^*Y \times T^*Z$, i.e.*

$$(SS(K_1) \times_{T^*Y} SS(K_2)) \cap p_{13}^{a-1}(p_X, p_Z^a) \subseteq \{((p_X, p_Y^a), (p_Y, p_Z^a))\}$$

in a neighborhood of $((p_X, p_Y^a), (p_Y, p_Z^a))$. Then there is a morphism

$$\begin{aligned} & \mu hom(K_1, \mathcal{O}_{X \times Y}^{t(0, d_Y)})_{(p_X, p_Y^a)} \otimes \mu hom(K_2, \mathcal{O}_{Y \times Z}^{t(0, d_Z)})_{(p_Y, p_Z^a)} \\ & \rightarrow \mu hom(K_1 \circ_{\mu} K_2, \mathcal{O}_{X \times Z}^{t(0, d_Z)})_{(p_X, p_Z^a)}[-d_Y] \end{aligned}$$

Proof. The result follows thanks to the morphisms defined in the previous section and adapting the proof of Proposition 3.3.12 of [1]. $\square$

7.4 Contact transformations

Let X, Y be two complex analytic manifolds of the same complex dimension n and let $\Omega_X \subset T^*X$, $\Omega_Y \subset T^*Y$ be two open subanalytic subsets. Let χ be a contact transformation from Ω_X to Ω_Y . We set $\Lambda \subset \Omega_X \times \Omega_Y^a$ be the Lagrangian manifold associated to the graph of χ (i.e. $(p_X, p_Y^a) \in \Lambda$ if $p_Y = \chi(p_X)$). We denote by p_1 and p_2^a the projections from Λ to Ω_X and Ω_Y respectively.

Let $(p_X, p_Y) \in \Omega_X \times \Omega_Y$ and consider $K \in D_{\mathbb{C}\text{-c}}^b(X \times Y, (p_X, p_Y^a))$ satisfying the following properties:

$$(7.16) \quad \begin{cases} SS(K) \subset \Lambda, \\ K \text{ is simple with shift 0 along } \Lambda. \end{cases}$$

In this situation we have the following results of [15] and [1].

Proposition 7.4.1 *Let $K \in D_{\mathbb{C}\text{-c}}^b(X \times Y, (p_X, p_Y^a))$ satisfying (7.16). Set $K^* = r_* R\mathcal{H}om(K, \omega_{X \times Y|Y})$, where $r : X \times Y \rightarrow Y \times X$ is the canonical map. Then the functor $\Phi_K^\mu : D^b(X_{sa}, p_X) \rightarrow D^b(Y_{sa}, p_Y)$ and the functor $\Phi_{K^*}^\mu : D^b(Y_{sa}, p_Y) \rightarrow D^b(X_{sa}, p_X)$ are equivalence of categories inverse to each other.*

Lemma 7.4.2 *Let $K \in D_{\mathbb{C}\text{-c}}^b(X \times Y, (p_X, p_Y^a))$ satisfying (7.16). Then $\mu hom(K, \mathcal{O}_{X \times Y}^t)$ is concentrated in degree zero.*

Proposition 7.4.3 *Let $K \in D_{\mathbb{C}\text{-c}}^b(X \times Y, (p_X, p_Y^a))$ satisfying (7.16) and let $s \in \mu hom(K, \mathcal{O}_{X \times Y}^{t(0, n)})_{(p_X, p_Y^a)}$.*

(i) For each $F \in D_{\mathbb{R}\text{-c}}^b(Y, p_Y)$ and $G \in D_{\mathbb{R}\text{-c}}^b(X, p_X)$ there are morphisms induced by s

$$\begin{aligned}\varphi_s : \mu hom(F, \mathcal{O}_Y^t)_{p_Y}[n] &\rightarrow \mu hom(\Phi_K^\mu F, \mathcal{O}_X^t)_{p_X} \\ \psi_s : \mu hom(D'(\Phi_K^\mu G), \mathcal{O}_Y^w)_{p_Y}[n] &\rightarrow \mu hom(D'F, \mathcal{O}_X^w)_{p_X}.\end{aligned}$$

(ii) Let Z be a n -dimensional complex analytic manifold, $\Omega_Z \subset T^*Z$ and let $\chi' : \Omega_Y \rightarrow \Omega_Z$ be a contact transformation. Let Λ' be the Lagrangian submanifold associated to the graph of χ' . Let $K' \in D_{\mathbb{C}\text{-c}}^b(Y \times Z, (p_Y, p_Z^a))$ satisfying (7.16) and $s \in \mu hom(K', \mathcal{O}_{Y \times Z}^{t(0,n)})$. Then $\varphi_s \circ \varphi'_{s'} = (\varphi \circ \varphi')_{s \circ s'}$ and $\psi_s \circ \psi'_{s'} = (\psi \circ \psi')_{s \circ s'}$, where $s \circ s'$ is the image of $s \otimes s'$ by the morphism

$$\begin{aligned}\mu hom(K, \mathcal{O}_{X \times Y}^t)_{(p_X, p_Y^a)} \otimes \mu hom(K', \mathcal{O}_{Y \times Z}^t)_{(p_Y, p_Z^a)} \\ \rightarrow \mu hom(K \circ_\mu K', \mathcal{O}_{X \times Z}^t)_{(p_X, p_Z^a)}.\end{aligned}$$

(iii) Let $P \in \mathcal{E}_{X, p_X}^{\mathbb{R}, f}$ and $Q \in \mathcal{E}_{Y, p_Y}^{\mathbb{R}, f}$ such that $Ps = sQ$. Then:

$$P \circ \varphi_s = \varphi_s \circ Q$$

(and similarly for ψ_s).

Proof. (i) Similar to Proposition 5.2.1 (i) of [1]. There exists a neighborhood Ω of (p_X, p_Y^a) such that $s \in \Gamma(\Omega, \mu hom(K, \mathcal{O}_{X \times Y}^{t(0,n)})$ and we may suppose that Λ is closed in Ω . Set $\mathcal{K} = \mu hom(K, \mathcal{O}_{X \times Y}^{t(0,n)})$. Then

$$(7.17) \quad s \in \Gamma(\Omega, \mathcal{K}) \simeq \text{Hom}(\mathbb{C}_\Lambda, \mathcal{K}).$$

Moreover we can find relatively compact neighborhoods V_Y and V_X of $\pi_Y(p_Y)$ and $\pi_X(p_X)$ respectively such that $\Phi_K^\mu F = \Phi_{K_{X \times V_Y}} F = K_{X \times V_Y} \circ F$ and $\Phi_K^\mu G = \Phi_{K_{V_X \times Y}} G = K_{V_X \times Y} \circ G$. Now set

$$\begin{aligned}\mathcal{F}_1 &= \mu hom(\Phi_{K_{X \times V_Y}} F, \mathcal{O}_X^t), & \mathcal{G}_1 &= \mu hom(D'G, \mathcal{O}_X^w), \\ \mathcal{F}_2 &= \mu hom(F, \mathcal{O}_Y^t)[n], & \mathcal{G}_2 &= \mu hom(D'(\Phi_{K_{V_X \times Y}} G), \mathcal{O}_Y^w)[n].\end{aligned}$$

Then the morphisms φ_s and ψ_s are given by the diagrams

$$\begin{aligned}\mathcal{F}_2|_{\Omega_Y} &\xrightarrow{\sim} (\mathbb{C}_\Lambda^a \circ \mathcal{F}_2)|_{\Omega_X} \longrightarrow (\mathcal{K}^a \circ \mathcal{F}_2)|_{\Omega_X} \longrightarrow \mathcal{F}_1|_{\Omega_X} \\ \mathcal{G}_2|_{\Omega_Y} &\xrightarrow{\sim} (\mathbb{C}_\Lambda^a \circ \mathcal{G}_2)|_{\Omega_X} \longrightarrow (\mathcal{K}^a \circ \mathcal{G}_2)|_{\Omega_X} \longrightarrow \mathcal{G}_1|_{\Omega_X}\end{aligned}$$

where the first arrows are given by (7.17) and the second ones by (7.14) and (7.15).

- (ii) The arrow follows from (i) and the associativity of the composition.
- (iii) See [1], Proposition 5.2.1 (iii).

□

Theorem 7.4.4 *Let χ be a contact transformation from Ω_X to Ω_Y and let Λ be the Lagrangian manifold associated to the graph of χ . Then there exists $K \in D_{\mathbb{C}\text{-c}}^b(X \times Y, (p_X, p_Y^a))$ satisfying (7.16) and $s \in \mu\text{hom}(K, \mathcal{O}_{X \times Y}^{t(0,n)})_{(p_X, p_Y^a)}$ such that:*

- (i) *the correspondence $\mathcal{E}_{X, p_X} \ni P \mapsto Q \in \mathcal{E}_{Y, p_Y}$ such that $Ps = sQ$ is an isomorphism of rings,*
- (ii) *for each $F \in D_{\mathbb{R}\text{-c}}^b(Y, p_Y)$ and $G \in D_{\mathbb{R}\text{-c}}^b(X, p_X)$ the morphisms induced by s*

$$\begin{aligned} \varphi_s : \mu\text{hom}(F, \mathcal{O}_Y^t)_{p_Y}[n] &\rightarrow \mu\text{hom}(\Phi_K^\mu F, \mathcal{O}_X^t)_{p_X} \\ \psi_s : \mu\text{hom}(D'(\Phi_K^\mu G), \mathcal{O}_Y^w)_{p_Y}[n] &\rightarrow \mu\text{hom}(D'G, \mathcal{O}_X^w)_{p_X}. \end{aligned}$$

are isomorphisms compatible with (i).

Proof. The proof is similar to the proof of Proposition 5.2.2 of [1].

□

Remark 7.4.5 *Set $G = D'\Phi_K^\mu F$ with $F \in D_{\mathbb{R}\text{-c}}^b(Y, p_Y)$, then $D'G = \Phi_K^\mu F$ and*

$$D'\Phi_K^\mu G \simeq D'\Phi_K^\mu D'\Phi_K^\mu F \simeq \Psi_K^\mu D'D'\Phi_K^\mu F \simeq \Psi_K^\mu \Phi_K^\mu F \simeq F,$$

where the last isomorphism follows from Theorem 7.1.2 of [11]. Hence we obtain the isomorphism

$$\mu\text{hom}(F, \mathcal{O}_Y^w)_{p_Y}[n] \xrightarrow{\sim} \mu\text{hom}(\Phi_K^\mu F, \mathcal{O}_X^w)_{p_X}.$$

8 Cauchy-Kowaleskaya-Kashiwara theorem

Here we prove the Cauchy-Kowaleskaya-Kashiwara theorem for holomorphic functions with growth conditions. The idea of the proof is the following: we divide the proof in two parts. In the first one we prove that for holomorphic

functions with growth conditions the characteristic variety of a coherent $\mathcal{D}$ -module coincides with the microsupport of the complex of solutions. In the second part we use inverse image formulas for $\mathcal{O}_Y^\lambda$ to finish the proof of the theorem.

8.1 Microsupport and characteristic variety

We are going to study the relation between microsupport of subanalytic sheaves and the characteristic variety of a $\mathcal{D}$ -module.

Lemma 8.1.1 *Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ and let $G \in D^b(\mathbb{C}_X)$. Then*

$$D'F \otimes \rho_! G \simeq R\mathcal{H}om(F, \rho_! G).$$

Proof. It follows from the following isomorphism of [23]: let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$, $K \in D^b(\mathbb{C}_{X_{sa}})$ and $G \in D^b(\mathbb{C}_X)$. Then

$$R\mathcal{H}om(F, K) \otimes \rho_! G \simeq R\mathcal{H}om(F, K \otimes \rho_! G).$$

Setting $K = \mathbb{C}_X$ we obtain the result. □

Let us recall the notion of elliptic pair of [27]. Let $\mathcal{M}$ be a coherent $\mathcal{D}$ -module and $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$, then $(F, \mathcal{M})$ is an elliptic pair if

$$SS(F) \cap \text{Char}(\mathcal{M}) \subseteq T_X^* X.$$

We consider the sheaf $\mathcal{O}_X^\lambda$, for $\lambda = \emptyset, t, w, \omega$.

Proposition 8.1.2 *Let $(F, \mathcal{M})$ be an elliptic pair. Let $G \in D^b(\rho_! \mathcal{D}_X)$ such that $H^k \mu hom(F, G)$ is a $\mathcal{E}_X$ -module for each $k \in \mathbb{Z}$. Then we have the isomorphism*

$$R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, D'F \otimes \rho^{-1} G) \xrightarrow{\sim} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \rho^{-1} R\mathcal{H}om(F, G)).$$

Proof. Let $\delta : \Delta \rightarrow X \times X$ be the embedding and let us consider the Sato's triangle

$$\begin{aligned} \delta^{-1} R\mathcal{H}om(q_1^{-1} F, q_2^! G) \otimes \omega_{\Delta|X \times X} &\rightarrow \delta^! R\mathcal{H}om(q_1^{-1} F, q_2^! G) \\ (8.1) \quad &\rightarrow R\dot{\pi}_* \mu hom^{sa}(F, G) \xrightarrow{+}. \end{aligned}$$

We have $\delta^! R\mathcal{H}om(q_1^{-1}F, q_2^!G) \simeq R\mathcal{H}om(F, G)$. Moreover

$$\begin{aligned} \delta^{-1} R\mathcal{H}om(q_1^{-1}F, q_2^!G) \otimes \omega_{\Delta|X \times X} &\simeq \delta^{-1} R\mathcal{H}om(q_1^{-1}F, q_2^{-1}G) \\ &\simeq \delta^{-1}(D'F \boxtimes G) \\ &\simeq D'F \otimes G \end{aligned}$$

where the second isomorphism follows from Lemma 5.3.8. Hence applying ρ^{-1} we obtain

$$D'F \otimes \rho^{-1}G \rightarrow \rho^{-1} R\mathcal{H}om(F, G) \rightarrow R\dot{\pi}_* \mu hom(F, G) \xrightarrow{\pm}.$$

Applying the functor $R\mathcal{H}om_{\mathcal{D}_X}(\rho^! \mathcal{M}, \cdot)$ we obtain

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, D'F \otimes \rho^{-1}G) &\rightarrow R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \rho^{-1} R\mathcal{H}om(F, G)) \\ &\rightarrow R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, R\dot{\pi}_* \mu hom(F, G)) \xrightarrow{\pm}. \end{aligned}$$

Then it is enough to prove that $R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, R\dot{\pi}_* \mu hom(F, G)) = 0$. First remark that we have by adjunction

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, R\dot{\pi}_* \mu hom(F, G)) \\ \simeq R\dot{\pi}_* R\mathcal{H}om_{\pi^{-1}\mathcal{D}_X}(\pi^{-1}\mathcal{M}, \mu hom(F, G)). \end{aligned}$$

Let $k \in \mathbb{Z}$. Since $H^k \mu hom(F, G)$ is a $\mathcal{E}_X$ -module for each $k \in \mathbb{Z}$ we have

$$\begin{aligned} R\dot{\pi}_* R\mathcal{H}om_{\pi^{-1}\mathcal{D}_X}(\pi^{-1}\mathcal{M}, H^k \mu hom(F, G)) \\ \simeq R\dot{\pi}_* R\mathcal{H}om_{\mathcal{E}_X}((\mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1}\mathcal{M}), H^k \mu hom(F, G)). \end{aligned}$$

We have $\text{supp}(H^k \mu hom(F, G)) \subseteq \text{SS}(F)$ for each $k \in \mathbb{Z}$ by Corollary 5.3.5 and $\text{supp}(\mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1}\mathcal{M}) = \text{Char}(\mathcal{M})$. Hence

$$(8.2) \quad R\dot{\pi}_* R\mathcal{H}om_{\pi^{-1}\mathcal{D}_X}(\pi^{-1}\mathcal{M}, H^k \mu hom(F, G)) = 0$$

for each $k \in \mathbb{Z}$ since the pair $(F, \mathcal{M})$ is elliptic. Let us suppose that the length of the bounded complex $\mu hom(F, G)$ is n and let us argue by induction on the truncation $\tau^{\leq i} \mu hom(F, G)$. If $i = 0$ the result follows from (8.2). Let us consider the distinguish triangle

$$\tau^{\leq n-1} \mu hom(F, G) \rightarrow \mu hom(F, G) \rightarrow H^n \mu hom(F, G) \xrightarrow{\pm}$$

and apply the functor $R\dot{\pi}_* R\mathcal{H}om_{\pi^{-1}\mathcal{D}_X}(\pi^{-1}\mathcal{M}, \cdot)$. The first term becomes zero by the induction hypothesis and the third one is zero by (8.2). Hence

$R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, R\pi_*\mu hom(F, G)) = 0$ and the result follows. $\square$

Setting $G = \mathcal{O}_X^\lambda$ in Proposition 8.1.2, $\lambda = \emptyset, t, w, \omega$ we obtain the following result.

Theorem 8.1.3 *Let $(F, \mathcal{M})$ be an elliptic pair. Then we have the isomorphism*

$$(8.3) \quad R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, D'F \otimes \mathcal{O}_X) \xrightarrow{\sim} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \rho^{-1}R\mathcal{H}om(F, \mathcal{O}_X^\lambda)).$$

Proof. If $\lambda = \omega$ this is a consequence of Lemma 8.1.1. If $\lambda = \emptyset, t, w$ then $H^k\mu hom(F, \mathcal{O}_X^\lambda)$ is a $\mathcal{E}_X$ -module for each $k \in \mathbb{Z}$ by Corollaries 7.2.5 and 7.2.18 and the result follows from Proposition 8.1.3. $\square$

Example 8.1.4 *Let M be a real analytic manifold and let X be a complexification of M . Let $\mathcal{M}$ be an elliptic system on M . Then*

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{A}_M) &\simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{C}_M^\infty) \\ &\simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{D}b_M) \\ &\simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{B}_M). \end{aligned}$$

This follows from Theorem 8.1.3 setting $F = D'\mathbb{C}_M$ and applying ρ^{-1} to the isomorphism (8.3). In fact

$$R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \rho^{-1}R\mathcal{H}om(D'\mathbb{C}_M, \mathcal{O}_X^\lambda)) \simeq \begin{cases} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{A}_M) & \text{if } \lambda = \omega, \\ R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{C}_M^\infty) & \text{if } \lambda = w, \\ R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{D}b_M) & \text{if } \lambda = t, \\ R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{B}_M) & \text{if } \lambda = \emptyset \end{cases}$$

and they are all isomorphic since we have $\rho^{-1}R\mathcal{H}om_{\rho!\mathcal{D}_X}(\rho!\mathcal{M}, D'D'\mathbb{C}_M \otimes \mathcal{O}_X^\lambda) \simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathbb{C}_M \otimes \mathcal{O}_X) \simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{A}_M)$ for $\lambda = \emptyset, t, w, \omega$.

Example 8.1.5 *Let X be a smooth submanifold of Y , let $\mathcal{O}_Y|_X$ be the formal completion of X along Y and let $\mathcal{B}_{X|Y}$ be the algebraic cohomology of $\mathcal{O}_Y$ with support in X . Let $\mathcal{M}$ be a $\mathcal{D}_Y$ -module such that $T_X^*Y \cap \text{Char}(\mathcal{M}) \subseteq T_Y^*Y$. Then we have the isomorphisms*

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{O}_Y|_X) &\simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \widehat{\mathcal{O}_Y|_X}), \\ &\simeq R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{B}_{X|Y}). \end{aligned}$$

This follows from Theorem 8.1.3 setting $F = D'\mathbb{C}_Y$, applying ρ^{-1} to the isomorphism (8.3) and arguing as in Example 8.1.4.

Let $\mathcal{M}$ be a $\mathcal{D}_X$ -module and let $\lambda = \emptyset, t, w, \omega$. One sets for short

$$\text{Sol}^\lambda(\mathcal{M}) := R\mathcal{H}om_{\rho_!\mathcal{D}_X}(\rho_!\mathcal{M}, \mathcal{O}_X^\lambda).$$

Corollary 8.1.6 *Let $\mathcal{M}$ be a coherent $\mathcal{D}_X$ -module. Then*

$$SS(\text{Sol}^\lambda(\mathcal{M})) = \text{Char}(\mathcal{M}).$$

Proof. Recall that $SS(\text{Sol}(\mathcal{M})) = \text{Char}(\mathcal{M})$.

(i) $\text{Char}(\mathcal{M}) \subseteq SS(\text{Sol}^\lambda(\mathcal{M}))$ follows from the fact that $\rho^{-1}\text{Sol}^\lambda(\mathcal{M}) = \text{Sol}(\mathcal{M})$ and $SS(\rho^{-1}G) \subseteq SS(G)$ for each $G \in D^b(\mathbb{C}_{X_{sa}})$.

(ii) $\text{Char}(\mathcal{M}) \supseteq SS(\text{Sol}^\lambda(\mathcal{M}))$. Let $(x, \xi) \notin \text{Char}(\mathcal{M}) = SS(\text{Sol}(\mathcal{M}))$ and let U be a conic open neighborhood of (x, ξ) , $U \cap \text{Char}(\mathcal{M}) = \emptyset$, such that for each $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ with $\text{supp}(F) \subset\subset \pi(U)$ and $SS(F) \subset U \cup T_X^*X$ we have $\text{Hom}_{D^b(\mathbb{C}_X)}(F, \text{Sol}(\mathcal{M})) = 0$. By Theorem 8.1.3 the complexes

$$\begin{aligned} R\mathcal{H}om(F, R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{O}_X)) &\simeq \rho^{-1}R\mathcal{H}om(F, R\mathcal{H}om_{\rho_!\mathcal{D}_X}(\rho_!\mathcal{M}, R\rho_*\mathcal{O}_X)) \\ &\simeq \rho^{-1}R\mathcal{H}om(F, R\mathcal{H}om_{\rho_!\mathcal{D}_X}(\rho_!\mathcal{M}, \mathcal{O}_X^\lambda)) \end{aligned}$$

are all quasi-isomorphic for $\lambda = \emptyset, t, w, \omega$. Hence $\text{Hom}_{D^b(\mathbb{C}_{X_{sa}})}(F, \text{Sol}^\lambda(\mathcal{M})) = 0$ and $(x, \xi) \notin SS(\text{Sol}^\lambda(\mathcal{M}))$. □

8.2 Cauchy-Kowaleskaya-Kashiwara theorem

Now we apply the preceding results to prove the Cauchy-Kowaleskaya-Kashiwara theorem for holomorphic functions with growth conditions $\lambda = \emptyset, t, w, \omega$. We refer to [10] for the statement and proof of the Cauchy-Kowaleskaya-Kashiwara theorem for holomorphic functions.

Let $f : X \rightarrow Y$ be a morphism of complex manifolds. Set $d = \dim_{\mathbb{C}} X - \dim_{\mathbb{C}} Y$. We recall the inverse image isomorphisms

$$(8.4) \quad f^!\mathcal{O}_Y^t \simeq \rho_!\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes}_{\rho_!\mathcal{D}_X} \mathcal{O}_X^t[d],$$

$$(8.5) \quad f^!\mathcal{O}_Y^w \simeq R\mathcal{H}om_{\rho_!\mathcal{D}_X}(\rho_!\mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w)[2d].$$

Proposition 8.2.1 *Let $\mathcal{M}$ be a coherent $\mathcal{D}_Y$ -module, and suppose that f is non characteristic for $\mathcal{M}$. Then we have the following isomorphism for $\lambda = \emptyset, t, w, \omega$:*

$$f^! R\mathcal{H}om_{\rho_! \mathcal{D}_Y}(\rho_! \mathcal{M}, \mathcal{O}_Y^\lambda) \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \underline{f}^{-1} \mathcal{M}, \mathcal{O}_X^\lambda)[2d].$$

Proof. (i) Let $\lambda = t$. Recall that if $\mathcal{M}$ is a coherent $\mathcal{D}_Y$ -module and f is non characteristic, then $\underline{f}^{-1} \mathcal{M}$ is a coherent $\mathcal{D}_X$ -module and

$$\underline{f}^{-1} R\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{M}, \mathcal{D}_Y) \simeq R\mathcal{H}om_{\mathcal{D}_X}(\underline{f}^{-1} \mathcal{M}, \mathcal{D}_X)[d].$$

We have the chain of isomorphisms

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(\rho_! \underline{f}^{-1} \mathcal{M}, \mathcal{O}_X^t)[2d] &\simeq \rho_! R\mathcal{H}om_{\mathcal{D}_X}(\underline{f}^{-1} \mathcal{M}, \mathcal{D}_X) \otimes_{\rho_! \mathcal{D}_X} \mathcal{O}_X^t[2d] \\ &\simeq \rho_! \underline{f}^{-1} R\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{M}, \mathcal{D}_Y) \otimes_{\rho_! \mathcal{D}_X} \mathcal{O}_X^t[d] \\ &\simeq \rho_! f^{-1} R\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{M}, \mathcal{D}_Y) \otimes_{\rho_! f^{-1} \mathcal{D}_Y} f^! \mathcal{O}_Y^t \\ &\simeq f^! (\rho_! R\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{M}, \mathcal{D}_Y) \otimes_{\rho_! \mathcal{D}_Y} \mathcal{O}_Y^t) \\ &\simeq f^! R\mathcal{H}om_{\rho_! \mathcal{D}_Y}(\rho_! \mathcal{M}, \mathcal{O}_Y^t), \end{aligned}$$

where the first and the last isomorphisms follow from the coherence of $\underline{f}^{-1} \mathcal{M}$ and $\mathcal{M}$, and the third one follows from (8.4).

(ii) Let $\lambda = w$. We have the chain of isomorphisms

$$\begin{aligned} f^! R\mathcal{H}om_{\rho_! \mathcal{D}_Y}(\mathcal{M}, \mathcal{O}_Y^w) &\simeq R\mathcal{H}om_{\rho f^{-1} \mathcal{D}_X}(\rho_! f^{-1} \mathcal{M}, f^! \mathcal{O}_Y^w) \\ &\simeq R\mathcal{H}om_{\rho_! f^{-1} \mathcal{D}_X}(\rho_! f^{-1} \mathcal{M}, R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w))[2d] \\ &\simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \underline{f}^{-1} \mathcal{M}, \mathcal{O}_X^w)[2d], \end{aligned}$$

where the second isomorphism follows from (8.5).

(iii) Let $\lambda = \emptyset, \omega$. Since $\mathcal{M}$ is coherent and f is non characteristic the result follows from the isomorphism

$$f^! R\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{M}, \mathcal{O}_Y) \simeq R\mathcal{H}om_{\mathcal{D}_X}(\underline{f}^{-1} \mathcal{M}, \mathcal{O}_X)[2d].$$

□

Theorem 8.2.2 *Let $\mathcal{M}$ be a coherent $\mathcal{D}_Y$ -module, and suppose that f is non characteristic for $\mathcal{M}$. Then we have the following isomorphism for $\lambda = \emptyset, t, w, \omega$:*

$$f^{-1} R\mathcal{H}om_{\rho_! \mathcal{D}_Y}(\rho_! \mathcal{M}, \mathcal{O}_Y^\lambda) \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \underline{f}^{-1} \mathcal{M}, \mathcal{O}_X^\lambda).$$

Proof. By Corollary 8.1.6 f is non characteristic for $SS(Sol^\lambda(\mathcal{M}))$. Hence by Proposition 5.3.7

$$f^! R\mathcal{H}om_{\rho_! \mathcal{D}_Y}(\rho_! \mathcal{M}, \mathcal{O}_Y^\lambda) \simeq f^{-1} R\mathcal{H}om_{\rho_! \mathcal{D}_Y}(\rho_! \mathcal{M}, \mathcal{O}_Y^\lambda)[2d].$$

Then the result follows from Proposition 8.2.1. □

A Appendix

A.1 Review on subanalytic sets

We recall briefly some properties of subanalytic subsets. Reference are made to [3] and [18] for the theory of subanalytic subsets and to [4] and [31] for the more general theory of o-minimal structures. Let X be a real analytic manifold.

Definition A.1.1 *Let A be a subset of X .*

- (i) *A is said to be semi-analytic if it is locally analytic, i.e. each $x \in A$ has a neighborhood U such that $X \cap U = \cup_{i \in I} \cap_{j \in J} X_{ij}$, where I, J are finite sets and either $X_{ij} = \{y \in U_x; f_{ij} > 0\}$ or $X_{ij} = \{y \in U_x; f_{ij} = 0\}$ for some analytic function f_{ij} .*
- (ii) *A is said to be subanalytic if it is locally a projection of a relatively compact semi-analytic subset, i.e. each $x \in A$ has a neighborhood U such that there exists a real analytic manifold Y and a relatively compact semi-analytic subset $A' \subset X \times Y$ satisfying $X \cap U = \pi(A')$, where $\pi : X \times Y \rightarrow X$ denotes the projection.*
- (iii) *Let Y be a real analytic manifold. A continuous map $f : X \rightarrow Y$ is subanalytic if its graph is subanalytic in $X \times Y$.*

Let us recall some result on subanalytic subsets.

Proposition A.1.2 *Let A, B be subanalytic subsets of X . Then $A \cup B$, $A \cap B$, $\overline{A}$, ∂A and $A \setminus B$ are subanalytic.*

Proposition A.1.3 *Let A be a subanalytic subsets of X . Then the connected components of A are locally finite.*

Proposition A.1.4 *Let $f : X \rightarrow Y$ be a subanalytic map. Let A be a relatively compact subanalytic subset of X . Then $f(A)$ is subanalytic.*

Definition A.1.5 *A simplicial complex (K, Δ) is the data consisting of a set K and a set Δ of subsets of K satisfying the following axioms:*

- S1 any $\sigma \in \Delta$ is a finite and non-empty subset of K ,*
- S2 if τ is a non-empty subset of an element σ of Δ , then τ belongs to Δ ,*
- S3 for any $p \in K$, $\{p\}$ belongs to Δ ,*
- S4 for any $p \in K$, the set $\{\sigma \in \Delta; p \in \sigma\}$ is finite.*

If (K, Δ) is a simplicial complex, an element of K is called a vertex. Let $\mathbb{R}^K$ be the set of maps from K to $\mathbb{R}$ equipped with the product topology. To $\sigma \in \Delta$ one associate $|\sigma| \subset \mathbb{R}^K$ as follows:

$$|\sigma| = \left\{ x \in \mathbb{R}^K; x(p) = 0 \text{ for } p \notin \sigma, x(p) > 0 \text{ for } p \in \sigma \text{ and } \sum_p x(p) = 1 \right\}.$$

As usual we set:

$$|K| = \bigcup_{\sigma \in \Delta} |\sigma|,$$

$$U(\sigma) = \bigcup_{\tau \in \Delta, \tau \supset \sigma} |\tau|,$$

and for $x \in |K|$:

$$U(x) = U(\sigma(x)),$$

where $\sigma(x)$ is the unique simplex such that $x \in |\sigma|$.

Theorem A.1.6 *Let $X = \bigsqcup_{i \in I} X_i$ be a locally finite partition of X consisting of subanalytic subsets. Then there exists a simplicial complex (K, Δ) and a subanalytic homeomorphism $\psi : |K| \xrightarrow{\sim} X$ such that*

- (i) for any $\sigma \in \Delta$, $\psi(|\sigma|)$ is a subanalytic submanifold of X ,*
- (ii) for any $\sigma \in \Delta$ there exists $i \in I$ such that $\psi(|\sigma|) \subset X_i$.*

Let us recall the definition of a subfamily of the subanalytic subsets of $\mathbb{R}^n$ which have some very good properties.

Definition A.1.7 A subanalytic subset A of $\mathbb{R}^n$ is said to be globally subanalytic if it is subanalytic in the projective space $\mathbb{P}^n(\mathbb{R})$. Here we identify $\mathbb{R}^n$ with a submanifold of $\mathbb{P}^n(\mathbb{R})$ via the map $(x_1, \dots, x_n) \mapsto (1 : x_1 : \dots : x_n)$.

An equivalent way to define globally subanalytic subsets is by means of the map $\tau_n : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by

$$\tau_n(x_1, \dots, x_n) := \left(\frac{x_1}{\sqrt{1+x_1^2}}, \dots, \frac{x_n}{\sqrt{1+x_n^2}} \right).$$

In particular relatively compact subanalytic subsets are globally subanalytic.

Definition A.1.8 A map $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be globally subanalytic if its graph is globally subanalytic.

Proposition A.1.9 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a globally subanalytic map. Let A be a globally subanalytic subset of $\mathbb{R}^n$. Then $f(A)$ is globally subanalytic.

Now we recall the notion of cylindrical cell decomposition, a useful tool to study the geometry of a subanalytic subset. We refer to [4] and [31] for a complete exposition.

A cylindrical cell decomposition (ccd for short) of $\mathbb{R}^n$ is a finite partition of $\mathbb{R}^n$ into subanalytic subsets, called the cells of the ccd. It is defined by induction on n :

$n=1$ a ccd of $\mathbb{R}$ is given by a finite subdivision $a_1 < \dots < a_\ell$ of $\mathbb{R}$. The cells of $\mathbb{R}$ are the points $\{a_i\}$, $1 \leq i \leq \ell$, and the intervals (a_i, a_{i+1}) , $0 \leq i \leq \ell$, where $a_0 = -\infty$ and $a_{\ell+1} = +\infty$.

$n>1$ a ccd of $\mathbb{R}^n$ is given by a ccd of $\mathbb{R}^{n-1}$ and, for each cell D of $\mathbb{R}^{n-1}$, continuous analytic functions

$$\zeta_{D,1} < \dots < \zeta_{D,\ell_D} : D \rightarrow \mathbb{R}.$$

The cells of $\mathbb{R}^n$ are the graphs

$$\{(x, \zeta_{D,i}(x)); x \in D\}, \quad 1 \leq i \leq \ell_D,$$

and the bands

$$\{(x, y) \in D \times \mathbb{R}; \zeta_{D,i}(x) < y < \zeta_{D,i+1}(x)\}$$

for $0 \leq i \leq \ell_D$, where $\zeta_{D,0} = -\infty$ and $\zeta_{D,\ell_D+1} = +\infty$.

Theorem A.1.10 *Let $A_1, \dots, A_k$ be globally subanalytic subsets of $\mathbb{R}^n$. There exists a ccd of $\mathbb{R}^n$ such that each A_i is a union of cells.*

We end this section with the following useful result.

Lemma A.1.11 *Let U be a globally subanalytic subset of $\mathbb{R}^n$ and denote by $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ the projection. Then U admits a finite open covering $\{U_i\}$ such that each U_i is simply connected and the intersection of each U_i with the fibers of π is contractible or empty.*

Proof. Up to take the image of U by the homeomorphism

$$\begin{aligned} \varphi : \mathbb{R}^n &\rightarrow (-1, 1)^n \\ (x_1, \dots, x_n) &\mapsto \left(\frac{x_1}{\sqrt{1+x_1^2}}, \dots, \frac{x_n}{\sqrt{1+x_n^2}} \right) \end{aligned}$$

we may assume that U is bounded. Then it follows from a result of [34] that U can be covered by finitely many open cells, and cells satisfy the desired properties. □

A.2 Ind-sheaves and subanalytic sites

Let us recall some results of [14].

One denotes by $I(k_X)$ the category of ind-sheaves of k -vector spaces on X , that is $I(k_X) = \text{Ind}(\text{Mod}^c(k_X))$, where $\text{Mod}^c(k_X)$ denotes the full subcategory of $\text{Mod}(k_X)$ consisting of sheaves with compact support on X . We denote by $D^b(I(k_X))$ the bounded derived category of $I(k_X)$.

There are three functors relating ind-sheaves and classical sheaves:

$$\begin{aligned} \iota : \text{Mod}(k_X) &\rightarrow I(k_X) & F &\mapsto \varinjlim_{U \subset \subset X} F_U, \\ \alpha : I(k_X) &\rightarrow \text{Mod}(k_X) & \varinjlim_i F_i &\mapsto \varinjlim_i F_i, \\ \beta : \text{Mod}(k_X) &\rightarrow I(k_X) & &\text{left adjoint to } \alpha. \end{aligned}$$

These functors satisfy the following properties:

- the functor ι is fully faithful, exact and commutes with $\varprojlim$,

- the functor α is exact and commutes with $\varinjlim$ and $\varprojlim$,
- the functor β is fully faithful, exact and commute with $\varinjlim$,
- (α, ι) and (β, α) are pairs of adjoint functors.

Since ι is fully faithful and exact we identify $\text{Mod}(k_X)$ (resp. $D^b(k_X)$) with a full abelian subcategory of $\text{I}(k_X)$ (resp. $D^b(\text{I}(k_X))$).

The category $\text{I}(k_X)$ admits an internal hom denoted by $\mathcal{I}hom$ and this functor admits a left adjoint, denoted by $\otimes$. One can also define an external $\mathcal{H}om : \text{I}(k_X) \times \text{I}(k_X) \rightarrow \text{Mod}(k_X)$ and one has

$$\mathcal{H}om(F, G) = \alpha \mathcal{I}hom(F, G) \quad \text{and} \quad \text{Hom}_{\text{I}(k_X)}(F, G) = \Gamma(X; \mathcal{H}om(F, G)).$$

The functor $\otimes$ is exact while $\mathcal{I}hom$ and $\mathcal{H}om$ are left exact and admit right derived functors $R\mathcal{I}hom$ and $R\mathcal{H}om$.

Consider a morphism of real analytic manifolds $f : X \rightarrow Y$. One defines the external operations

$$\begin{aligned} f^{-1} : \text{I}(k_Y) &\rightarrow \text{I}(k_X) & \varinjlim_i G_i &\mapsto \varinjlim_{i, U \subset \subset X} (f^{-1} G_i)_U \\ f_* : \text{I}(k_X) &\rightarrow \text{I}(k_Y) & \varinjlim_i F_i &\mapsto \varprojlim_{U \subset \subset X} \varinjlim_i f_* \Gamma_U F_i \\ f_{!!} : \text{I}(k_X) &\rightarrow \text{I}(k_Y) & \varinjlim_i F_i &\mapsto \varinjlim_i f_! F_i \end{aligned}$$

where the notation $f_{!!}$ is chosen to stress the fact that $f_{!!} \circ \iota \neq \iota \circ f_!$ in general.

While f^{-1} is exact, the others functors admit right derived functors. One can show that the functor $Rf_{!!}$ admits a right adjoint denoted by $f^!$ and we get the usual formalism of the six Grothendieck operations. Almost all the formulas of the classic theory of sheaves remain valid for ind-sheaves.

There is a strict relations between ind-sheaves and sheaves on the subanalytic site associated to X . Set $\text{I}_{\mathbb{R}\text{-c}}(k_X) = \text{Ind}(\text{Mod}_{\mathbb{R}\text{-c}}^c(k_X))$ for short.

Theorem A.2.1 *One has an equivalence of categories*

$$\begin{aligned} \text{I}_{\mathbb{R}\text{-c}}(k_X) &\xrightarrow{\sim} \text{Mod}(k_{X_{sa}}) \\ \varinjlim_i F_i &\mapsto \varinjlim_i \rho_* F_i. \end{aligned}$$

Let us recall the following functor defined in [14]:

$$\begin{aligned} I_{\mathcal{T}} : \text{Mod}(k_{X_{sa}}) &\rightarrow \text{I}(k_X) \\ \varinjlim_i \rho_* F_i &\mapsto \varinjlim_i F_i. \end{aligned}$$

It is fully faithful, exact and commutes with $\varinjlim$ and $\otimes$. It admits a right adjoint

$$J_{\mathcal{T}} : \mathbf{I}(k_X) \rightarrow \mathbf{Mod}(k_{X_{sa}})$$

satisfying, for each $U \in \mathbf{Op}(X_{sa})$, $\Gamma(U; J_{\mathcal{T}}F) = \mathbf{Hom}_{\mathbf{I}(k_X)}(k_U, F)$. This functor is right exact and commutes with filtrant inductive limits. Moreover we have $RJ_{\mathcal{T}} \circ I_{\mathcal{T}} \simeq \text{id}$ and

$$RJ_{\mathcal{T}} R\mathcal{I}hom(I_{\mathcal{T}}F, G) \simeq R\mathcal{H}om(F, RJ_{\mathcal{T}}G).$$

We have the following relations:

$$\begin{aligned} RJ_{\mathcal{T}} \circ \iota &\simeq R\rho_* \quad \text{and} \quad \alpha \simeq \rho^{-1} \circ J_{\mathcal{T}} \\ \alpha \circ I_{\mathcal{T}} &\simeq \rho^{-1} \quad \text{and} \quad I_{\mathcal{T}} \circ \rho_! \simeq \beta \end{aligned}$$

Let $f : X \rightarrow Y$ be a morphism of real analytic manifolds and let U be an open subanalytic subset of X .

Lemma A.2.2 *Let $F \in D^b(k_{X_{sa}})$ and $G \in D^b(k_{Y_{sa}})$. We have*

- (i) $I_{\mathcal{T}} \circ Rf_{!!}F \simeq Rf_{!!} \circ I_{\mathcal{T}}F$,
- (ii) $I_{\mathcal{T}} \circ f^{-1}G \simeq f^{-1} \circ I_{\mathcal{T}}G$,
- (iii) $I_{\mathcal{T}} \circ f^!G \simeq f^! \circ I_{\mathcal{T}}G$,
- (iv) $I_{\mathcal{T}}F_U \simeq (I_{\mathcal{T}}F)_U$,
- (v) $I_{\mathcal{T}} \circ R\Gamma_U F \simeq R\Gamma_U \circ I_{\mathcal{T}}F$.

A.3 Inverse image for tempered holomorphic functions

Let $f : M \rightarrow N$ be a morphism of oriented real analytic manifolds of dimension d_M and d_N . Set $d = d_N - d_M$. Denote by $\mathcal{A}_M$ the sheaf of analytic functions on M .

Lemma A.3.1 *Let F be an $\mathcal{A}_M$ -module locally free of finite rank. Then $R^k f_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! F) = 0$ for $k \neq 0$.*

Proof. It is a consequence of the fact that $\mathcal{D}b_M^t$ is quasi-injective and Proposition 1.6.5 of [23]. □

Lemma A.3.2 *Let M and N be orientable real manifolds. There is a natural morphism of complexes*

$$f_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! \Omega_M^\bullet)[d_M] \rightarrow \mathcal{D}b_N^t \otimes_{\rho_! \mathcal{A}_N} \rho_! \Omega_N^\bullet[d_N].$$

Proof. Let $U \in \text{Op}^c(N_{sa})$. We have the chain of morphisms

$$\begin{aligned} \Gamma(U; f_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! \Omega_M^{d_M-i})) &\simeq \Gamma(N; \mathcal{H}om(\mathbb{C}_U, f_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! \Omega_M^{d_M-i}))) \\ &\simeq \Gamma(N; f_! T\mathcal{H}om(f^{-1} \mathbb{C}_U, \mathcal{D}b_M \otimes_{\mathcal{A}_M} \Omega_M^{d_M-i})) \\ &\rightarrow \Gamma_c(N; T\mathcal{H}om(G, \mathcal{D}b_N \otimes_{\mathcal{A}_N} \Omega_N^{d_N-i})) \\ &\simeq \Gamma(U; \mathcal{D}b_N \otimes_{\mathcal{A}_N} \Omega_N^{d_N-i}), \end{aligned}$$

where the third morphism follows from Proposition 4.3 of [13]. $\square$

Proposition A.3.3 *There is a natural morphism in $D^b(\rho_! \mathcal{D}_M^{op})$:*

$$(A.1) \quad Rf_{!!}(\mathcal{D}b_M^{t\vee} \otimes_{\rho_! \mathcal{D}_M}^L \rho_! \mathcal{D}_{M \rightarrow N}) \rightarrow \mathcal{D}b_N^{t\vee}.$$

Proof. The Spencer resolution of $\mathcal{D}_{M \rightarrow N}$ give rise to the quasi-isomorphism

$$\mathcal{D}_{M \rightarrow N} \xleftarrow{\sim} \mathcal{D}_M \otimes_{\mathcal{A}_M} \bigwedge^\bullet \Theta_M \otimes_{\mathcal{A}_M} \mathcal{D}_{M \rightarrow N} \simeq \mathcal{D}_M \otimes_{\mathcal{A}_M} \bigwedge^\bullet \Theta_M \otimes_{f^{-1} \mathcal{A}_M} f^{-1} \mathcal{D}_N$$

from which we obtain the following quasi-isomorphism for $\mathcal{D}b_M^{t\vee} \otimes_{\rho_! \mathcal{D}_M} \rho_! \mathcal{D}_{M \rightarrow N}$ in $D^b(\rho_! f^{-1} \mathcal{D}_N^{op})$:

$$\begin{aligned} \mathcal{D}b_M^{t\vee} \otimes_{\rho_! \mathcal{A}_M} \rho_! \mathcal{D}_{M \rightarrow N} &\simeq (\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{D}_M} \rho_! \Omega_M) \otimes_{\rho_! \mathcal{D}_M} (\rho_! (\mathcal{D}_M \otimes_{\mathcal{A}_M} \bigwedge^\bullet \Theta_M \otimes_{f^{-1} \mathcal{A}_N} f^{-1} \mathcal{D}_N)) \\ &\simeq \mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! (\Omega_M \otimes_{\mathcal{A}_M} \bigwedge^\bullet \Theta_M \otimes_{f^{-1} \mathcal{A}_N} f^{-1} \mathcal{D}_N) \\ &\simeq \mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! (\Omega_M^\bullet \otimes_{f^{-1} \mathcal{A}_N} f^{-1} \mathcal{D}_N)[d_M]. \end{aligned}$$

Applying $Rf_{!!}$ we obtain:

$$\begin{aligned}
Rf_{!!}(\mathcal{D}b_M^{t\vee} \otimes_{\rho_! \mathcal{D}_M} \rho_! \mathcal{D}_{M \rightarrow N}) &\simeq Rf_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_!(\Omega_M^\bullet \otimes_{f^{-1} \mathcal{A}_N} f^{-1} \mathcal{D}_N))[d_M] \\
&\simeq Rf_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! \Omega_M^\bullet) \otimes_{\rho_! \mathcal{A}_N} \rho_! \mathcal{D}_N[d_M] \\
&\simeq f_{!!}(\mathcal{D}b_M^t \otimes_{\rho_! \mathcal{A}_M} \rho_! \Omega_M^\bullet) \otimes_{\rho_! \mathcal{A}_N} \rho_! \mathcal{D}_N[d_M] \\
&\rightarrow \mathcal{D}b_N^t \otimes_{\rho_! \mathcal{A}_N} \rho_! \Omega_N^\bullet \otimes_{\rho_! \mathcal{A}_N} \rho_! \mathcal{D}_N[d_N] \\
&\simeq \mathcal{D}b_N^t \otimes_{\rho_! \mathcal{A}_N} \rho_! \Omega_N \\
&= \mathcal{D}b_N^{t\vee},
\end{aligned}$$

where the third isomorphism follows from Lemma A.3.1 and the morphism from Lemma A.3.2. $\square$

By adjunction we get a morphism

$$(A.2) \quad \mathcal{D}b_M^{t\vee} \otimes_{\rho_! \mathcal{D}_M}^L \rho_! \mathcal{D}_{M \rightarrow N} \rightarrow f^! \mathcal{D}b_N^{t\vee}.$$

Theorem A.3.4 *The morphism (A.2) is an isomorphism.*

Proof. Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$ with compact support. We have the chain of isomorphisms

$$\begin{aligned}
\mathrm{RHom}(F, f^! \mathcal{D}b_N^{t\vee}) &\simeq \mathrm{RHom}(Rf_{!!} F, \mathcal{D}b_N^{t\vee}) \\
&\simeq \mathrm{R}\Gamma(N, T\mathcal{H}om(Rf_! F, \mathcal{D}b_N^\vee)) \\
&\simeq \mathrm{R}\Gamma(N, Rf_!(T\mathcal{H}om(F, \mathcal{D}b_M^\vee) \otimes_{\mathcal{D}_M}^L \mathcal{D}_{M \rightarrow N})) \\
&\simeq \mathrm{R}\Gamma(M, T\mathcal{H}om(F, \mathcal{D}b_M^\vee) \otimes_{\mathcal{D}_M}^L \mathcal{D}_{M \rightarrow N}) \\
&\simeq \mathrm{RHom}(F, \mathcal{D}b_M^{t\vee} \otimes_{\rho_! \mathcal{D}_M}^L \rho_! \mathcal{D}_{M \rightarrow N}),
\end{aligned}$$

where the third isomorphism follows from Theorem 4.4 of [13]. $\square$

By the equivalence between left and right $\mathcal{D}$ -modules, we have an isomorphism

$$(A.3) \quad \rho_! \mathcal{D}_{N \leftarrow M} \otimes_{\rho_! \mathcal{D}_M}^L \mathcal{D}b_M^t \xrightarrow{\sim} f^! \mathcal{D}b_N^t.$$

Corollary A.3.5 *When f is smooth we have an isomorphism*

$$f^{-1}\mathcal{D}b_N^t \xrightarrow{\sim} R\mathcal{H}om_{\rho_!\mathcal{D}_M}(\rho_!\mathcal{D}_{M \rightarrow N}, \mathcal{D}b_M^t).$$

Proof. The result is obtained by the following isomorphisms

$$\begin{aligned} R\mathcal{H}om_{\rho\mathcal{D}_M}(\rho_!\mathcal{D}_{M \rightarrow N}, \mathcal{D}b_M^t) &\simeq \rho_! R\mathcal{H}om(\mathcal{D}_{M \rightarrow N}, \mathcal{D}_M) \overset{L}{\underset{\rho_!\mathcal{D}_M}{\otimes}} \mathcal{D}b_M^t \\ &\simeq \rho_!\mathcal{D}_{N \leftarrow M} \overset{L}{\underset{\rho_!\mathcal{D}_M}{\otimes}} \mathcal{D}b_M^t[d] \\ &\simeq f^!\mathcal{D}b_N^t[d] \\ &\simeq f^{-1}\mathcal{D}b_N^t. \end{aligned}$$

The first isomorphism is obtained by replacing $\mathcal{D}_{M \rightarrow N}$ with its Koszul complex. The second follows from the smoothness of f and the isomorphism

$$R\mathcal{H}om_{\mathcal{D}_M}(\mathcal{D}_{M \rightarrow N}, \mathcal{D}_M) \simeq \mathcal{D}_{N \leftarrow M}[d].$$

The last isomorphism follows since when f is smooth we have the isomorphism $f^!(\cdot)[d] \simeq f^{-1}(\cdot)$. □

From now on X will be a complex manifold, with structure sheaf $\mathcal{O}_X$. We denote by $\overline{X}$ the complex conjugate manifold (with structure sheaf $\mathcal{O}_{\overline{X}}$), and $X_{\mathbb{R}}$ the underlying real analytic manifold, identified with the diagonal of $X \times \overline{X}$. Let $\mathcal{O}_X^t$ be the sheaf of tempered holomorphic functions on X . We also consider the sheaf $\Omega_X^t \in D^b(\rho_!\mathcal{D}_X^{op})$:

$$\Omega_X^t := \mathcal{D}b_{X_{\mathbb{R}}}^{t\vee} \overset{L}{\underset{\rho_!\mathcal{D}_{\overline{X}}}{\otimes}} \rho_!\mathcal{O}_{\overline{X}}[-d_X].$$

Proposition A.3.6 *Let $f : X \rightarrow Y$ be a holomorphic map between complex manifolds. Then*

$$(A.4) \quad \Omega_X^t \overset{L}{\underset{\rho_!\mathcal{D}_{\overline{X}}}{\otimes}} \rho_!\mathcal{D}_{X \rightarrow Y}[d_X] \simeq f^!\Omega_Y^t[d_Y].$$

Proof. We have the chain of isomorphisms

$$\begin{aligned}
f^!(\mathcal{D}b_{Y_{\mathbb{R}}}^{t\vee} \otimes_{\rho_! \mathcal{D}_{\overline{X}}}^L \rho_! \mathcal{O}_{\overline{Y}}) &\simeq f^! \mathcal{D}b_{Y_{\mathbb{R}}}^{t\vee} \otimes_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}^L \rho_! f^{-1} \mathcal{O}_{\overline{Y}} \\
&\simeq \mathcal{D}b_{X_{\mathbb{R}}}^{t\vee} \otimes_{\rho_! \mathcal{D}_{X_{\mathbb{R}}}}^L \rho_! \mathcal{D}_{X_{\mathbb{R}} \rightarrow Y_{\mathbb{R}}} \otimes_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}^L \rho_! f^{-1} \mathcal{O}_{\overline{Y}} \\
&\simeq (\mathcal{D}b_{X_{\mathbb{R}}}^{t\vee} \otimes_{\rho_! \mathcal{D}_X}^L \rho_! \mathcal{D}_{X \rightarrow Y}) \otimes_{\rho_! \mathcal{D}_{\overline{X}}}^L \rho_! \mathcal{D}_{\overline{X} \rightarrow \overline{Y}} \otimes_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}^L \rho_! f^{-1} \mathcal{O}_{\overline{Y}} \\
&\simeq (\mathcal{D}b_{X_{\mathbb{R}}}^{t\vee} \otimes_{\rho_! \mathcal{D}_X}^L \rho_! \mathcal{D}_{X \rightarrow Y}) \otimes_{\rho_! \mathcal{D}_{\overline{X}}}^L \rho_! \mathcal{O}_{\overline{X}} \\
&\simeq (\mathcal{D}b_{X_{\mathbb{R}}}^{t\vee} \otimes_{\rho_! \mathcal{D}_{\overline{X}}}^L \rho_! \mathcal{O}_{\overline{X}}) \otimes_{\rho_! \mathcal{D}_X}^L \rho_! \mathcal{D}_{X \rightarrow Y},
\end{aligned}$$

where the second isomorphism follows from Proposition A.3.4. $\square$

By the equivalence between left and right $\mathcal{D}$ -modules, we have an isomorphism

$$(A.5) \quad \rho_! \mathcal{D}_{Y \leftarrow X} \otimes_{\rho_! \mathcal{D}_X}^L \mathcal{O}_X^t \xrightarrow{\sim} f^! \mathcal{O}_Y^t.$$

Corollary A.3.7 *When f is smooth we have an isomorphism*

$$f^{-1} \mathcal{O}_Y^t \xrightarrow{\sim} R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^t).$$

Proof. The proof is similar to that of Corollary A.3.5. $\square$

A.4 Inverse image for Whitney holomorphic functions

Let $f : M \rightarrow N$ be a morphism of oriented real analytic manifolds of dimension d_M and d_N . Set $d = d_N - d_M$.

Lemma A.4.1 *The sheaf $f^! \mathcal{C}_N^{\infty, w}[d]$ is concentrated in degree zero.*

Proof. If f is smooth, then $f^!(\cdot)[d] \simeq f^{-1}$, and the result is clear. Let f be a closed embedding. Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$. We have the chain of isomorphisms

$$\begin{aligned}
R\mathcal{H}om(D'F; f^! \mathcal{C}_N^{\infty, w}[d]) &\simeq R\mathcal{H}om(f_! D'F[-d]; \mathcal{C}_N^{\infty, w}) \\
&\simeq R\mathcal{H}om(D'(f_! F); \mathcal{C}_N^{\infty, w}) \\
&\simeq R\Gamma(Y; f_! F \overset{w}{\otimes} \mathcal{C}_N^{\infty}).
\end{aligned}$$

The fourth isomorphism follows since $Rf_*DF \simeq D(Rf_!F)$ if $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$ and $Rf_* \simeq Rf_! \simeq f_!$ since f is a closed embedding. Let $U \in \text{Op}^c(M_{sa})$ be locally cohomologically trivial. We have $D'\mathbb{C}_{\overline{U}} \simeq \mathbb{C}_U$, and we get

$$R^{k+d}\Gamma(U; f^!\mathcal{C}_N^{\infty, w}) \simeq R^k\Gamma(X; f_!\mathbb{C}_{\overline{U}} \overset{w}{\otimes} \mathcal{C}_N^{\infty}) = 0$$

if $k \neq 0$ since $f_!\mathbb{C}_{\overline{U}} \overset{w}{\otimes} \mathcal{C}_N^{\infty}$ is soft. Hence $f^!\mathcal{C}_N^{\infty, w}[d]$ is concentrated in degree zero on a basis for the topology of M_{sa} and the result follows. $\square$

Proposition A.4.2 *There is a natural morphism in $\text{Mod}(\mathbb{C}_{M_{sa}})$*

$$f^!\mathcal{C}_N^{\infty, w}[d] \rightarrow \mathcal{C}_M^{\infty, w}$$

Proof. Let $U \in \text{Op}^c(M_{sa})$ be locally cohomologically trivial. We have the chain of morphisms

$$\begin{aligned} \Gamma(U; f^!\mathcal{C}_N^{\infty, w}[d]) &\simeq R\Gamma(N; Rf_!\mathbb{C}_{\overline{U}} \overset{w}{\otimes} \mathcal{C}_N^{\infty}) \\ &\rightarrow R\Gamma(M; f^{-1}Rf_!\mathbb{C}_{\overline{U}} \overset{w}{\otimes} \mathcal{C}_M^{\infty}) \\ &\rightarrow R\Gamma(M; \mathbb{C}_{\overline{U}} \overset{w}{\otimes} \mathcal{C}_M^{\infty}) \\ &\simeq \Gamma(U; \mathcal{C}_M^{\infty, w}), \end{aligned}$$

where the first isomorphism has been proved in Lemma A.4.1 and the first arrow follows from Theorem 3.3 of [13]. $\square$

Proposition A.4.3 *There is a natural morphism in $D^b(\rho_!\mathcal{D}_M)$:*

$$(A.6) \quad \rho_!\mathcal{D}_{M \rightarrow N} \overset{L}{\underset{\rho_!f^{-1}\mathcal{D}_M}{\otimes}} f^!\mathcal{C}_N^{\infty, w}[d] \rightarrow \mathcal{C}_M^{\infty, w}.$$

Proof. The Spencer resolution of $\mathcal{D}_{M \rightarrow N}$ gives rise to the quasi-isomorphism

$$\mathcal{D}_{M \rightarrow N} \overset{\sim}{\leftarrow} \mathcal{D}_M \otimes_{\mathcal{A}_M} \bigwedge^{\bullet} \Theta_M \otimes_{\mathcal{A}_M} \mathcal{D}_{M \rightarrow N} \simeq \mathcal{D}_M \otimes_{\mathcal{A}_M} \bigwedge^{\bullet} \Theta_M \otimes_{f^{-1}\mathcal{A}_M} f^{-1}\mathcal{D}_M$$

from which we obtain

$$\begin{aligned}
& \rho_! \mathcal{D}_{M \rightarrow N} \bigotimes_{\rho_! f^{-1} \mathcal{D}_M}^L f^! \mathcal{C}_N^{\infty, w}[d] \\
& \simeq \rho_! \mathcal{D}_M \bigotimes_{\rho_! \mathcal{A}_M} \rho_! \bigwedge^{\bullet} \Theta_M \bigotimes_{\rho_! f^{-1} \mathcal{A}_M} \rho_! f^{-1} \mathcal{D}_M \bigotimes_{\rho_! f^{-1} \mathcal{D}_M}^L f^! \mathcal{C}_N^{\infty, w}[d] \\
& \simeq \rho_! \mathcal{D}_M \bigotimes_{\rho_! \mathcal{A}_M} \rho_! \bigwedge^{\bullet} \Theta_M \bigotimes_{\rho_! f^{-1} \mathcal{A}_M} f^! \mathcal{C}_N^{\infty, w}[d] \\
& \rightarrow \rho_! \mathcal{D}_M \bigotimes_{\rho_! \mathcal{A}_M} \rho_! \bigwedge^{\bullet} \Theta_M \bigotimes_{\rho_! f^{-1} \mathcal{A}_M} \mathcal{C}_M^{\infty, w} \\
& \simeq \mathcal{C}_M^{\infty, w}.
\end{aligned}$$

□

By adjunction we get a morphism

$$(A.7) \quad f^! \mathcal{C}_N^{\infty, w}[d] \rightarrow R\mathcal{H}om_{\rho_! \mathcal{D}_M}(\rho_! \mathcal{D}_{M \rightarrow N}, \mathcal{C}_M^{\infty, w}).$$

Theorem A.4.4 *The morphism (A.7) is an isomorphism.*

Proof. Let $F \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_M)$. We have the chain of isomorphisms

$$\begin{aligned}
R\mathcal{H}om(D'F, f^! \mathcal{C}_N^{\infty, w}[d]) & \simeq R\Gamma(Y; Rf_! F \overset{w}{\otimes} \mathcal{C}_N^{\infty}) \\
& \simeq R\mathcal{H}om_{\mathcal{D}_M}(\mathcal{D}_{M \rightarrow N}, F \overset{w}{\otimes} \mathcal{C}_M^{\infty}) \\
& \simeq R\mathcal{H}om_{\mathcal{D}_M}(\mathcal{D}_{M \rightarrow N}, \rho^{-1} R\mathcal{H}om(D'F, \mathcal{C}_M^{\infty, w})) \\
& \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_M}(\rho_! \mathcal{D}_{M \rightarrow N}, R\mathcal{H}om(D'F, \mathcal{C}_M^{\infty, w})) \\
& \simeq R\mathcal{H}om(D'F, R\mathcal{H}om_{\rho_! \mathcal{D}_M}(\rho_! \mathcal{D}_{M \rightarrow N}, \mathcal{C}_M^{\infty, w})),
\end{aligned}$$

where the second isomorphism follows from Theorem 3.5 of [13].

□

Corollary A.4.5 *When f is smooth we have an isomorphism*

$$f^{-1} \mathcal{C}_N^{\infty, w} \xrightarrow{\sim} R\mathcal{H}om_{\rho_! \mathcal{D}_M}(\rho_! \mathcal{D}_{M \rightarrow N}, \mathcal{C}_M^{\infty, w}).$$

Proof. It follows from the fact that $f^!(\cdot)[d] \simeq f^{-1}$ when f is smooth.

□

From now on X will be a complex manifold of complex dimension d_X , with structure sheaf $\mathcal{O}_X$. We denote by $\overline{X}$ the complex conjugate manifold (with structure sheaf $\mathcal{O}_{\overline{X}}$), and $X_{\mathbb{R}}$ the underlying real analytic manifold, identified with the diagonal of $X \times \overline{X}$. Let $\mathcal{O}_X^w$ be the sheaf of Whitney holomorphic functions on X .

Theorem A.4.6 *Let $f : X \rightarrow Y$ be a morphism of complex manifolds. Then*

$$(A.8) \quad f^! \mathcal{O}_Y^w[2d_Y] \xrightarrow{\sim} R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w)[2d_X].$$

Proof. Remark that, if $\mathcal{M} \in D^b(\rho_! \mathcal{D}_{X_{\mathbb{R}}})$ we have

$$\begin{aligned} & R\mathcal{H}om_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}(\rho_! f^{-1} \mathcal{O}_{\overline{Y}}, R\mathcal{H}om_{\rho_! \mathcal{D}_{X_{\mathbb{R}}}}(\rho_! \mathcal{D}_{X_{\mathbb{R}} \rightarrow Y_{\mathbb{R}}}, \mathcal{M})) \\ & \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, R\mathcal{H}om_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}(\rho_! f^{-1} \mathcal{O}_{\overline{Y}}, R\mathcal{H}om_{\mathcal{D}_{\overline{X}}}(\rho_! \mathcal{D}_{\overline{X} \rightarrow \overline{Y}}, \mathcal{M}))) \\ & \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! (\mathcal{D}_{\overline{X} \rightarrow \overline{Y}} \overset{L}{\otimes}_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}} f^{-1} \mathcal{O}_{\overline{Y}}), \mathcal{M})) \\ & \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mathcal{M})). \end{aligned}$$

We have the chain of isomorphisms

$$\begin{aligned} f^! \mathcal{O}_Y^w[2d_Y] & \simeq f^! R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{Y}}}(\rho_! \mathcal{O}_{\overline{Y}}, \mathcal{C}_{Y_{\mathbb{R}}}^{\infty, w})[2d_Y] \\ & \simeq R\mathcal{H}om_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}(f^{-1} \rho_! \mathcal{O}_{\overline{Y}}, f^! \mathcal{C}_{Y_{\mathbb{R}}}^{\infty, w})[2d_Y] \\ & \simeq R\mathcal{H}om_{\rho_! f^{-1} \mathcal{D}_{\overline{Y}}}(\rho_! f^{-1} \mathcal{O}_{\overline{Y}}, R\mathcal{H}om_{\rho_! \mathcal{D}_{X_{\mathbb{R}}}}(\rho_! \mathcal{D}_{X_{\mathbb{R}} \rightarrow Y_{\mathbb{R}}}, \mathcal{C}_{X_{\mathbb{R}}}^{\infty, w}))[2d_X] \\ & \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X}}}(\rho_! \mathcal{O}_{\overline{X}}, \mathcal{C}_{X_{\mathbb{R}}}^{\infty, w}))[2d_X] \\ & \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w)[2d_X]. \end{aligned}$$

□

Corollary A.4.7 *When f is smooth we have an isomorphism*

$$f^{-1} \mathcal{O}_Y^w \xrightarrow{\sim} R\mathcal{H}om_{\rho_! \mathcal{D}_X}(\rho_! \mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^w).$$

Proof. The proof is similar to that of Corollary A.4.5.

□

References

- [1] E. Andronikof, Microlocalisation tempérée, *Mémoires Soc. Math. France* **57** (1994).
- [2] A. Beck, Continuous flows in the plane, *Grundlehren der Math.* **201**, Springer-Verlag, New York-Heidelberg (1974).
- [3] E. Bierstone, D. Milmann, Semianalytic and subanalytic sets, *Publ. I.H.E.S.* **67** pp. 5-42 (1988).
- [4] M. Coste, An introduction to o-minimal geometry, *Dip. Mat. Univ. Pisa, Dottorato di Ricerca in Matematica, Istituti Editoriali e Poligrafici Internazionali*, Pisa (2000).
- [5] V. Colin, Formal microlocalization, *C. R. Acad. Sci. Paris Math.* **327**, pp. 289-293 (1998).
- [6] V. Colin, Specialization of formal cohomology and asymptotic expansions, *Publ. RIMS, Kyoto Univ.* **37** pp. 37-69 (2001).
- [7] A. D'Agnolo, P. Schapira, Leray's quantization of projective duality, *Duke Math. Journal* **84** pp. 453-496 (1996).
- [8] M. Edmundo, G. Jones and N. Peatfield, Sheaf cohomology in o-minimal structures, *J. Math. Logic* **6** pp. 163-179 (2006).
- [9] M. Kashiwara, The Riemann-Hilbert problem for holonomic systems, *Publ. RIMS, Kyoto Univ.* **20** pp. 319-365 (1984).
- [10] M. Kashiwara, $\mathcal{D}$ -modules and microlocal calculus, *Translations of Math. Monog.* **217**, Iwanami Series in Modern Math., American Math. Soc., Providence (2003).
- [11] M. Kashiwara, P. Schapira, Sheaves on manifolds, *Grundlehren der Math.* **292**, Springer-Verlag, Berlin (1990).
- [12] M. Kashiwara, P. Schapira, Categories and sheaves, *Grundlehren der Math.* **332**, Springer-Verlag, Berlin (2006).
- [13] M. Kashiwara, P. Schapira, Moderate and formal cohomology associated with constructible sheaves, *Mémoires Soc. Math. France* **64** (1996).
- [14] M. Kashiwara, P. Schapira, Ind-sheaves, *Astérisque* **271** (2001).

- [15] M. Kashiwara, P. Schapira, Microlocal study of ind-sheaves I: micro-support and regularity, *Astérisque* **284**, pp. 143-164 (2003).
- [16] M. Kashiwara, P. Schapira, F. Ivorra, I. Waschkes, Microlocalization of ind-sheaves, *Studies in Lie theory, Progress in Math.* **243**, Birkhäuser pp. 171-221 (2006).
- [17] S. Łojaciewicz, Sur le problème de la division, *Studia Mathematica* **8** pp. 87-136 (1959).
- [18] S. Łojaciewicz, Sur la géométrie semi- et sous-analytique, *Ann. Inst. Fourier* **43** pp. 1575-1595 (1993).
- [19] B. Malgrange, Ideals of differentiable functions, Tata Institute, Oxford University Press (1967).
- [20] B. Malgrange, Équations différentielles à coefficients polynomiaux, *Progress in Math.* **96**, Birkhäuser (1991).
- [21] A. R. Martins, Functorial properties of the microsupport and regularity for ind-sheaves, *Math. Zeitschrift* Vol. **260** N. **3**, pp. 541-556 (2008).
- [22] L. Prelli, Sheaves on subanalytic sites, PHD thesis, Universities of Padova and Paris 6 (2006).
- [23] L. Prelli, Sheaves on subanalytic sites, *Rend. Sem. Mat. Univ. Padova* Vol. **120**, pp. 167-216 (2008).
- [24] L. Prelli, Microlocalization of subanalytic sheaves, *C. R. Acad. Sci. Paris Math.* **345**, pp. 127-132 (2007).
- [25] L. Prelli, Cauchy-Kowaleskaya-Kashiwara theorem with growth conditions, *Math. Zeitschrift* Vol. **265** N. **1**, pp. 115-124 (2010).
- [26] P. Schapira, Microdifferential systems in the complex domain, *Grundlehren der Math.* **269**, Springer-Verlag, Berlin (1985).
- [27] P. Schapira, J. P. Schneiders, Index theorem for elliptic pairs, *Astérisque* **224** (1994).
- [28] J. P. Schneiders, An introduction to $\mathcal{D}$ -modules, *Bull. Soc. Royale des Sciences de Liège* **63**, pp. 223-295 (1994).
- [29] Y. Sibuya, Linear ordinary differential equation in the complex domain: problems of analytic continuation, *Transl. of Math. Monog.*, **82**, AMS, Providence (1990).

- [30] G. Tamme, Introduction to étale cohomology, Universitext, Springer-Verlag, Berlin (1994).
- [31] L. Van Den Dries, Tame topology and o-minimal structures, London Math. Society Lecture Notes Series **248**, Cambridge University Press, Cambridge (1998)
- [32] L. Van Den Dries, C. Miller, Geometric categories and o-minimal structures, Duke Math. Journal **84** vol. **2** pp. 497-540 (1996).
- [33] I. Waschies, Microlocal perverse sheaves, Bull. Soc. Math. de France **132** vol. **3** pp. 397-462 (2004).
- [34] A. J. Wilkie, Covering definable open sets by open cells. In M. Edmundo, D. Richardson, and A. Wilkie, editors, O-minimal Structures, Proceedings of the RAAG Summer School Lisbon 2003, Lecture Notes in Real Algebraic and Analytic Geometry, Cuvillier Verlag (2005).
- [35] SGA4: Sém. Géom. Algébrique du Bois-Marie by M. Artin, A. Grothendieck, J. L. Verdier, Lecture Notes in Math. **269**, Springer-Verlag, Berlin (1972).